

ASSEMBLAGE AND DISTRIBUTION OF EARTHWORMS IN THE TALACAUVERY WILDLIFE SANCTUARY, WESTERN GHATS

M.P. Krishna*

*Department of Zoology, Field Marshal K. M. Cariappa Mangalore University College,
Madikeri 571201, Karnataka, India.

mcmpkrishna@gmail.com

ABSTRACT

Earthworms have the potential to biodegrade and biotransform chemical pollutants, there by converting them to less toxic substances in their bodies. Earthworms play an important role in soil ecology as they have the tendency to reach high densities. They are recognized as soil engineers, and occupy a vital position in soil food-webs. They improve the structure of the soil by tilting and mixing, Earthworms play important role in humus formation and increasing the water holding capacity of the soil. The distribution of earthworms is usually diverse and their numbers fluctuate in relation to the different abiotic factors and land use patterns of the soil. The fertility of soil depends on the biological diversity and soil faunal biomass. The present study was carried out in Talacauvery wildlife sanctuary of the Central Western Ghats. It is spread over an area of about 105sq km and is present at 1525m above sea level. Temperature in the region varies from 10° – 28° C. Annual rainfall ranges between 6500mm – 7500mm. The sampling was conducted during the months of May to October 2017. Earthworms were collected from selected site two to four quadrats were randomly laid, earthworms were sampled from different land use pattern like agricultural land, forest, gardens and nurseries by the hand-sorting method up to 30 cm deep by using quadrates (30 × 30 cm² area). In the present study twelve earthworm species have been recorded belonging to nine genera and five families. As per the ecological category, one species anecic two species are epigeic and nine endogeic species are identified. Out of these Family Megascolecidae six species, Moniligastridae three species, Family Glossoscolecidae, Acanthodrilidae and Eudrilidae one each species were recorded.

Keywords: Assemblage, Biodegrade, Distribution, Earthworms, Wildlife Sanctuary

INTRODUCTION

Talacauvery wildlife sanctuary of the Central Western Ghats is spread over an area of about 105sq km and is present at 1525m above sea level. Temperature in the region varies from 10° – 28° C. Annual rainfall ranges between 6500mm – 7500mm. It is a catchment area for Cauvery River and serves as an important catchment for perennial streams like Dodda Hole, Nadumale Hole, Bettamale Hole, Kume Kolli and

Mundra Hole. The hills and valleys in the area are dense evergreen forests 63% of the evergreen trees reported are endemic. The diversity of forests provides varied habitats for a host of mammals, birds, reptiles and amphibians. And is threatened by increasing anthropogenic pressure on its natural resources. Soil biodiversity is key to sustainable organic farming (Ramakrishnan et al., 2005) and earth worms are the most dominant component of soil biota in terms of biomass and crucial for maintaining soil fertility (Dash, 1978; Senapati and Dash, 1981; Julka and Paliwal, 2005a and b; Dash et al, 2009; Bhadauria et al., 2012; Dash, 2012). Western Ghats region of India, the areas distinguished globally for their highly valuable biodiversity and ecosystem services. On the basis of available data, the Western Ghats and West Coastal plains would stand out as the region with the highest level of earthworm species richness. The Western Ghats region is home to 53% species known from India. The Western Ghats harbour 193 native species. The distribution of earthworms is usually diverse and their numbers fluctuate in relation to the different abiotic factors and land use patterns of the soil. The fertility of soil depends on the biological diversity and soil faunal biomass. Earthworms are known to be the most important soil fauna biomass in humid soils of temperate and tropical regions (Lee, 1985). The beneficial role of earthworms in the breakdown of dead plant material in the forest litter was first documented by (Darwin, 1881). For a long time, earthworms have been known as the farmer's friend, natural ploughmen, soil ecosystem engineers and intestines of earth. Earthworms can significantly influence soil physical, chemical and biological properties, hence improving the fertility and structure of soil (Doan et al., 2013; Singh et al., 2016). Earthworms also play an important role in mixing of mineral soils and plant materials. Various studies reported that the disturbance and degradation of natural forest affect the number of earthworms and their distribution (Baretta et al., 2007; Chandran et al., 2012). The distribution of earthworm is usually heterogeneous (Guild, 1952; Satchell, 1955; Svendsen, 1957) and their numbers fluctuate in relation to the abiotic factors of the soil (Evans and Guild, 1948). Environmental factors like moisture, temperature, pH and soil texture also affect the distribution of earthworms. (Bhadauria and Ramakrishnan 1989) determined that regional earthworm biodiversity and species dispersal pattern was influenced by a variety of biotic and abiotic forces such as soil properties, surface litter inputs, surface vegetation type, dynamic land management history, local or regional climate and human pressure. The significance of diverse soil habitats is one of the most influencing factors affecting the overall earthworm distribution (Rajkhowa et al., 2014). Changes in land use patterns have also directly affected the composition and population structure of earthworm communities in different agro-climatic regions (Blanchart and Julka, 1997; Behera et al., 1999; Bhadauria et al., 2000; Lalthanzara et al. 2011). Endogeic earthworm appears a key feature of soil functioning in the urban context through their roles on organic matter transformation, the formation and maintenance of soil structure (Amosse et al., 2015). The Western Ghats have 219 (193 native, 26 exotic) documented species (Siddaraju et al., 2013), which is about 52.4% of total numbers in India, followed by the Eastern Himalayas and Northeast Hills Region with 85 native and 25 exotic species (Julka, 2010). Therefore, current study of earthworm diversity and distribution in Talacauvery Wildlife Sanctuary, Western Ghats were carried out.

MATERIALS AND METHODS

The present study was carried out in Talacauvery wildlife sanctuary Western Ghats (**Fig. 1**). The sampling was conducted during the months of May to October 2017. Earthworms were collected from selected site two to four quadrats were randomly laid, earthworms were sampled from different land use pattern like agricultural land, forest, gardens and nurseries by the hand-sorting method up to 30 cm deep by using quadrates (30 × 30 cm² area). A global positioning system (GPS) was also used to mark the latitude and longitude of each sampling site. Moisture content was measured with a digital soil moisture meter (Micro make). The collected samples of earthworms with appropriate amount of soil were placed in polythene bags labelled with place name, date of collection, surrounding soil biota etc. and brought to the research lab for further study. Earthworms were washed in fresh water and sorted on the presence or absence of clitellum. Earthworms were narcotized in 70% ethyl alcohol and fixed in 5% formalin for 6-8 hours and finally preserved in 5% formalin. The preserved samples of earthworms were identified upto the species level by Dr (Mrs) P. Kathireswari, an earthworm taxonomist and also by referring the descriptions and keys of identification of earthworms published by (Stephenson 1915, 1923, Michaelsen 1910 and Julka 1988). The identified earthworms were deposited in the Museum of Zoology Department, Field Marshall K. M. Cariappa College, Madikeri.

RESULTS AND DISCUSSION

In the present study twelve earthworm species have recorded (**Table. 1**) belonging to nine genera and five families. As per the ecological category, one species anecic two species are epigeic and nine endogeic species are identified. Out of these Family Megascolecidae six species, Moniligastridae three species, Family Glossoscolecidae, Acanthodrilidae and Eudrilidae one each species were recorded. A great variety of vegetation types coupled with high rainfall and moderate temperature provide many different niches for the occurrence of earthworms. The Western Ghats and west coast of the country, is considered as the richest in terms of earthworm biodiversity (Julka and Paliwal 2005). The overall findings showed that Talacauvery wildlife sanctuary, Western Ghats had higher number of native earthworm species. This is an indication that their habitat is less disturbed, habitat alteration and disturbance leads to the invasion of exotic species (Bhadauria and Saxena 2007). These results suggest that habitat plays a major role in the earthworm species diversity. Total number of species identified in this study showed a decreasing tendency at higher altitude, this observation coincides with the earlier reports on earthworm diversity (Decaens, 2010). Many studies shown that the same species occupy different altitudes, for example *Drawida ghatensis* was reported at an altitude 1,524-2438 m by (Stephenson 1923) and the same has been observed by (Julka and Chandra 1986) at 940-1,40 m in silent valley. On the contrary, the exotic species can tolerate a wide range of soil and environmental conditions. Therefore, any kind of disturbances to the natural ecosystems and vegetation of an area may lead to the elimination of native earthworm species and subsequent invasion of exotic species (Bhadauria and

Ramakrishnan 1991; Fragoso et al., 1997). The successful colonisation of the exotic earthworm fauna is mainly due to their inherent ability to withstand disturbance and interference system (Julka 1988).




Fig. 1 Talacauvery Wildlife Sanctuary Western Ghats

Sl no.	Family	Species	Exotic/ Native	Ecological category
1	Glossoscolecidae	<i>Pontoscolex corethrurus</i>	Exotic	Endogeic
2	Acanthodrilidae	<i>Plutellus variabilis</i>	Native	Epigeic
3	Megascolecidae	<i>Lampito mauritii</i>	Native	Anecic
		<i>Amyntas corticis</i>	Exotic	Endogeic
		<i>Megascolex konkanensis</i>	Native	Endogeic
		<i>Megascolex trarancorensis</i>	Native	Endogeic
		<i>Notoscolex tenmalai</i>	Native	Endogeic
		<i>Metaphire houlleti</i>	Native	Endogeic
4	Moniligastridae	<i>Drawida pellucida</i>	Native	Endogeic
		<i>Drawida travancorensis</i>	Native	Endogeic
		<i>Drawida ghatensis</i>	Native	Endogeic
5	Eudrilidae	<i>Eudrilus eugeniae</i>	Exotic	Epigeic

Table: 1 List of families and exotic/native species of earthworms sampled from Talacauvery Wildlife sanctuary, together with their ecological categories

ACKNOWLEDGMENTS

I would like to record my sincere thanks to Karnataka Forest Department, Kodagu district, for granting permission to carry out this study. I also thank the Principal, FMKMC College, Madikeri, for support, encouragement and facility provided. Special thanks to Dr. Siddaraju for the help during field visit, identification of earthworms and photography.

REFERENCES

- Amosse, J., Turberg, P., Kohler-Milleret, R., Gobat, J., Le Bayon, R., 2015. Effects of endogeic earthworms on the soil organic matter dynamics and the soil structure in urban and alluvial soil materials. *Geoderma* 243–244, 50–57.
- Baretta, D., Brown, G.G., James, S.W., Bran, E.J., Cardoso, N., 2007. Earthworm populations sampled using collection methods in Atlantic forests with *Araucaria angustifolia*. *Sci. Agric.* 64, 384–392.
- Behera, B. Giri, S., Dash, N. C., Sahu, J. and Senapati, B. K. 1999. Earthworm bioindication of forest land use pattern in western Orissa. *Indian Forester*. 125: 272-281.
- Bhadauria, T. and Ramakrishnan, P. S. 1989. Earthworm population dynamics and contribution to nutrient cycling during cropping and fallow phases of shifting agriculture (Jhum) in northeast India. *J. Applied Ecology*. 26: 505-520.
- Bhadauria T, Ramakrishnan PS 1991. Population dynamics of earthworms and their activity in forest ecosystems of north-east India. *J Trop Ecol* 7:305–318
- Bhadauria T, Ramakrishnan PS, Srivastva KN 2000. Diversity and distribution of endemic and exotic earthworm in natural regeneration ecosystems in the Central Himalaya, India. *Soil Biol Biochem* 32:2045–2054.
- Bhadauria T, Saxena, KG 2007. Influence of landscape modification on earthworm biodiversity in the Garhwal region of central Himalayas. *Proceedings of Indo-US workshop on Vermitechnology in human welfare, Coimbatore.* pp 80–95.
- Bhadauria, T., Kumar, P., Kumar, R., Maikhuri, R. K., Rao, K. S. and Saxena, K. G. 2012. Earthworm populations in a traditional village landscape in Central Himalaya, India. *Applied Soil Ecology*. 53: 83-93.
- Blanchart, E. and Julka, J.M. 1997. Influence of forest disturbance on earthworm (Oligochaeta) communities in the western ghats (South India). *Soil Biol. Biochem.*, 29(3/4): 303–6.
- Chandran, M.S.S., Sujatha, S., Mohan, M., Julka, J.M., Ramasamy, E.V., 2012. Earthworm diversity at Nilgiri biosphere reserve, Western Ghat, India. *Biodivers. Conserv.* 21, 3342–3353.
- Darwin, C., 1881. *The formation of vegetable mould through the action of worms, with observation on their habits* Murray, p. 326, London.
- Dash, M. C. 1978. The role of earthworms in the decomposer system, Pages 399-409, In: Singh, J. S. and Gopal, B. (Eds) *Glimpses of Ecology*, International Scientific Publishers, Jaipur, India.
- Dash, M. C., Saxena, K. G. and Giri, S. 2009. Vermitechnology for Wasteland Reclamation, Plant Productivity and Composting: A Review in Indian Context. *International J. Ecology and Environmental Sciences*. 35(2-3):163-185.
- Dash, M. C. 2012. *Charles Darwin's Plough. Tools for Vermitechnology*. I. K. International Publishing House Pvt. Ltd. New Delhi. p. 185.
- Decaens T 2010. Macroecological patterns in soil communities. *Global Ecol Biogeogr* 19:287–302.

- Doan, T.T., Ngo, P.T., Rumpel, C., Nguyen, B.V., Jouquet, P., 2013. Interactions between compost, vermicompost and earthworms influence plant growth and yield: a one year greenhouse experiment. *Sci. Hortic. Amsterdam* 160, 148–154.
- Evans, A.V. and Guild, W.J.M.C.L. 1948. Studies on the relationships between earthworms and soil fertility IV. On the life-cycle of some British Lumbricidae. *Ann. Appl. Biol.*, 35: 471–84.
- Fragoso C, Brown GG, Patro'n JC, Blanchart E, Lavelle P, Pashanasi P, Senapati B, Kumar T. 1997. Agricultural intensification, soil biodiversity and agroecosystem function in the tropics the role of earthworms. *Appl Soil Ecol* 6:17–35
- Guild, W. J. Mc.L 1952. Variations in earthworm numbers within field population. *J. Animal Ecology*. 21:169.
- Julka JM 1988. The fauna of Indian and the adjacent countries: Megadrile Oligochaeta (Earthworms): Haplotaxida: Lumbricina: Megascolecoidea: Octochaetidae. *Zoological Survey of India, Calcutta*, p 399.
- Julka JM. 2010. Role of earthworms in soil ecosystem. *Advances in Environmental Sciences—a resource material*, School of Environmental Sciences, Mahatma Gandhi University, Kottayam pp 80–85.
- Julka JM 1988. The fauna of Indian and the adjacent countries: Megadrile Oligochaeta (Earthworms): Haplotaxida: Lumbricina: Megascolecoidea: Octochaetidae. *Zoological Survey of India, Calcutta*, p 399.
- Julka JM, Paliwal R 2005. Distribution of earthworms in different agroclimatic regions of India. In: Ramakrishnan PS, Saxena KG, Swift M, Raoks J, Maikhuri RK (eds) *Soil biodiversity, ecological processes and landscape*. Oxford and ABH Publications Co. Pvt. Ltd., New Delhi, pp 3–13.
- Julka, J. M. and Paliwal, R. 2005a. Diversity and Biogeography of Indian Earthworms, In: proceedings of National Level Workshop on Vermitechnology Transfer to NSS programme officers, Department of Biochemistry and Zoology, Kongunadu Arts and Science College, Coimbatore, India. pp. 5-21.
- Julka, J. M. and Paliwal, R. 2005b. Distribution of earthworms in different agro-climatic regions of India. In: *Soil Biodiversity, Ecological Processes and Landscape Management*. Edited by P.S. Ramakrishnan, K.G. Saxena, M.J. Swift, K.S. Rao and R.K. Maikhuri. Oxford and IBH Publishing Company Private Limited, New Delhi. pp. 3-13.
- Julka JM, Chandra M 1986. On a small collection of earthworms and leeches (annelida) from the Silent Valley, Kerala, India. *Rec Zool Surv India* 84(1–4):165–171
- Lalthanzara, H., Ramanujam, S.N., Jha, L.K., 2011. Population dynamics of earthworm in relation to soil physico-chemical parameters in agroforestry systems if Mizoram, India. *J. Environ. Biol.* 32, 599–605.
- Lee, K.E., 1985. *Earthworms: Their Ecology and relationship with soil and land use*. Academic Press, Sydney, Australia. 444p.
- Michaelsen, W. 1910. Die oligochattem fauna der vorde indish-ceylonischen region. *Abh Geb Naturw Hamburg*, 19(5): 1–108.
- Rajkhowa, D.J., Bhattacharyya, P.N., Sarma, A.K., Mahanta, K., 2014. Diversity and distribution of Earthworms in different soil habitats of Assam, North-East India, an Indo-Burma Biodiversity Hotspots. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.*

- Ramakrishnan, P. S. Saxena, K. G., Swift, M. J., Rao, K. S. and Maikhuri, R. K. 2005. Soil Biodiversity, Ecological Processes and Landscape Management. Oxford and IBH Publishing Company Private Limited, New Delhi. p.302.
- Satchell, J.E., 1955. Some aspects of earthworm ecology. In: Kevan, D. K., Butterworths, Mc E. (Eds.), Soil Zoology, pp. 180–201, London.
- Senapati, B. K. and Dash, M. C.1981. Effect of grazing on the elements of production in vegetation and Oligochaeta of a tropical pasture. Review of Ecology and Biology of Soil. 18: 487-505.
- Siddaraju, M., Sreepada, K.S. and Krishna, M.P. 2013. Recorded distribution of earthworms of the family Octochaetidae in Dakshina Kannada district, south-west coast, Karnataka. *Int. J. Scientific Res. Publ.*, 3(6): 108–16.
- Singh, S.P., Singh, J., Vig, A.P., 2016. Earthworm as ecological engineers to change the physico-chemical properties of soil: Soil vs vermicast. *Ecol. Eng.* 90, 1–5.
- Stephenson J 1923. The fauna of British India including Ceylon and Burma, Oligochaeta. Taylor and Francis, London IV, p 518.
- Svendsen, J.A., 1957. The distribution of Lumbricidae in an area of Penine moorland (moor House, Nature reserve). *J. Anim. Ecol.* 26 (2), 411–421.

