

SOIL STRUCTURE ANALYSIS OF AGRICULTURE LAND OF VIJAPUR AND VISNAGAR TALUKAS OF MEHSANA DISTRICT, NORTHGUJARAT, INDIA

H.A. Patel

Assistant Professor
Physics Department

Sir P. T. Science College, Modasa, Gujarat, India

Abstract : Soil is important for growing of different crops. So It is necessary to study soil parameters. In present research work, we have made chemical parameters analysis. of agricultural soil of different villages of Vijapur and Visanagar Talukas of Mehsana district of North Gujarat. My primary focus is based on randomly selected 20 soil samples. Soil samples were collected by locally trained farmers and brought for soil test at Laboratory. Soil analysis is done by standard methods. The aim of this work is to study the fertility of soil of different agricultural land of Mehsana district by measuring soil parameters.

Key words: Soil Parameters , Agricultural , Analysis, Chemical

I. INTRODUCTION

For analysis of soil, it is necessary to know the fundamental needs of soil. Soil is a natural medium which provides water, nutrients, air and heat to the plants for its wholesome growth and give mechanical support to the plant. Soil is a reservoir of major nutrients required by the crops but it doesn't provide all the necessary nutrient immediately as requirement of plants. The aim of this analysis is to assess the surplus, adequacy or deficiency of available nutrients for growth of crop and to monitor change by farming of crop. Plant growth depends on fertility of soil and soil fertility is determined by the availability of nutrients either in the form of macro or micro.

The area of this study is agricultural land of different taluka of Mehsana district which is situated in North Gujarat in India. Its area is more than 4,500 km². The temperature variation is 45°C (max.) to 12°C (min.) and average rainfall is 800-900 mm.

Major soils are medium black, sandy and hydromorphic type. Major field crops are bajra, green gram, castor, groundnut, cotton, wheat, moth bean and major horticultural crops are mango, papaya, banana, cucurbits etc.

II. MATERIALS AND METHODS:

We have selected 20 soil samples from different agricultural land of Mehsana district for this study. The detail of the collected soil samples are shown in table 1.

TABLE NO-1 IDENTIFICATION OF SOIL SAMPLES

SAMPLE NO.	NAME OF VILLAGE	SAMPL E NO.	NAME OF VILLAGE	SAMPL E NO.	NAME OF VILLAGE	SAMPLE NO.	NAME OF VILLAGE
1	Devipura	6	Kada	11	Mandali	16	Ladol
2	Motipura	7	Kharod	12	Gundrasan	17	Manipur
3	Chadasana	8	Hirpura	13	Malosan	18	Pamol
4	Gerita	9	Sunshi	14	Tatosan	19	Rampura
5	Ubakhal	10	Ghagaret	15	Morvad	20	Kamana

III. TOOLS AND TECHNIQUES:

Mean, Maximum, Minimum and Standard Deviation are calculated from the measured soil parameters. Descriptive statistical analysis is used to analyze the data of soil samples. Variables like Phosphorus (P), Potassium (K), Electrical Conductivity (EC), Organic Carbon (OC) and pH are included in this study for analysis.

IV. SOIL SAMPLING AND ANALYSIS:

These samples were collected by a systematic sampling strategy at 0 to 15 cm depth below the surface of soil. The samples were dried and passed through a 2 mm sieve to prepare them for testing. All the samples were tested using standard Manual-Soil Testing Methods used in India.

Physical parameters like electrical conductivity (EC), phosphorus (P), potassium (K), Organic carbon (OC) and pH are measured for different collected soil samples.

Table 2: Soil characteristics of selected samples from study area

Sample No.	P	K	OC	pH	EC
1	42	262	0.65	7.7	0.28
2	38	248	0.66	7.3	0.27
3	42	252	0.57	7.4	0.32
4	46	270	0.73	6.9	0.42
5	37	232	0.75	8.5	0.33
6	30	242	0.66	8.1	0.55
7	35	268	0.55	8.2	0.42
8	46	291	0.35	7.6	0.45
9	47	281	0.64	8.7	0.47
10	52	244	0.47	7.0	0.36
11	48	256	0.48	7.3	0.28
12	45	266	0.72	8.1	0.38
13	42	264	0.68	7.6	0.41
14	37	257	0.54	7.9	0.47
15	37	245	0.59	8.2	0.52
16	32	278	0.65	7.3	0.38
17	36	232	0.63	7.3	0.37
18	38	271	0.74	7.6	0.44
19	52	263	0.71	7.8	0.43
20	48	283	0.52	7.2	0.53
Mean	41.5	260.2	0.58	7.69	0.40
Minimum	30	232	0.35	6.9	0.27
Maximum	52	291	0.76	8.7	0.55
Standard Deviation	6.37	16.49	0.11	0.49	0.083

TABLE – 3 GENERAL INTERPRETATION OF SOIL PARAMETERS

Parameters	pH	EC dS/cm	OC %	K Kg/ha	P Kg/ha	S
Interpretation	< 4.6 Extremely Acidic	0 – 2 Sault free	< 0.5 Low	< 108 Low	< 10 Low	8 – 10 (ppm) Critical limit
	4.6 – 5.5 Strongly Acidic	4– 8 Slightly Saline	0.5 – 0.75 Medium	108 –280 Medium	10 –24.6 Medium	
	5.6 – 6.5 Moderate Acidic	9-15 Moderate Saline	> 0.75 High	>280 High	>24.6 High	
	6.6 – 6.9 Slight Acidic	>15 Highly Saline				
	7 Neutral					
	7.1 -8.5 Moderate Alkaline					
	> 8.5 Strongly Alkaline					

V. THE RESULTS AND DISCUSSION:

Statistical analysis of parameters of soil samples are shown in table 2. The pH parameter of collected soil samples varies between the range 8.7 and 6.9 with the mean value of 7.69. As presented in table-2 and table -3, pH value indicates soils are neutral to alkaline in reaction. (6.9 -8.7).Majority samples are alkaline and 5% samples are neutral. No sample is acidic in nature (pH < 6.5). The Electrical conductivity (EC) is varied from 0.27 to 0.55 dScm⁻¹ with a mean value of 0.40 dScm⁻¹. The values of EC is in the range 0 - 2. This shows that all samples are salt free (ref: table 3). Organic carbon (OC) of the soil is varied from 0.35– 0.76 with a mean value of 0.58 as in table-2. It is very low (< 0.50) in 5% soil samples, 90% soil samples are medium and 5% samples are with high value (> 0.75). Phosphorus(P) parameter in the samples is 30–52 kg/ha.

It is observed that phosphorous in all the samples are of high range (> 24.6). It can be due to high fertilizer practice. In case of potassium no sample is in low range. Among 20 samples, 85% contains medium amount (108 – 280) whereas remaining 15% are with very high amount (> 280). It may be due to use of over fertilizers.

VI. CONCLUSION :

Study shows that the study area is free from salt, not even acidic in nature. It is moderately neutral in nature. Very high value of potassium and phosphorus indicates much use of fertilizers. EC and pH of study area are fairly good for agriculture.

VII. ACKNOWLEDGMENT:

Authors are thankful to Dr. M S Jangid prof. Baotany Department, Sir P.T.Science College, Modasa and Dr. R S Patel prof. Baotany Department Maninagar Science college for providing the necessary guidance, laboratory facilities and encouragement and also thankful to farmers of this area for their kind co-operation.

VIII. REFERENCES:

1. Gharekhan A, Arora S, et al. Distinguishing auto-fluorescence of Normal, Benign and Cancerous Breast Tissues through Wavelet Domain Correlation Studies. *Journal of Biomedical Optics*, 16(8):087003, 2011.
2. Vijayakumar R, Arokiaraj A, Martin DP. Micronutrients and their Relationship with Soil Properties of Natural Disaster Prone Coastal Soils. *Res J Chem Sci*. 1(1):pp.8-12, 2011.
3. Mico C, Recatala L, Peris M, Sanchez J. Assessing heavy metal sources in agricultural soil of an European Mediterranean area by multivariate analysis. *Chemosphere*, 65:863-72, 2006.
4. Mali VS, Zende NA, Verma UK. Correlation between soil physico-chemical properties and available micronutrients in salt affected soils, 17th WCSS 2002; Thailand
5. Gharekhan A, Oza AN, Sureshkumar MB, Pradhan A, Panigrahi PK. Study of polarized spectral features of human breast tissues.
6. Jackson ML. *Soil Chemical Analysis* 1973; Prentice Hall of India Pvt. Ltd. New Delhi.
7. Chauhan JS. Fertility status of soils of Birla Panchayat Samiti of Jodhpur district Rajasthan. M.Sc.(Ag.) 2011; Thesis: MPUAT Udaipur
8. Also published in *Virtual Journal of Biological Physics Research* 2011; 22(4) Kutch. vibrantgujarat.com Soils of Gujarat - <http://goo.gl/CF9Rb> Agriculture Contingency Plan for District, Kutch, Gujarat.
9. "Methods Manual-Soil Testing in India", Department of Agriculture & Cooperation Ministry of Agriculture Government of India, 2011.
10. Singh RP, Mishra SK. Available macronutrients (N, P, K and S) in the soils of Chirgaon block of district Varansi (U.P.) in relation to soil Characteristics. *Indian J Sci Res*, 3(1), pp.97-100, 2012.