

K- Shell Ionization of Neon and Magnesium due to impact of alpha particles

Santosh Kumar^{1,*}, Sandeep Kumar Prasad², Omprakash³ and Dr. Ravi Ranjan⁴

¹Research Scholar, University Department of Physics, B.R.A. Bihar University, Muzaffarpur 842001, Bihar

²Assistant Professor, Department of Physics, M.P.S. Science College, Muzaffarpur

³Research Scholar, University Department of Physics, B.R.A. Bihar University, Muzaffarpur

⁴Assistant Professor, Department of Physics, Nitishwar Mahavidyalaya, Muzaffarpur

Abstract : In the present work, K-shell ionization cross-sections of Neon and Magnesium due to alpha impact have been calculated through the Binary Encounter Approximation model. The effects of Coulomb deflection of the projectile under the influence of target nucleus and increase in binding energy of the target electron because of presence of the projectile inside K- shell of the target have been incorporated. The cross-sections have been averaged over Hartree-Fock momentum distribution for the target electron. The calculated cross-sections have been found in satisfactory agreement with the available experimental values and other calculated results.

Keywords: K-shell ionization, Binary Encounter Approximation, Alpha impact, ECPSSR.

I. INTRODUCTION

Study of ionization of atoms and molecules due to impact of charged particles is important because of its applications in the field of gaseous discharge, astrophysics, atmospheric physics, plasma physics, laser physics etc. Moreover accurate values of ionization cross sections for inner shells (particularly K- shell) of atoms induced by heavy charged particles e.g. protons, alpha particles are important for trace element analysis of many elements¹. However, experimental measurements of cross sections are not always available at the desired values of impact energies and hence theoretical studies of such processes become important. Further these studies provide a test for suitability of the theories for charged particle impact direct ionization of atoms.

There are a number of theoretical methods for calculation of inner shell ionization cross sections due to heavy charged particles e.g. Plane Wave Born Approximation², Semi Classical Approximation³, Binary Encounter Approximation (BEA)⁴. Out of these, the BEA is simple and required less computational efforts but yields reasonable results.

II. TYPE STYLE AND FONTS

In the present work we have calculated K-shell ionization cross sections of atomic neon and magnesium due to impact of alpha particles in the BEA. In the present calculations, we have incorporated the effects of the following two physical processes on the ionization cross sections.

(i) Coulomb deflection effect:

A positively charged particle coming close to the target nucleus experiences repulsion due to the target nucleus. This reduces the velocity of the incident particle as well as changes its trajectory during collision. These two changes cause a reduction in the inner shell ionization cross sections. In the present work the above effects have been incorporated following Thomas and Garcia⁵ through the relation

$$\sigma(E_1) = \sigma(E_1') \left[\frac{1}{2} + \frac{1}{2} \left(1 - \frac{Z_1 Z_2 k e^2}{E_1 a_{2k}} \right)^2 \right]^2 \quad ; \quad \dots\dots (1)$$

where $\sigma(E_1')$ is the ionization cross section at the reduced energy.

$$E_1' = E_1 - \frac{Z_1 Z_2 e^2}{a_{2k}} \quad \dots\dots (2)$$

And $Z_{2k} = Z_2 - S_{2k}$

Z_1 and Z_2 are the nuclear charges of the projectile and the target respectively, a_{2k} is radiation of K- shell of the target whereas S_{2k} is screening constant for K-shell.

(ii) Effect of increase in binding of the target electron:

For ejection of an electron from an inner shell, the projectile has to penetrate deep into the shell. This causes an increase in the binding energy of the target electron. Under this condition, the electron response time is much shorter than the collision time which allows the electron to adjust to the presence of the projectile and thereby increases the binding energy of the electron. This feature reduces ionization cross sections at low impact energies and the effect gradually decreases with increase in the impact energy².

The above described effect has been incorporated in the expression for ionization cross sections by replacing the unperturbed binding energy U_{2k} of the atomic K-shell by $U_c = \epsilon U_{2k}$, U_c being the corrected binding energy, ϵ is a correction factor given by Brandt and Lapicki².

$$\epsilon = 1 + \left(\frac{2Z_1}{Z_{2k} \theta_{2k}} \right)^g \tag{3}$$

θ_{2k} is the reduced binding energy for the K-shell and is given by

$$\theta_{2k} = \frac{U_{2k}}{Z_{2k}^2}$$

The factor g is an impact velocity dependent term and is given by⁶

$$g = \frac{1 + 5x + 7.14x^2 + 4.27x^3 + 0.947x^4}{(1 + x)^5} \tag{4}$$

where

$$x = \frac{V_1}{\left(\frac{1}{2} \theta_{2k} V_{2k} \right)}$$

Vriens' expressions⁷ for ionization cross section, incorporating the above mentioned effects can be written as

$$Q_I(s, t) = \frac{(s+s')^2 Z_1^2}{s^2 s'^2 U_c^2} \left[1 + \frac{2t^2}{3} - \frac{1}{4(s'^2 - t^2)} \right] (\pi a_0^2); \quad 1 \leq 4s'(s'-t) \tag{5}$$

$$= \frac{(s+s')^2 Z_1^2}{2s^2 s'^2 U_c^2 t} \left[\frac{1}{4(s'+t)} + t + \frac{2}{3} \left\{ 2s'^3 + t^3 - (1+t^2)^{3/2} \right\} \right] (\pi a_0^2); \tag{6}$$

$$4s'(s'-t) \leq 1 \leq 4s'(s'+t) \\ = 0; \quad 1 \geq 4s'(s'+t) \tag{7}$$

In the above equations s and t are dimensionless variables defined as

$$t^2 = \frac{V_{2k}^2}{V_0^2} \text{ and } s^2 = \frac{V_1^2}{V_0^2}$$

whereas,

$$(s')^2 = s^2 - \frac{1.058 Z_1 Z_{2k}}{1836 M a_{2k} U_c}$$

V_0^2 being corrected binding energy in Rydberg, while V_1 , V_{2k} and M (mass of the projectile) are expressed in atomic units.

The above expressions have been averaged over Hartree-Fock velocity distribution for the target electron. The Hartree-Fock radial functions given by Clementi and Roetti⁸ have been used to construct the momentum distribution for target electrons. The quantum mechanical values of K-shell energy given by Clementi and Roetti⁸, have been used in the present work. The quantum mechanical values of the points of maximum radial probability reported by Desclaux⁹ have been used as the shell radii. The screening constant for the K-shell has been taken equal to 0.3

Table.1 Alpha Particle Impact K-shell Ionization Cross sections of Ne


Impact Energy (MeV)	Expt. Ion. Cross sections (10 ⁻²² cm ²) Ref:10	Expt. Ion. Cross sections (10 ⁻²² cm ²) Ref:11	Cal. ECPSSR Ion. Cross sections (10 ⁻²² cm ²) Ref:12	Present Calculations (10 ⁻²² cm ²)		
				(A)	(B)	(C)
0.116	0.195		0.09	2.93	2.50	0.137
0.135	0.314		0.187	4.65	4.05	0.244
0.200		0.767	1.17	14.7	13.4	1.05
0.250		2.03	3.18	27.2	25.5	2.37
0.300		3.94	7.06	44.5	42.2	4.52
0.400		12.2	23.9	92.9	89.4	12.2
0.500		30.0	57.8	159.0	154.0	25.5
0.600		62.2	114.0	240.0	234.0	45.4
0.800		172.0	298.0	438.0	431.0	107.0
1.000		343.0	567.0	671.0	656.0	198.0
1.200		548.0	890.0	922.0	913.0	314.0

A- Without modification

B - Including Coulomb effect

C - Including Coulomb and Binding effect

Figure-2: K- shell ionization cross-sections of Neon




- A: Present calculated cross sections without any modification
- B: Present calculated cross sections including only Coulomb effect
- . - C: Present calculated cross sections including both the processes
- ECPSSR cross sections Ref.:12
- Experimental ionization cross sections
- Ref.:10
- ▲ Ref.:11

Table.2 K-Shell Ionization Cross sections of Magnesium

Impact Energy (MeV)	Expt. Ion. Cross sections (10 ⁻²¹ cm ²) Ref:13	Cal. ECPSSR Ion. Cross sections (10 ⁻²¹ cm ²) Ref:12	Present Calculations (10 ⁻²¹ cm ²)		
			(A)	(B)	(C)
1.0	9.6	7.8	13.0	12.8	3.62
1.5	16.7	24.9	28.4	28.1	10.5
2.5	57.7	71.0	67.0	66.3	31.5
3.0	79.7	93.1	105.0	105.0	45.5
3.5	105.0	112.0	131.0	131.0	71.6
4.0	118.0	128.0	149.0	149.0	91.1
4.5	135.0	142.0	163.0	162.0	106.0
5.0	153.0	153.0	173.0	173.0	118.0

Figure-2: K- shell ionization cross-sections of Magnesium



.....

- . - . - .

—————

■

A: Present calculated cross sections without any modification

B: Present calculated cross sections including only Coulomb effect

C: Present calculated cross sections including both the processes

ECPSSR cross sections Ref.:12

Experimental ionization cross sections

Ref.:13

III. RESULTS AND DISCUSSION

3.1 K – shell ionization cross sections of Neon

We have calculated K – shell ionization cross sections of Neon due to impact of alpha particles in the energy range from 0.116 MeV to 1.2 MeV and the results have been shown in the table 1 and figure 1. Observed results have been compared using the several of experimental results to Harrison et.al.¹⁰ and Langenberg and vanEck¹¹. We have presented a comparison of our results and the experimental observation and the theoretical calculations in ECPSSR¹². At low impact energies our results are slightly underestimate the experimental values but with increase in impact energy the agreement improves. This is a general trend of BEA cross sections. Overall agreement of our results with the experiments is satisfactory; they are always within a factor of two. The present results are also in reasonable agreement with the values of cross sections obtained in ECPSSR theory.

3.2 K – shell ionization cross sections of Magnesium

We have performed K – shell ionization cross sections for Magnesium due to impact of alpha particle in the range of 1.0 MeV to 5.0 MeV and the results have been shown in the table-2 and figure-2. Observed results have been compared using the several of experimental results to Sellers et.al.¹³ and calculations of ECPSSR¹². Experimental results are available only in a small range of impact energy. Our calculations overestimated experiments but the agreement improves with its increase in the impact energy. Present results are also in reasonably good agreement with ECPSSR¹² calculations.

IV. CONCLUSION

A critical study of the results presented in section 3 leads to the conclusion that the present method gives reasonable values of alpha impact K- shell ionization cross sections of Neon and Magnesium with very small computational effort. The present results are also in close agreement with the cross sections obtained using more improved method namely ECPSSR.

REFERENCES

- [1] A. P. L. Bertol, J. Trincavelli, R. Hinrichs and M. A. Z. Vascon cellos; Nucl. Inst. Methods in Phys. Res. B. **318**, 19 (2014).
- [2] W. Brandt and G. Lapicki; Phys. Rev. A **20**, 465 (1979).
- [3] J. Bang and J. M. Hansteen; Mat. Phys. Medd. **37**, No. 13 (1950).
- [4] J. D. Garcia; Phys Rev. A **1**, 280 (1970).
- [5] B. K. Thomas and J. D. Garcia; Phys. Rev. A **179**, 94 (1969).
- [6] G. Basbas, W. Brandt and R. Laubert; Phys. Rev. A **7**, 1971 (1973).
- [7] L. Vriens; Proc. Phys. Soc. **90**, 935 (1967).
- [8] E. Clementi and C. Roetti; At.Data Nucl.Data Tables **14**, 187 (1974).
- [9] J. P. Desclaux; At.Data Nucl.Data Tables **12**, 325 (1973).
- [10] K. G. Harrison, H. Tawara and F. J. De Heer, Physica**66**, 6 (1973).
- [11] A. Langenberg and J. venEck, J. Phys. B **9**, 2421 (1976).
- [12] G. Lapicki, J. Phys. Chem. Ref Data **18** No.1 (1989).
- [13] B. Sellers, F. A. Hansen and H. H. Wilson, Phys. Rev. **182**, 90 (1969).