

EFFECTS OF AEROBIC EXERCISE ON SELECTED PHYSIOLOGICAL PARAMETERS AMONG INACTIVE YOUNG WOMEN

* P. Anisha¹ & **Dr. A. Subradeepan²

¹Ph.D. Scholar, Dept. of Physical Education, Annamalai University, Chidambaram, Tamilnadu, India,

²Assistant Professor, Dept. of Physical Education, Annamalai University, Chidambaram, Tamilnadu, India.

The purpose of the study was to find out the effects of Aerobic exercise on selected physiological parameters among inactive young women. To achieve the purpose of the study thirty inactive young women were selected randomly as subjects from Women's Christian College, Nagercoil, Kanyakumari District, Tamilnadu, India and their age were ranged from 18 to 25 years. The subjects divided into two groups in equal numbers (N=20). Group I underwent Aerobic exercise group and group II acted as control group who did not attended any special training other than their daily college schedule curriculum. The duration of the training period was restricted into six weeks for three alternative days per week. The pre and post tests data were collected before and after the training period. The dependent variables pulse rate, and VO₂ Max were tested by standardized test items respectively. The collected data from the two groups prior to and after the experimental treatments on selected variables pulse rate, and VO₂ Max were statistically analyzed by using the statistical technique of dependent 't' test and analysis of covariance (ANCOVA). In all the cases the level of confidence was fixed at 0.05 significant. The result of the study indicated that the experimental group had shown significantly improved in pulse rate, and VO₂ Max variables among inactive young women due to the effects of aerobic exercise. However the control group did not shown any significant improvement on selected variables such as pulse rate, and VO₂ Max.

Keywords: Aerobic exercise, physiological parameters, pulse rate, and VO₂ Max.

Introduction

Aerobic exercise is physical exercise of low to high intensity that depends primarily on the aerobic energy-generating process "Aerobic" means "relating to, involving, or requiring free oxygen", and refers to the use of oxygen to adequately meet energy demands during exercise via aerobic metabolism. Generally, light-to-moderate intensity activities that are sufficiently supported by aerobic metabolism can be performed for extended periods of time what is generally called aerobic exercise might be better termed "solely aerobic", because it is designed to be low-intensity enough so that all carbohydrates are aerobically turned into energy. And bulk of the energy in this type of exercise is due to mitochondria ATP production which relies on oxygen for the metabolism of carbs, proteins and fats for energy **Smith (2007).most health power full workout takes this training very usefully for women society.**

Statement of the Problem

The purpose of this study was to find out the effects of Aerobic exercise on selected physiological parameters among inactive young women.

Hypothesis

1. It was hypothesized that there may be a significant difference on selected among inactive young women.
2. There would be a significant difference among experimental and control groups on selected Pulse Rate
3. There would be a significant difference among experimental and control groups on selected VO₂ max test.

Methodology

The purpose of this study was to find out the effects of Aerobic exercise on selected physiological parameters among inactive young women. To achieve the purpose of the study thirty inactive young women were selected randomly as subjects from Women's Christian College, Nagercoil, Kanyakumari District, Tamilnadu, India and their age were ranged from 18 to 25 years. The inactive young women were assigned at random into two groups of each ten (N=10). Group-I underwent Aerobic exercise and Group-II acted as control group who did not attended any special training other than their regular daily college schedule curriculum. The duration of the training period was restricted to six week for three alternative days per week. The pre and post data were collected before and after the training period. The dependent variables physiological parameters were tested by standardized tests Pulse Rate and VO₂ max.

Analysis of the data

The effects of Aerobic exercise on selected physiological parameters among inactive young women. Were analyzed and presented below.

Back strength Test

The t-test on Pulse Rate test (**second**) of the pre and post test scores of Aerobic exercise group and control group have been analyzed and presented in table I.

TABLE-I
THE PRE TEST AND POST TEST SCORES OF EXPERIMENTAL AND CONTROL GROUP ON PULSE RATE TEST

Group	Pre Mean	SD	Post Mean	SD	Obtained t-ratio
Experimental	65.06	3.03	61.20	2.44	6.81*
Control	65.12	2.10	66.03	1.69	1.01

*Significant at .05 level. (The table value required for 0.05 level of significance with df 9 is 2.26)

The table I show that the pre-test mean value of Aerobic exercise group and control group are 65.06 and 65.12 respectively and the post test means are 61.20 and 66.03 respectively. The obtained dependent t-ratio values between the pre and post test means of Aerobic exercise group and control group are 6.81 and 1.01 respectively. The table value required for significant difference with df 9 at 0.05 level is 2.26. Since, the obtained 't' ratio value of Aerobic exercise group was greater than the table value, it is understood that Aerobic exercise group had significantly improved the pulse rate test. However, the control group has not improved significantly. The 'obtained t' value is less than the table value, as they were not subjected to any specific training.

Analysis of covariance (ANCOVA) on Pulse Rate test of experimental and control groups have been analyzed and presented in table II.

TABLE II
ANALYSIS OF COVARIANCE (ANCOVA) ON PULSERATE TEST OF EXPERIMENTAL GROUP AND CONTROL GROUP


Adjusted Post Test Means		Source of variance	Sum of squares	Ddf	Mean square	F – ratio
Experimental Group	Control Group	Between	2022.26	1	2022.26	30.01*
62.63	65.57	Within	1280.31	19	67.38	

* Significant at 0.05 level. (The table value required for significance at 0.05 levels with df 1 and 19 is 4.38)

Table II shows that the adjusted post test means values on Pulse Rate test. The obtained f- ratio of 30.01 for adjusted post test mean is greater than the table value 4.38 with df 1 and 19 required for significance at 0.05 level of confidence. The results of the study indicate that there is a significant mean difference exist between the adjusted post test means of Aerobic exercise and control groups on back strength test.

The bar diagram shows the mean values of pre test, post test and adjusted post test on Pulse Rate test of Aerobic exercise group and control group.

FIGURE I
PRE TEST, POST TEST AND ADJUSTED POST TEST MEAN VALUES OF AEROBIC EXERCISE AND CONTROL GROUPS ON PULSE RATE TEST



VO2 max test

The t-test on **VO2 max test** of the pre and post test scores of Aerobic exercise group and control group have been analyzed and presented in table I.

TABLE-II
THE PRE TEST AND POST TEST SCORES OF EXPERIMENTAL AND CONTROL GROUP ON VO2 max TEST

Group	Pre Mean	SD	Post Mean	SD	Obtained t-ratio
Experimental	50.60	1.65	55.46	1.60	13.39*
Control	50.59	1.63	50.61	1.50	1.42

*Significant at .05 level. (The table value required for 0.05 level of significance with df 9 is 2.26)

The table I shows that the pre-test mean value of Aerobic exercise group and control group are 50.60 and 50.59 respectively and the post test means are 55.46 and 50.61 respectively. The obtained dependent t-ratio values between the pre and post test means of Aerobic exercise group and control group are 13.39 and 1.42 respectively. The table value required for significant difference with df 1 and 9 at 0.05 level is 2.26. Since, the obtained 't' ratio value of Aerobic exercise group was greater than the table value, it is understood that Aerobic exercise group had significantly improved VO2 max test. However, the

control group has not improved significantly. The ‘obtained t’ value is less than the table value, as they were not subjected to any specific training.

Analysis of covariance (ANCOVA) on **VO2 max test** of experimental and control groups have been analyzed and presented in table II.

TABLE II
ANALYSIS OF COVARIANCE (ANCOVA) ON VO2 max TEST OF EXPERIMENTAL GROUP AND CONTROL GROUP


Adjusted Post Test Means		Source of variance	Sum of squares	Ddf	Mean square	F – ratio
Experimental Group	Control Group	Between	3926.10	1	3926.10	22.40*
51.53	50.14	Within	2481.05	19	130.58	

* Significant at 0.05 level. (The table value required for significance at 0.05 levels with df 1 and 19 is 4.38)

Table II shows that the adjusted post test means values on **VO2 max test**. The obtained f- ratio of 22.40 for adjusted post test mean is greater than the table value 4.38 with df 1 and 19 required for significance at 0.05 level of confidence. The results of the study indicate that there is a significant mean difference exist between the adjusted post test means of Aerobic exercise and control groups on **VO2 max test**.

The bar diagram shows the mean values of pre test, post test and adjusted post test on **wallsit test** of Aerobic exercise group and control group.

FIGURE II
PRE TEST, POST TEST AND ADJUSTED POST TESTMEAN VALUES OF AEROBIC EXERCISE AND CONTROL GROUPS ON VO2 max TEST



Conclusions

1. There was significant improvement on Pulse Rate test due to the effects of Aerobic exercise among inactive young women
2. There was significant improvement on VO2 max test due to effects of Aerobic exercise among inactive young women
3. However the control group had not shown any significant improvement on any of the selected variables.

Acknowledgement

I acknowledge sincere gratitude to my guide who support and guided in a proper way. I would like to place my sincere thanks to the Department of physical Education; Annamalai University provided study supportive materials. Also I express my sincere thanks to the volunteers of this study.

References

- Akazawa N, Tanahashi K, Kosaki K, Ra SG, Matsubara T, Choi Y, Zempo-Miyaki A Maid S.(2018). "Aerobic exercise training enhances cerebrovascular pulsatility response to acute aerobic exercise in older adults". *Physiol Rep.* Apr;6(8):e13681. doi: 10.14814/phy2.13681
- Abdelmalki A, Fimbel S, Mayet-Sornay MH, Sempore B, Favier R.(1996). "Aerobic capacity and skeletal muscle properties of normoxic and hypoxic rats in response to training". *Pflugers Arch.* Mar;431(5):671-9.
- Chandler, T.J. (1994). "Physiology of aerobic fitness/endurance." *Instr Course Lect.* 1994 ;43 :11-5
- Gopinath, V. (2008). "Effects of Aerobic Exercise (Submaximal) on Biochemical Parameters", A Paper Presented at the International Conference of Exhibition on Sports Technology.
- Jothi and Mangayarkarasi, (2004), "Effect of Aerobic Exercise on Selected Physiological Variables of Pregnant Women", *Journal of Physical Education and Exercise Sciences.*
- Nilsson BB, Hellens B.(2008). "Group-based aerobic interval training in patients with chronic heart failure: Norwegian Ullevaal Model." *PhysTher.* Apr;88(4):523-35
- Martinez-Rodriguez A.(2014). "Effects of diet and aerobic or anaerobic exercises on eating disorders" *Nutr Hosp.* Oct 19;31(3): 1240-5. doi: 10.3305 / nh.2015.31.3.8131
- Nilsson BB, Hellens B.(2008). "Group-based aerobic interval training in patients with chronic heart failure: Norwegian Ullevaal Model." *PhysTher.* Apr;88(4):523-35
- Pattyn N, Beulque R, Cornelissen V.(2018). "Aerobic Interval vs. Continuous Training in Patients with Coronary Artery Disease or Heart Failure: An Updated Systematic Review and Meta-Analysis with a Focus on Secondary Outcomes". *Sports Med.* May;48(5):1189-1205. doi: 10.1007/s40279-0180885-5.
- Park HY, Hwang H, Park J, Lee S, Lim K.(2016). "The effects of altitude/hypoxic training on oxygen delivery capacity of the blood and aerobic exercise capacity in elite athletes - a meta-analysis" *Journal Exercise Nutrition Biochemistry.* Mar 31;20(1):15-22. doi: 10.20463/jenb.2016.03.20.1.3.
- Rowland, A. (1995). "Aerobic response to endurance exercise training in children. *Pediatrics.*" Oct;96(4 Pt 1):654-658.
- Selvan V. Sudha, (2008). "Effect of Aerobic Exercise on Selected Physiological Variables among College girls" A Paper Presented at National Seminar on Professional and Scientific Approaches in Physical Education and Sports Sciences.
- Saghiv MS, Sira DB, Goldhammer E, Sagiv M.(2017). "The effects of aerobic and anaerobic exercises on circulating soluble-Klotho and IGF-I in young and elderly adults and in CAD patients" *Journal CircBiomark.* Sep 28;6:1849454417733388. doi: 10.1177/1849454417733388.
- Selvalakshmi, (2007). "Effect of varied aerobic training programme on obese women working in IT companies", A Paper Presented at the National Seminar on Multidimensional Futuristic Approach to excellence in physical education and Sports.