

# Determination of molar refraction and polarizability Constant of Substituted Thiazolyl Schiff's Bases in Binary Solvent Mixtures with Varying Concentration.

<sup>1</sup>Dr. Rupali S. Talegaonkar, <sup>2</sup>Dr. A.S. Burghate

<sup>1</sup>Assistant Professor, <sup>2</sup> Associate Professor

<sup>1</sup>Department of Chemistry, Mahatma Fule Arts, Commerce and Sitaramji Chaudhari Science Mahavidyalaya, Warud, Dist-Amravati.

**Abstract:** Density, Refractive indices of pure solvent 1,4-Dioxane, Acetone and DMSO and ligand solution of 2-[3-(4-methoxy-phenyl)-1-(4-phenyl-thiazol-2-ylimino)-allyl]-phenol in 70% 1,4-Dioxane, Acetone and DMSO have been studied at 30±0.1°C. Temperature with varying concentrations. The data obtained is utilized to determine the molar refraction and polarizability constant. Calculated data is used to study the solute-solute, solute-solvent and solvent-solvent interaction in the system.

**Key Words :** Refractive indices, Densities, Molar refraction and Polarizability constant.

## INTRODUCTION:-

Refractive index is one of the most important properties of liquid. The measurement of the refractive index of liquids is an important work in engineering and science. Transmission and reflection detections near critical angles related to total internal reflections are common methods in refractive index measurement.

When a ray of light passes from one medium to another, it suffers refraction, that is change in direction. If it passes from a less dense to a denser medium, it is refracted towards the normal so that the angle of refraction ( $r$ ) is less than the angle of incidence ( $i$ ) the refractive index ( $n$ ) of the medium is the ratio of velocity of light in vacuum to the velocity of light in the medium. Refractive index is an important additive property of the structural arrangement of atom in molecule

Refractive index can be measured easily with high degree of accuracy. The values of refractive index depend on the temperature as well as on wavelength of light used.

Oswal et.al<sup>1</sup> have been studied refractivity properties of some homologues series such as n-ethanoate, methyl alkanoates, and ethyl alkanoates etc. A.N. Sonar<sup>2</sup> and N.S. Pawar have studied the molar refraction and polarizability constant of substituted heterocyclic compounds in different media from refractive indices. Burghate et.al<sup>3</sup> have studied the molar refraction and polarizability constant of substituted chalcones in different percentage of acetone-water mixture.

J.D. Pandey et.al<sup>4</sup> have studied the refractometric and dielectric studies of binary liquid mixtures at different temperature.

A.B.Nikumbh et.al<sup>5</sup> have studied the molar refraction and molarisability constant of aqueous solutions of  $\text{NH}_4\text{NO}_3$  and  $\text{KBrO}_3$  at different temperature.

R.A. Synowicki et.al<sup>6</sup> implemented two different fluid measurement techniques to determine the refractive index of fluids on a commercial spectroscopic ellipsometer system. In first technique they use roughened glass to which liquid is applied. And in second they use the prism minimum deviation technique in a hollow prism cell. The advantages and disadvantages of both the techniques discussed.

The present work deals with the study of molar refraction and polarizability constant of substituted thiazolyl Schiff's base such as 2-[3-(4-methoxy-phenyl)-1-(4-phenyl-thiazol-2-ylimino)-allyl]-phenol in non-aqueous solvent such as 1,4-dioxane, acetone and dmsol (with different concentrations).

**EXPERIMENTAL:-**

Above solution of the various compositions i.e. of ethanol, acetone and dioxane were prepared by dissolving an appropriate amount by weight. For density measurement, all the weighings were made on contech balance having accuracy (0.001gm). The refractive index of solvent mixtures/solutions were measured using different percentages by Abbe's refractometer ranging reading from 1.3000 to 1.70. The temperature of prism box was maintained constant by circulating water from thermostat at 30°C. ( $\pm 0.1^\circ\text{C}$ ). The refractometer was calibrated using glass test pieces of known refractive index supplied with the instrument.

The molar refraction of binary by used mixtures such as dioxane-water, acetone-water and ethanol-water mixture were determined from

$$R_{a-w} = X_1R_1 + X_2R_2$$

Where  $R_1$  and  $R_2$  are molar refractions of medium and water respectively.

The molar refraction of solvent and solution were determined using Lorentz-Lorentz<sup>7</sup> equation.

$$R_m = \frac{(n^2 - 1)}{(n^2 + 2)} \left\{ \frac{[x_1m_1 + x_2m_2 + x_3m_3]}{d} \right\}$$

Where  $R_m \rightarrow$  molar refraction

$\eta \rightarrow$  refractive index of solution

$x_1 \rightarrow$  mole fraction of medium

$x_2 \rightarrow$  mole fraction of water

$x_3 \rightarrow$  mole fraction weights of solute

$M_1, M_2$  &  $M_3 \rightarrow$  molecular weights of medium water and solute respectively

$d \rightarrow$  density of solution

The polarizability constant ( $\alpha$ ) of ligand is calculated from the following equation.

$$R_{\text{lig}} = \frac{4}{3}\pi N_0 \alpha$$

Where  $N_0$  is the Avagadro's number.

The calculated values of molar refraction and polarizability constant are shown in table.

**The values of molar refraction and polarizability constant at  $30 \pm 0.1^\circ\text{C}$  temp. System : 2-[3-(4-methoxy phenyl)-1-(4-phenyl-thiazol-2-ylimino)-allyl] – phenol.**

**Table 1 : for 70% dioxane-water mixture.**

SR. NO.	Conc.	Density	R.I.	Rm(mix) Cm <sup>3</sup> mole <sup>-1</sup>	$\alpha \times 10^{-23}$
1	0.01	8.938	1.418	1.787	0.0709
2	0.02	9.05	1.417	1.762	0.0698
3	0.03	9.20	1.416	1.729	0.0685
4	0.04	9.35	1.416	1.703	0.0675
5	0.05	9.49	1.415	1.673	0.0663

Table 2 : for 70% acetone-water mixture


SR. NO.	Conc.	Density	R.I.	Rm(mix) Cm <sup>3</sup> mole <sup>-1</sup>	$\alpha \times 10^{-23}$
1	0.01	7.382	1.362	1.427	0.0566
2	0.02	7.458	1.363	1.416	0.0561
3	0.03	7.498	1.364	1.413	0.0560
4	0.04	7.578	1.364	1.397	0.0554
5	0.05	7.658	1.365	1.386	0.0549

Table 3 : for 70% dms0-water mixture

SR. NO.	Conc.	Density	R.I.	Rm(mix) Cm <sup>3</sup> mole <sup>-1</sup>	$\alpha \times 10^{-23}$
1	0.01	10.1	1.468	1.611	0.0639
2	0.02	10.12	1.467	1.605	0.0636
3	0.03	10.14	1.466	1.599	0.0634
4	0.04	10.26	1.465	1.578	0.0625
5	0.05	10.32	1.464	1.566	0.0621

The values of molar refraction and polarizability constant at 30±0.1 °C temp.

System: 2-[3-(4-methoxy-phenyl)-1-(4-phenyl-thiazol-2-ylimino)-allyl]-4-methyl phenol(L<sub>3</sub>)



Graph 1 : variation of polarizability constant with respect to concentration of solvent.

## RESULTS AND DISCUSSION:-

The present investigation considers the density and refractive index . measurement of 2-[3-(4-methoxy-phenyl)-1-(4-phenyl-thiazol-2-ylimino)-allyl] -phenol in binary mixtures dioxane-water, acetone-

water and dmsol-water. The results obtained for variation in concentration of binary mixtures are reported in above tables no.1-3 and respective graphical representation is shown in graph.

In the present investigation, there is decrease in polarizability as well as molar refraction with increase in concentration of binary mixture with respect to more polar solvent. This may be due to dispersion force. It is the molecular force which arises from temporary dipole moment. The cumulative dipole-dipole interaction may create weak dispersion force resulting in decrease in molar refraction and polarizability.

From the results it may be predicted that for binary liquid mixtures on addition of mentioned compound there is decrease in molar refraction as well as polarizability. This may be due to the fact that the solvent-solvent interaction may be more strong than solute-solvent interaction.

#### ACKNOWLEDGMENT:

The authors are thankful to the Head Department of Chemistry and Principal of Mahatma Fule Arts, Commerce and Sitaramji Chaudhari Science Mahavidyalaya, Warud, District Amravati for providing facilities to carry out this research work.

#### REFERENCES.

1. Oswal, S.L. Oswal, P. Modi, P.S. Dave, J.P. and Gardas. R.L. 2004. Thermachemic Acta, 410:1
2. Sonar, A.N. and Pawar, N.S. 2010. Rasayan J. Chem. Vol. 3(2) : 250-254
3. Burghate, A. S. Agrawal, P.B. Quazi, S.W. and Narwade. M. L. 2001. Asian Journal of chemistry Vol. 13(4) : 1652-1654
4. Pandey, J.D. Chhabra, J. Soni, N.K. Tiwari, K.K. and Mishtra. R.K. 2006. Indian Journal of chemistry, 45 A : 653-656
5. Nikumbh, A. B. Rathi, M. V. 2016. American Journal of Engineering Research Vol.5 (11):195-200
6. Synowicki, R. A. Prihil, G. K. Cooney, G. Herringer, C.M. and Green. S.E. 2004. Journal of vac sci Technol B. 22 (6) : 3450-3453
7. Kittel, C. Introduction to solid state physics 8<sup>th</sup> edition : 464