

# COMPARATIVE STUDY OF POTENTIAL ANTIMICROBIAL ACTIVITY OF ROOT AND LEAVES OF GINGER AGAINST SELECTED BACTERIA.

Rahul D. More\* Satish V. Gaikwad

Department of Chemistry, Siddharth Arts, Commerce & Science College, Jafrabad, Jalna-431206(M.S.)

\*Department of Chemistry, B. Raghunath Arts, Commerce & Science College, Parbhani-431401(M.S.)

**Abstract :** The main objective of present study was to evaluate comparative potential anti microbial activity of aqueous extract, ethanol extract and n hexane extract obtained from root and leaves of Ginger(*Zingiber officinale*) against selected bacteria *E coli*, *Salmenolla typhi*, & *Shingella SP*. Different ginger extracts were prepared. 6 extracts of roots & leaves of Ginger were prepared with three solvents Aqueous, Ethanol & n-Hexane and antimicrobial activity checked by agar disc diffusion method. Generally, all extracts obtained by Ginger showed antimicrobial activity against all selected microorganisms. The study showed that aqueous, ethanol & n-Hexane extracts shows near about same properties but leaves of ginger more effective properties than root. Concluded that *Ginger root and leaves both* may be a potential source for the curing of various diseases caused to animals

**Keywords:** *Zingiber officinale*, Ginger extract, Antimicrobial, disc diffusion

## I. INTRODUCTION

A number of medicinal properties can be shown by various parts of the *Ginger* tree. Almost all the parts of this plant: root, leaf, flowers etc. have been used for various ailments in the indigenous medicine of various countries [1]. Antimicrobial agents is very important for the control of pathogenic microbes, especially for the treatment of infections caused by resistant microbes. Medicinal herbs with antimicrobial activities are considered a potent source of novel antimicrobial functions. *Zingiber officinale* is widely used as a vegetable, functional food and medicinal plant that has rich nutritional composition with diverse pharmacological activities [2-4]. In recent times, focus on plant research has increased all over the world and a large body of evidence has collected to show immense potential of medicinal plants used in various traditional systems.[5]. Ginger is a natural antioxidant, antidiabetic, antiulcer, antiobesity, antibiotic, outstanding immune builder used in many countries from ancient times. A few researchers have investigated the antimicrobial activity of *GINGER* (Lam) extracts against some pathogenic bacteria .[7-8] In this present study aqueous extracts, n hexane extracts and ethanol extracts obtained from leaves and roots for their antimicrobial activity.

## II. MATERIAL AND METHODS

### Collection of Plant Material

Fresh Ginger root and leaves were obtained from campus of Siddhartha College, Jafrabad, Jalna (Maharashtra). The leaves and roots were identified and confirmed by the Botanist at the Botany Department of the present institution. All materials were washed with tap water to remove impurities. It were dried under shade for 1 week, all dried materials were ground in a mixer grinder separately, which were easily grinded into the powder form. Same processes repeated 4 to 5 times and it stored in an air tight container for further use.

### Preparation of Extracts

From each 30 gm of the powdered sample of roots and leaves were separately extracted in 500ml conical flasks with 100 ml of deionised distilled water (aqueous extraction) while 100ml each of n-Hexane, and ethanol, (solvent extraction). The conical flasks were plugged with rubber corks, then shaken at 120 rpm for 30 min and allowed to stand at room temperature for 6 day. The extracts were separately filtered using Whatman filter paper no. 1. The resulting filtrates were centrifuged. After centrifugation, supernatants were labeled as "Extract AQL" and "Extract AQR" for the aqueous extracts of leaves and roots respectively. Similarly labeled for solvent extraction as "Extract NHX" and "Extract ETH" for n-Hexane and ethanol respectively. Totally six extract of Ginger were prepared with different solvent and different part. All extracts were properly labeled and kept at 4°C until use.

### Antimicrobial properties of extracts

The antimicrobial activity of the various extracts was tested using the disc diffusion method and minimum inhibitory concentration (MIC) determination.

### The pathogenic microorganism

The three selected common pathogenic microorganism were used in the study, among those one was gram negative *Escherichia Coli*, *Salmonella Typhi* and *Shingella SP*

### Preparation of microbial cultures

All these bacterias were cultured on the Muller Hinton Agar. On this agar a loop full of each bacterial strain was inoculated in 6 ml of nutrients both. These plate agars were incubated in incubator for 24 hours at room temperature. And these were used for incolumns.

**Inoculum preparation:**

To prepared bacterial inoculums , pure culture of micro-organisms was inoculated into 5 ml of sterile nutrient broth followed by incubation at 37°C. Till moderate turbidity developed.

The density of suspension inoculated onto the media for susceptibility test was determined by comparison with 0.5 McFarland standard of Barium sulphate solution. [10]

**Antimicrobial activity by Kirby-Bauer agar disc diffusion method**

The antimicrobial activity of each plant extracts was determined by disc agar diffusion technique described by Kirby Bauer et al (1966)[9] . The nutrient broth cultures of test bacteria were spreaded on the Muller Hinton Agar media in petriplates and microbes broth culture were applied on media by swabbing. The Whatmann filter paper discs 6mm were dipped in the 3 different concentrations of the plant extracts and were placed on to 3 different spots surface of agar plat. The diameter of zone of inhibition was measured in mm using a ruler and results were recorded.

**III. RESULT AND DISCUSSION**

Table.1 Antibacterial activity of different Ginger extracts against E.Coli, S.Typhi, Shingella SP

| Type of extract | Zone of Inhibition |                |                     |
|-----------------|--------------------|----------------|---------------------|
|                 | <i>E.Coli</i>      | <i>S.Typhi</i> | <i>Shingella SP</i> |
| AQUL            | 9                  | 14             | 16                  |
| AQUR            | 10                 | 12             | 14                  |
| NHXL            | 8                  | 6              | 8                   |
| NHXR            | 13                 | 11             | 13                  |
| ETHL            | 11                 | 15             | 17                  |
| ETHR            | 12                 | 13             | 15                  |

**IV. CONCLUSION**

The result of this research has demonstrated that *Z. Officinale* could become promising natural antimicrobial agents with potential application in therapeutic drugs for controlling the pathogenic bacteria. Inhibition of both Gram-positive and Gram-negative organisms by this plant extract depicts that it can serve as a source of broad spectrum antibiotics, which justified the traditional use of this plant for therapeutic purposes.

**REFERENCES**

- [1] Falodun A, Okenroba LO, Uzoamaka N (2006). Phytochemical Screening and anti-inflammatory evaluation of mjethandic and aqueous extracts of *Euphorbia heterophylla* Linn (Ephorbiaceae), *Afr. J.Biotechnol.*, 5(6): 529-531.
- [2] Goto C, Kasuya S, Koga K, Ohtomo H, Kagei N. 1990. Lethal efficacy of extract from *Zingiber officinale* (traditional Chinese medicine) or [6]-shogaol and [6]-gingerol in *Anisakis* larvae in vitro. *Parasitol Res* 76: 653–656.
- [3] Hamza, I. S., Ahmed, S. H., Aoda, H. (2009). Study the antimicrobial activity of Lemon grass leaf extracts . Ministry of science & Technology Abstract, 2009, 198–212.
- [4]Habsah, M., Amran, M., Mackeen, M. M., Lajis, N. H., Kikuzaki, H., Nakatani, N., et al. (2000).Screening of *Zingiberaceae* extracts for antimicrobial and antioxidant activities. *Journal of Ethnopharmacol- ogy*, 72, 403–410.
- [5]Hoffman T (2007). Antimicrobial activity of some medicinal plants from India. *Hawaii Med. J.*, 66: 326-327.
- [6]Hirasawa M, Shouji N, Neta T, Fukushima K, Takada K. 1999.Three kinds of antibacterial substances from *Lentinusedodes* (Berk.) Sing. (*Shiitake*, an edible mushroom). *Int J Anti- microb Agents* 11: 151–157.
- [7] Hiserodt RD, FranzblauSG, Rosen RT. 1998. Isolation of 6-, 8-, and 10-gingerol from ginger rhizome by HPLC and preliminary evaluation of inhibition of *Mycobacterium avium* and *Mycobacterium tuberculosis*. *J Agric Food Chem*46: 2504– 2508.
- [8]Jang KC, Kim SC, Song EU et al. 2003. Isolation and structural identification of antimicrobial substances from the rhizome of *Zinzibermioga* Roscoe. *J Kor Soc Agric Chem Biotechnol* 46: 246–250.
- [9]Jiang H, Solyom AM, Timmermann BN, Gang DR. 2005. Characterization of gingerol-related compounds in ginger rhizome (*Zingiber officinale* Rosc.) by high-performance liquid chromatography/electrospray ionization mass spectro- metry. *Rapid Column Mass Spectrom*19: 2957–2964.
- [10]Katsura H, Tsukiyama RI, Suzuki A, Kobayashi M. 2001. In vitro antimicrobial activities of bakuchiol against oral micro- organisms. *Antimicrob Agents Chemother*45: 3009–3013.
- [11]Kawai T, Kinoshita K, Koyama K, Takahashi K. 1994. Anti-emetic principles of *Magnolia obovata* bark and *Zingiber officinale* rhizome. *Planta Med* 60: 17–20.
- [12]Kikuzaki H, Tsai SM, Nakatani N. 1992. Gingerdiol related com- pounds from rhizomes of *Zingiber officinale*. *Phytochemistry* 31: 1783–1786.
- [13]Kaleeswaran, B., Ilavenil, S., & Ravikumar, S. (2010). Screening of phytochemical properties and antibacterial activity of *Cynodon dactylon* L . *International Journal of Current Research*, 3, 83-88.
- [14]Kumar, A., Kashyap, P., Sawarkar, H., Muley, B., & Pandey, A. (2011). Evaluation of antibacterial activity of *Cynodon dactylon* ( L.) Pers . *International Journal of Herbal Drug Research*, I(Ii), 31–35.
- [15]Langner E, Griefenberg S, Gruenwald J (2008). Antimicrobial activity of Ginger (*Zingiber officinalis*) in vitro. *Adu.Their.*, 25: 44.
- [16]Li XC, Cai L, Wu CD. 1997. Antimicrobial compounds from *Ceanothusamericanus* against oral pathogens. *Phytochemis- try* 46: 97–102.
- [17]Mahady GB, Pendland SL, Yun GS, Lu ZZ, Stoa A. 2003. Ginger (*Zingiber officinale* Roscoe) and the gingerols inhibit the growth of Cag A+ strains of *Helicobacter pylori*. *Anticancer Res* 23: 3699–