

# Combinatorics and Triple Sequence

Vipin Verma\*

*“Department of Mathematics, School of Chemical Engineering and Physical Sciences Lovely Professional University, Phagwara, Punjab, India.”*

**Keywords:** Combinatorics, Triple sequence

**Abstract:** In this paper a we have established and proved new generalised properties on one of the schemes of multiplicative Triple sequence using combinatorics.

## 1. Introduction

Sequence and series have wide applications, combinatorics is a strong concept of Number theory in mathematics with the help of combinatorics many problems on mathematics have been solved. Many mathematicians have generalised many properties on well-known Fibonacci and Lucas sequence using combinatorics. The concept triple sequence was first introduced by Jin-Zai Lee & Jia-Sheng Lee [1] in 1987. There are different schemes possible for multiplicative triple sequence, in this paper we have established and prove new generalised identities by using combinatorics approach

## 2. Multiplicative Triple sequence

The one of the schemes of Multiplicative Triple sequence is defined by the recurrence relations

$$\alpha_{n+2} = \gamma_{n+1}\gamma_n, \quad \beta_{n+2} = \alpha_{n+1}\alpha_n, \quad \gamma_{n+2} = \beta_{n+1}\beta_n \quad (2.1)$$

for all integer  $n \geq 0$ , with initial conditions

$$\alpha_0 = a, \alpha_1 = d, \beta_0 = b, \beta_1 = e, \gamma_0 = c, \gamma_1 = f$$

Where  $a, d, b, e, c$  and  $f$  are real numbers

**Theorem 2.1** If  $\alpha_n$  and  $\gamma_n$  are define by equation (2.1) then (for  $n \geq 0$ )

$$\gamma_{n+10} = \prod_{i=n}^{n+5} \alpha_i^{\binom{5}{i-n}} \quad (2.2)$$

**Proof:** Theorem can be proved by mathematical induction method on  $n$

For  $n = 1$  by equations (2.1) and (2.2) and the fact that  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

$$\prod_{i=1}^6 \alpha_i^{\binom{5}{i-1}} = \alpha_1 \alpha_2^5 \alpha_3^{10} \alpha_4^{10} \alpha_5^5 \alpha_6$$

by using equation (2.1) we have

$$\prod_{i=1}^6 \alpha_i^{\binom{5}{i-1}} = \gamma_{11}$$

which proves for  $n = 1$

Suppose the theorem is true for  $n = k$ , so by equation (2.2)

$$\gamma_{k+10} = \prod_{i=k}^{k+5} \alpha_i^{\binom{5}{i-k}} \quad (2.3)$$

Now to prove for  $n = k + 1$ , by using equation (2.1), (2.2) and the fact that  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

$$\prod_{i=k+1}^{(k+1)+5} \alpha_i^{\binom{5}{i-(k+1)}} = \alpha_{k+1} \alpha_{k+2}^5 \alpha_{k+3}^{10} \alpha_{k+4}^{10} \alpha_{k+5}^5 \alpha_{k+6}$$

by using equation (2.1) we have

$$\prod_{i=k+1}^{(k+1)+5} \alpha_i^{\binom{5}{i-(k+1)}} = \gamma_{(k+1)+10}$$

which proves the theorem.

**Theorem 2.2** If  $\beta_n$  and  $\gamma_n$  are define by equation (2.1) then (for  $n \geq 0$ )

$$\beta_{n+10} = \prod_{i=n}^{n+5} \gamma_i^{\binom{5}{i-n}} \quad (2.4)$$

**Proof:** Theorem can be proved by mathematical induction method on  $n$

For  $n = 1$  by equations (2.1), (2.4) and the fact that  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

$$\prod_{i=1}^6 \gamma_i^{\binom{5}{i-1}} = \gamma_1 \gamma_2^5 \gamma_3^{10} \gamma_4^{10} \gamma_5^5 \gamma_6$$

by using equation (2.1) we have

$$\prod_{i=1}^6 \gamma_i^{\binom{5}{i-1}} = \beta_{11}$$

which proves for  $n = 1$

Suppose the theorem is true for  $n = k$ , so by equation (2.4)

$$\beta_{k+10} = \prod_{i=k}^{k+5} \gamma_i^{\binom{5}{i-k}} \quad (2.5)$$

Now to prove for  $n = k + 1$ , by using equation (2.1), (2.4) and the fact that  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

$$\prod_{i=k+1}^{(k+1)+5} \gamma_i^{\binom{5}{i-(k+1)}} = \gamma_{k+1} \gamma_{k+2}^5 \gamma_{k+3}^{10} \gamma_{k+4}^{10} \gamma_{k+5}^5 \gamma_{k+6}$$

by using equation (2.1) we have

$$\prod_{i=k+1}^{(k+1)+5} \gamma_i^{\binom{5}{i-(k+1)}} = \beta_{(k+1)+10}$$

which proves the theorem.

**Theorem 2.3** If  $\alpha_n$  and  $\beta_n$  are define by equation (2.1) then (for  $n \geq 0$ )

$$\alpha_{n+10} = \prod_{i=n}^{n+5} \beta_i^{\binom{5}{i-n}} \quad (2.6)$$

**Proof:** Theorem can be proved by mathematical induction method on  $n$

For  $n = 1$  by equations (2.1), (2.6) and the fact that  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

$$\prod_{i=1}^6 \beta_i^{\binom{5}{i-1}} = \beta_1 \beta_2^5 \beta_3^{10} \beta_4^{10} \beta_5^5 \beta_6$$

by using equation (2.1) we have

$$\prod_{i=1}^6 \beta_i^{\binom{5}{i-1}} = \alpha_{11}$$

which proves for  $n = 1$

Suppose the theorem is true for  $n = k$ , so by equation (2.6)

$$\alpha_{k+10} = \prod_{i=k}^{k+5} \beta_i^{\binom{5}{i-k}} \quad (2.7)$$

Now to prove for  $n = k + 1$ , by using equation (2.1), (2.6) and the fact that  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

$$\prod_{i=k+1}^{(k+1)+5} \beta_i^{\binom{5}{i-(k+1)}} = \beta_{k+1} \beta_{k+2}^5 \beta_{k+3}^{10} \beta_{k+4}^{10} \beta_{k+5}^5 \beta_{k+6}$$

by using equation (2.1) we have

$$\prod_{i=k+1}^{(k+1)+5} \beta_i^{\binom{5}{i-(k+1)}} = \alpha_{(k+1)+10}$$

which proves the theorem.

## References

- [1] Cerda-Morales, G., On the Third-Order Jabosthal and Third-Order Jabosthal-Lucas Sequences and Their Matrix Representations, arXiv:1806.03709v1 [math.CO], 2018.
- [2] Feinberg, M., Fibonacci–Tribonacci, The Fibonacci Quarterly, 1s: 3 (1963) pp. 71–74, 1963.
- [3] J.Z. Lee and J.S. Lee, Some Properties of the Generalization of the Fibonacci sequence, The Fibonacci Quarterly, 25(2), 111–117, 1987
- [4] K T Atanassov, V Atanassova, A G Shannon, J C Turner, New Visual Perspective on Fibonacci Numbers, World Scientific Publishing Co. Pte. Ltd., World Scientific Publishing Co. Pte. Ltd., 2002.
- [5] Koshy, T. Fibonacci and Lucas Numbers with Applications; John Wiley and Sons Inc.: New York, NY, USA, 2001.
- [6] Ozdemir, G.; Simsek, Y. Generating functions for two-variable polynomials related to a family of Fibonacci type polynomials and numbers, Filomat 969–975, 30, 2016.