

# ROLE OF DIFFERENT AGRONOMIC PRACTICE ON UPTAKE OF NUTRIENTS AND YIELD OF RICE.

M. Saravana Perumal, G. Parimala\*, G. Murugan, P. Stalin and  
A. Balasubramanian

Department of Agronomy, Faculty of Agriculture,  
Annamalai University,  
Annamalai Nagar – 608 002, Tamil Nadu, India

\* Corresponding Author, Department of Agronomy, Faculty of Agriculture, Annamalai University.

## ABSTRACT

Field experiments were conducted during Late Samba Season at Experimental farm, Annamalai University to study the system of Rice intensification (SRI) and their effect on nutrient uptake and Yield of Rice Crop. The treatment has the influence of combined effect of age of the seedlings, spacing, weed management by conoweeder, chemical weeding, manual weeding, nutrient management with organic and inorganic fertilizer and alternate wetting and drying on the growth character, (plant height, tillers hill<sup>-1</sup>, tiller m<sup>-2</sup>, leaf area index, dry matter production) of SRI were studied. Among the treatments, SRI (use of 8-12 days old seedling, 25 × 25 cm of spacing, saturation of water, four times conoweeding only inorganic) significantly increased the growth characters. This was followed by SRI (use of 8-12 days old seedlings, 25 × 25 cm of spacing, four times conoweeding, saturation water, use of 75% inorganic + 25% organic).

**Key words:** Rice – uptake of nutrients –Yield.

## INTRODUCTION:

More than 90 percent of world's rice is grown and consumed in Asia. SRI is a holistic agro-ecological crop management technique seeking alternatives to the conventional high input oriented agriculture through effective integration of crop soil water continuum. In SRI, one of the sound principles is wider spacing (25 × 25) of plant leading to greater root growth by creating aerobic condition and better tillering potential. Rice fields are kept moist rather than continuous flooding thereby minimizing anaerobic condition and it improves the root growth and aerobic microorganisms. SRI has the edge over others as the method offers not only water saving but also enhances the yield and helps in environmental protection (Norman Uphoff, 2006). SRI increases rice yield two or three fold compared to farmers current rice yields (Norman Uphoff, 2002). Excessive weed growth is one of the major constrains in intermittent irrigation practice followed under SRI. Early and frequent weeding by cono or rotary weeder solves this problem (Stoopet *et al.*, 2002). Burying the weeds in the fields rather than removing them improve the crop yield. Besides, the soil gets aerated and the weeds get decomposed in the soil and turn into organic

matter. Due to this, the root and plant grow healthier and higher yields can be achieved. In SRI, methodology productivity will not only increase but input use efficiency will also be enhanced (Ghosh *et al.*, 2007). The present investigation was conducted to evaluate the principles of SRI and their contribution towards enhancement of growth and yield.

## MATERIALS AND METHODS:

Field experiments were conducted at Annamalai University Experimental Farm, Annamalainagar during late samba season with the cultivars of CO 43 to evaluate the SRI and their contribution towards nutrient uptake and grain yield. The Experimental Farm is situated at 11° 24' North latitude and 79° 44' longitude at an altitude of + 5.79 m above mean sea level. The experiments were taken up in a Randomized Block Design with four replication with seven treatments *viz.*, (T1) - SRI (Use of 8-12 days old seedling raised bed nursery, carefully transplanting at a spacing of 25 × 25, weed management with conoweeder 4 times, saturation of water management use of 75% inorganic +25% of organic, (T2) - T1 with 20 to 25 days old seedlings, (instead of 8-12 days old seedlings), (T3) - T1 with 20 × 15 cm spacing, (instead of 25 × 25 spacing), (T4) - T1 with only inorganic (RDF) (instead of organic + inorganic, (T5) - T1 with herbicide + manual weeding (instead of conoweeding), (T6) - T1 with alternate wetting and drying (instead of saturation of water management), (T7) - conventional transplanting (location specific best management practice).

## RESULT AND DISCUSSION:

The data on crop dry matter production (DMP) recorded at harvest in Late samba season is given in Table 1. The highest DMP was recorded under the treatment 8 -12 days old seedling, spacing 25x25 cm with conoweeding four times, saturation of water and using only inorganic (T<sub>4</sub>) recorded significantly higher DMP of 11901.20 kg ha<sup>-1</sup> at harvest in Late samba season. The conventional method of transplanting recorded lowest DMP of 5762.00 kg ha<sup>-1</sup> at harvest stage.

Regarding the crop nutrient uptake of N,P,K by the rice crop is given in Table 1. The highest uptake of N,P,K were recorded under the treatment of 8 -12 days old seedling, spacing 25x25 cm with conoweeding four times, saturation of water using only inorganic (T<sub>4</sub>) of 126.95 kg ha<sup>-1</sup>, 26.21 kg ha<sup>-1</sup> and 297.12 kg ha<sup>-1</sup> respectively. Highest Nutrient uptake was observed in (T<sub>4</sub>) and this might be due to better establishment and growth of plants, resulting in higher DMP, competition free environment between plants for better absorption of nutrients, water, sunlight and space. The increased uptake might be due to more functional root system per unit area which absorbed the nutrients released from the nature source. The least N,P,K uptake of 93.10 kg ha<sup>-1</sup>, 16.42 kg ha<sup>-1</sup> and 275.67 kg ha<sup>-1</sup> were recorded under the conventional method of transplanting (T<sub>7</sub>) respectively and this might be due to reduced supply of nutrients. The report of Hugar *et al.*,(2009), Thekar Singh *et al.*,(2010) and Harish Deshpande *et al.*,(2010) falls in line with our findings.

**Table 1: Effect of different agronomic practice on uptake of nutrients and yield of rice.**

Treatment	DMP (kg ha <sup>-1</sup> ) (Late Samba)	Nutrient Uptake(kg ha <sup>-1</sup> )			Rice Grain Yield (kg ha <sup>-1</sup> ) Late Samba
		Nitrogen Uptake (kg ha <sup>-1</sup> )	Phosphorous Uptake (kg ha <sup>-1</sup> )	Potassium Uptake (kg ha <sup>-1</sup> )	
T <sub>1</sub>	11692.02	124.76	24.44	294.41	4445
T <sub>2</sub>	8782.37	99.21	18.80	281.22	3758
T <sub>3</sub>	9532.80	102.93	19.42	282.92	3937
T <sub>4</sub>	11901.20	126.95	26.21	297.12	4450
T <sub>5</sub>	11578.20	120.13	20.92	290.77	4437
T <sub>6</sub>	11281.00	122.34	23.44	292.12	4440
T <sub>7</sub>	5762.00	93.10	16.42	275.67	3538
S. Ed	104.56	1.01	0.76	0.67	231
C.D (0.05%)	209.13	2.19	1.67	12.31	385

**CONCLUSION:**

The results shows that under SRI by adapting transplanting with 8-12 days old seedlings, saturation of water, 25 × 25 cm of spacing and only inorganic with four times conoweeding effectively increased the nutrient uptake in Late samba season which paved the way for increasing the rice yield.

**REFERENCES:**

- Ghosh, A., K.S. Rao, M.P. Pandey and A. Poonam. 2007. System of rice intensification – A holistic management towards enhancing rice production in future. Towards a learning alliance- SRI in Orissa.
- Harish H. Deshpande, P. Devasenapathy and Nagaraj M. Naik. 2010. Microbial population dynamics as influenced by application of organic manures in rice field. **Green Farming**, **1**: 356 – 359.
- Hugar, A.Y.,H. Chandrappa, H.M. Jayadeva, A. Sathish and G.B. Mallikarjun. 2009. Comparative performance of different rice establishment methods in Bhadra Command area. **Karnataka J. Agric. Sci.**, **22**: 992 – 994.
- Norman Uphoff, 2002. System of rice intensification for enhancing the productivity of land, labour and water. **J. Agric. Resource Management**, **1**: 43-49.

Norman Uphoff, 2006. The system of rice intensification as a methodology for reducing water requirements in irrigated rice production. Paper for International Dialogue on rice and water: Exploring options for food security and sustainable environments, IRRI, Los Banos, Philippines, March 7-8, 2006.

Stoop, W., N. Uphoff and A. Kassam. 2002. A review of agricultural research issues raised by the system of rice intensification (SRI) from Madagascar. Opportunities for improving farming system for resource poor farmer. *Agric. Sys.*, **71**: 249-274.

Thakar Singh, Taranvir Pal Singh and Ranjiv Sikka. 2010. Effect of date of transplanting, plant population and nitrogen levels uptake and nutrients status of soil in basmati rice. *J. Crop Res.* **40**:12 -15.

