

IMPROVEMENT OF VASE LIFE IN ANTHURIUM (*Anthurium andreanum*) cv. ACROPOLIS BY USING CHEMICAL SOLUTIONS

Ajish Muraleedharan^{1*}, P. Karuppaiah¹, S. Ramesh kumar¹ and J. L. Joshi²

¹Department of Horticulture, Faculty of Agriculture, Annamalai University, Annamalai nagar, Tamil nadu.

² Department of Genetics and Plant Breeding Faculty of Agriculture, Annamalai University, Annamalai nagar, Tamil nadu, India - 608002.

*Author for correspondence- ajishm1000@gmail.com.

ABSTRACT

Anthurium is the largest genus of the Araceae family and very popular with flower arrangers because of lasting qualities of flower. The majority of these species are perennial herbaceous plants, cultivated for ornamental purposes due to their attractive inflorescences. For prolonging the vase life of cut flowers chemical preservatives are used in the holding solutions. The experiment was conducted at Flora-tech floriculture unit at Kottarakara, kollam Dist, kerala state, India during july 2014 to find out the best chemical preservative solution for extending the vase life of anthurium cut flowers, the variety used for the study is Tropical. The present experiment was conducted to find out the appropriate preservative solution for extending the vase life of anthurium cut flowers. The experiment was conducted in Completely Randomized Design with three replications. Maximum days taken for spadix necrosis, Days taken for spathe blueing, Physiological loss in weight, Solution uptake and Vase life were recorded. The treatment (T₈) Al₂(SO₄)₃ 200 ppm + sucrose 4%) recorded maximum results in extending the vase life and flower quality.

Keywords: Anthurium, vase life, preservative solution, sucrose

INTRODUCTION

Anthuriums are gaining popularity due to higher returns per unit area and their beautiful and attractive long lasting flowers. They have gained popularity as one of the most important commercial ornamental crop of the modern world. They are very popular with flower arrangers because of bold effect and lasting qualities of flowers. Under normal conditions, flower stay fresh without much deterioration for a period of about two weeks. The plant produces blooms throughout the year, one bloom emerging from the axil of every leaf (Croat, 1980). Vase life of cut flowers can be prolonged by the addition of chemical preservatives (Nowak and Rudnicki, 1990). Different factors affect the vase life of cut flowers are chemical and physiological factors such as the content of stored foods of flower, humidity, light, and temperature of the place where vase is kept. Cut flowers are forced to continue living with reserved carbohydrates, proteins and fat for their longevity. Factors affecting water uptake such as air embolism

and duration of vascular occlusion contribute to cut flower senescence in Anthurium flowers. The major reasons for less vase life may be due to nutrient deficiency, bacterial and fungal infection, water stress induced wilting and vascular blockage and the action of ethylene in plant cells. Ethylene serves as a hormone in plants by stimulating and regulating the opening of flowers and the shedding of flowers (Serek, *et al.*, 1996). By applying various chemicals the post-harvest life of cut flowers can be extended. Wilting is the main cause in the termination of the vase life of cut flowers.

Cut flowers will decay if they are not able to draw water from the vase solution. Also clogging of vascular tissues of the stem by a material produced by phloem will block the absorption of water (Lu, *et al.*, 2010). Another important factor which helps the vase life is its content of stored foods. Sucrose act as a source of energy required for the continuation of the vase life of the cut flowers and also helped for the improvement in the keeping quality value of Anthurium cut flowers. Among all the different types of sugars, sucrose has been found to be the most commonly used sugar in prolonging vase life of cut flowers. Sugar has an important role in the longevity of flowers, especially cut flowers, because after harvest they receive no nutritional and hormonal support from the mother plant (Van Staden, *et al.*, 1994). Therefore, the present experiment was conducted to find out the appropriate preservative solution for extending the vase life, quality and postharvest behaviour of flowers.

MATERIAL AND METHODS

The experiment was conducted at Flora-tech floriculture unit at Kottarakara, kollam Dist, kerala state, India during july 2014 to find out the best chemical preservative solution for extending the vase life of anthurium cut flowers, the variety used for the study is Acropolis. Ten treatments were used and the treatments are T₁ -8-HQS 100 ppm + sucrose 4% , T₂ - 8-HQS 200 ppm + sucrose 4%, T₃ – 8-HQS 300 ppm + sucrose 4%, T₄ - GA 100 ppm + sucrose 4%, T₅ - GA 200 ppm + sucrose 4%, T₆ - GA 300 ppm + sucrose 4 and T₇ - Al₂(SO₄)₃ 100 ppm + sucrose 4%, T₈ - Al₂(SO₄)₃ 200 ppm + sucrose 4% + sucrose 8%, T₉ – Al₂(SO₄)₃ 300 ppm + sucrose 4%, T₁₀ - Distilled water (control) without preservative solution and sucrose using Completely Randomized Design with three replications. Each treatment have three flowers with each flower as one replication. Observations on various parameters of postharvest life were recorded on Days taken for spadix necrosis, Days taken for spathe blueing, Physiological loss in weight, Solution uptake and Vase life.

RESULTS AND DISCUSSION

Chemical preservative solutions along with sucrose significantly influenced all the treatments and its performances on prolonging the vase life and quality of cut Anthurium flowers. Among the different chemical preservative solutions (T₈) Al₂(SO₄)₃ 200 ppm + sucrose 4% recorded the maximum results and enhanced the postharvest life and quality of Anthurium cut flowers (Table. 1). Al₂(SO₄)₃ 200 ppm + sucrose 4% delayed flower senescence compared to flowers either held in other treatments. Vase life was significantly increased to 25.48 days (Fig 2). Jowkar, *et al* (2012)

reported that aluminum sulfate treatment significantly increased vase life and improved postharvest visual quality by retaining freshness even at the end of vase life. Sucrose act as a source of energy required for the continuation of the vase life of the cut flowers and also helped for the improvement in the keeping quality value of Anthurium cut flowers. It is well known that sucrose supply increases the longevity of many cut flowers, since sucrose can act as a source of nutrition for tissues approaching carbohydrate starvation, flower opening and subsequent water relations (Kuiper *et al.*, 1995), similar finding were obtained by Lalonde *et al.*, (1999), Nichols (1973), Ichimura, 1998) and Dhaduk *et al.*, (2007). Sucrose act as a source of energy required for the continuation of the vase life of the cut flowers (Halevy and mayak, 1981, Singh and Tiwari. 2002, Reddy, 1988), and may also act as osmotically active molecule, thereby lead to the promoting of subsequent water relations and lengthening their vase life.

Data in Table 1 showed that flowers treated with $\text{Al}_2(\text{SO}_4)_3$ 200 ppm + 4% sucrose recorded the highest relative fresh weight among all treatments. Beginning of the senescence phase in cut flowers is characterized by decrease in fresh weight (Adachi *et al.*, 2000, Liu *et al.*, 2009, Ichimura *et al.*, 1999), aluminium sulphate treatments increased the relative fresh weight. Maximum relative fresh weight was related to T₈ in all days compared to control. $\text{Al}_2(\text{SO}_4)_3$ and sucrose combination had positive effect on solution uptake rate and remaining the flowers fresh weight (Liao *et al.*, 2001). The increased results in (T₈) $\text{Al}_2(\text{SO}_4)_3$ 200 ppm + sucrose 4% with a significant improvement in days taken for gloss loss and vase life of anthurium cut flowers which attained the best result compared to other treatments. Days taken for spadix necrosis as well as days taken for spathe blueing has been increased during the postharvest life as the result of using this combination treatment. $\text{Al}_2(\text{SO}_4)_3$ decreased ethylene production and also highly efficient in reducing bacterial growth in the vase solution and the cut stem ends of anthurium flowers which led to increase in the water uptake of the flower. Sucrose act as a source of energy required for the continuation of the vase life of the cut flowers (Behera, 1993).

Table. 1 Effect of chemical preservatives with sucrose 4% on extending the quality of anthurium cut flowers

Treatments	Days taken for gloss loss	Days taken for spadix necrosis	Days taken for spathe blueing	Solution uptake (ml)	Water loss (ml)
T ₁ – 8-HQS 100 ppm + sucrose 4%	20.32	19.67	23.28	3.25	20.46
T ₂ – 8-HQS 200 ppm + sucrose 4%	20.21	20.12	23.24	3.29	20.81
T ₃ – 8-HQS 300 ppm + sucrose 4%	20.12	19.34	24.04	3.01	21.83
T ₄ – GA 100 ppm + sucrose 4%	20.22	19.93	23.12	3.34	20.04
T ₅ – GA 200 ppm + sucrose 4%	22.05	20.10	23.45	3.21	21.82
T ₆ – GA 300 ppm + sucrose 4%	22.15	21.45	22.32	3.23	20.18
T ₇ – $\text{Al}_2(\text{SO}_4)_3$ 100 ppm + sucrose 4%	21.67	20.54	23.54	3.51	21.11
T ₈ – $\text{Al}_2(\text{SO}_4)_3$ 200 ppm + sucrose 4%	22.23	21.81	24.28	3.62	22.42
T ₉ – $\text{Al}_2(\text{SO}_4)_3$ 300 ppm + sucrose 4%	21.41	19.56	22.77	2.82	21.04
T ₁₀ - Distilled water (control)	14.28	12.84	15.03	1.21	13.29
SE (d)	1.12	0.74	1.15	0.10	1.19
CD (p=0.05)	2.23	1.51	2.21	0.21	2.29

Fig. 1 Effect of chemical preservatives with sucrose 4% on the spadix blackening of Anthurium cut flowers





Fig. 2 Effect of chemical preservatives with sucrose 4% on the Vase life of Anthurium cut flowers



CONCLUSION

Considering the experimental results it can be concluded that the increased results in (T₈) Al₂(SO₄)₃ 200 ppm + sucrose 4% with a significant improvement in days taken for gloss loss and vase life of anthurium cut flowers which attained the best result compared to other treatments. Days taken for spadix necrosis as well as Days taken for spathe blueing has been increased during the postharvest life as the result of using this combination treatment. Al₂(SO₄)₃ decreased ethylene production and also highly efficient in reducing bacterial growth in the vase solution and the cut stem ends of anthurium flowers which led to increase in the water uptake of the flower. Sucrose act as a source of energy required for the continuation of the vase life of the cut flowers.

REFERENCES

- Adachi, M., S. Kawabata, and R. Sakiyama. 2000. Effect of temperature and stem length in carbohydrate content in summer grown chrysanthemums during development and senescence. *Postharvest Biol. Technol.* 20: 63-70.
- Behera, P.K. 1993. Senescence in cut leafy flowering shoot: role of sucrose. *Plant Physiol. and Biochemistry*, New Delhi, 20: 59–59.
- Croat, T.B. 1980. Flowering behaviour of the neotropical genus *Anthurium* (Araceae). *Am. J. Bot.*, 67(6): 888-904.
- Dhaduk, B.K., S. Kumari, A. Singh and J.R. Desai. 2007. Response of gibberellic acid on growth and flowering attributes in anthurium. *Journal of Ornamental Horticulture.*, 10 (3): 187 – 189.
- Halevy, A.H. and S. Mayak. 1981. Senescence and postharvest physiology of cut flowers, part 2. *Hort. Rev.*, 3: 59-143.
- Van Staden, J., J. Upfold and F.E. Dewes. 1994. Effect of sea weed concentrate on growth and development of the marigold (*Tagetes patula*). *J. Appl. Phycol.*, 6: 427-428.
- Ichimura, K., 1998. Improvement of postharvest life in several cut flowers by the addition of sucrose. *Japan Agric. Res. Q.*, 32: 275-280.
- Ichimura, K., K. Kojima and R. Goto. 1999. Effect of temperature, 8-hydroxyquinoline sulphate and sucrose on the vase life of cut rose flowers. *Postharvest Biol. Technol.* 15, 33–40.
- Jowkar, M.M., M. Kafi, M. Khalighi, and N. Hasanzadeh. 2012. Evaluation of aluminum sulfate as vase solution biocide on postharvest microbial and physiological properties of ‘Cherry Brandy’ rose. *Annals of Biological Research*, 3: 1132-1144.
- Kuiper, D., S. Ribots, H.S. Van Reen and N. Marissenn. 1995. The effect of sucrose on the flower bud ripening of ‘Madelon: Cut roses. *Sci. Hort.*, 60: 325-336.
- Lalonde, S., E. Boles. H. Hellmann, L. Barker and J.W. Patrick. 1999. The dual function of sugar carriers: Transport and sugar sensing. *Plant Cell*, 11: 707-726.
- Liao, L. J., Y. H. Lin, K. L. Huang, and W. S. Chen. 2001. Vase life of *Eustoma grandiflorum* as affected by aluminum sulfate. *Bot. Bull. Acad. Sin.*, 42:35-38.
- Liu, J., S. He, Z. Zhang, J. Cao, P. Lv, S. He, G. Cheng and D.C. Joyce. 2009. Nano-silver. pulse treatment inhibit stem end bacteria on cut gerbera cv. Ruikou flowers. *Postharvest Biol. Tech.*, 54: 59-62.
- Lü P., J. Cao; S. He, J. Liu; H. Li, G. Chenga, Y. Dinga and D.C. Joyce. 2010. Nano-silver pulse treatments improve water relations of cut rose cv. Movie Star flowers. *Postharvest Biology and Technology* 57 (2010):66 196–202.
- Nichols, R. 1973. Senescence of cut carnation flowers: respiration and sugar status. *J. Hort. Sci.*, 8: 111-121.
- Nowak, J. and R.M. Rudnicki. 1990. *Postharvest Handling and Storage of Cut Flowers, Florist, Greens and Potted Plants*. 1st Edn., Timber Press, Inc., ISBN-10: 0881921564, pp: 210.
- Reddy, T., C. Nagarajaiah and B. Raju. 1988. Impregnating cut rose stem with nickel increases in vase life. *Hort. Abst.*, 59: 2360.
- Serek, M., E. Sisler and M. Reid. 1996. Ethylene and the postharvest performance of miniature roses. *Acta Hort.*, 424: 145-150.
- Singh, A.K. and A.K. Tiwari. 2002. Effect of pulsing on postharvest life of Rose v. Doris Tystermann. *South Indian Hort.*, 50: 140-144.