

“Studies On The Relationship Between Body Length And Haematocrit In Labeo rohita.”

*Dr. Ravi Kumar Saran Academy, Chapra *Dr. Monalisa S. K. M. University, Dumka *Pooja Kumari, Research Scholar, Jai Prakash University, Chapra *Dr. G. P. Singh H. O. D., P. G. Deptt. of Zoology, Jagdam College, Chapra

Abstract :

The relationship between length and haematocrit in Labeo rohita - Haematological parameters have been recognized as valuable tools for the monitoring of fish health. Blood directly or indirectly takes part in almost all the physiological activities of fishes and is a good indicator of proper health. The external and internal factors have been found to influence the physiology of the body and their health is closely related with their blood parameters. In this study of Labeo rohita, we analyse the relationship between haematocrit and body length (fish size) and found that ht value is increased with increasing length. In male L. rohita ht value increased from mean length 23.875 ± 0.4618 to 34.227 ± 0.2651 but decreased in fishes in the mean length 38.250 ± 0.1682 and again increased in the mean length 44.625 ± 0.4624 Male Labeo rohita showed a higher ht value than females.

Key Words :

Haematological parameters, body length, sex, L. rohita.

Introduction :

Studies of the haematology and blood biochemistry in different species of fish are of comparative physiological interest. They contribute to a greater understanding of habitat, food selection and mode of life.

A sound knowledge of the length – Weight relationship (LWR) of fishes are important in fishery biology because they allow the estimation of the average weight of the fish of a given length group by establishing a mathematical relation between the two (Beyer 1987). LWR has a number of important applications in fish stock assessment . various studies have been done on the length – weight relationship and food habits of fish species [Mortuza and Rahman 2006, Ayoade and Ikulala 2007, Ayoade et al; 2008, Hosseini et al; 2009, Jud et al; 2010, Lawson 2011]. In the present study the ht value increase with increasing fish size upto a certain length and decreased after that.

Materials and Methods :

Live specimens of the Labeo rohita were obtained from various ponds such as pond Inai, Rauza pond, Rajendra sarovar and local market in Chapra and transported in aerated containers to the laboratory. The fishes acclimatize to the laboratory conditions for at least 20 days prior to the experiment in a glass aquarium filled with dechlorinated water. The size of fish varied from 18 to 50 cm length and 100 to 1500 gm in weight. Both sexes were used.

The blood samples obtained from the caudal circulation with the aid of a heparinised 2 cm³ disposable plastic syringe and a 21 gauge disposable hypodermic needle.

Haematocrit (Ht/PCV) was determined by microhaematocrit centrifugation technique.

The haematocrit value or the packed cell volume were estimated by centrifuging it for 5 minutes at 10,000 rotation per minute (rpm).

Differences in haematological parameters between male and female fish were statistically analyzed by student's t – test.

Table – 1

Total length and blood haematocrit value of male *Labeo rohita*; BL. Body length groups (in cm); N. No. of fishes; ML. Mean length (in cm \pm SE); Ht. Haematocrit value (% \pm SE).

| BL | N | ML | Ht |
|---------------|----|---------------------|---------------------|
| 19.00 – 28.70 | 40 | 23.875 \pm 0.4618 | 35.036 \pm 0.1540 |
| 28.70 – 31.60 | 30 | 30.150 \pm 0.1607 | 39.407 \pm 0.1766 |
| 31.60 – 36.80 | 35 | 34.227 \pm 0.2651 | 41.651 \pm 0.0898 |
| 36.80 – 39.70 | 30 | 38.250 \pm 0.1682 | 35.865 \pm 0.3635 |
| 39.70 – 49.50 | 40 | 44.625 \pm 0.4624 | 38.538 \pm 0.4487 |

Table – 2

Total length and blood haematocrit value of female *Labeo rohita*; BL Body length groups (in cm); N. No. of fishes; ML. Mean length (in cm \pm SE); Ht. Haematocrit value (% \pm SE).

| BL | N | ML | Ht |
|---------------|----|----------------------|---------------------|
| 19.00 – 28.70 | 40 | 23.875 \pm 0.4618 | 33.726 \pm 0.1339 |
| 28.70 – 31.60 | 30 | 30.150 \pm 0.1607 | 37.445 \pm 0.2336 |
| 31.60 – 36.50 | 50 | 34.0500 \pm 0.2062 | 39.827 \pm 0.0539 |
| 36.50 – 39.50 | 30 | 37.950 \pm 0.1607 | 34.814 \pm 0.3111 |
| 39.50 – 49.25 | 40 | 44.375 \pm 0.4621 | 36.145 \pm 0.3297 |

Table – 3

Blood haematocrit value of *Labeo rohita* of different sexes and different body length groups: BL. Body length groups (in cm.); Ht. Haematocrit value (% \pm SE); N. Number of fishes; M. Male; F. female.

| BL | N | | Ht | | t.test | P | df |
|---------------|----|----|---------------------|---------------------|---------|--------|----|
| | M | F | M | F | | | |
| 19.00 – 28.70 | 40 | 40 | 35.036 \pm 0.1540 | 33.725 \pm 0.1339 | 6.4218 | < 0.05 | 77 |
| 28.70 – 31.60 | 30 | 30 | 39.406 \pm 0.1766 | 37.445 \pm 0.2336 | 6.6832 | < 0.05 | 54 |
| 31.60 – 36.80 | 35 | 50 | 41.651 \pm 0.0898 | 39.827 \pm 0.0539 | 17.4072 | < 0.05 | 58 |
| 36.80 – 39.70 | 30 | 30 | 35.865 \pm 0.3635 | 34.814 \pm 0.3111 | 2.1967 | < 0.05 | 57 |
| 39.70 – 49.50 | 40 | 40 | 38.538 \pm 0.4487 | 36.145 \pm 0.3297 | 4.2967 | < 0.05 | 72 |

Results & Discussions :

In the present study the haematocrit value in male *Labeo rohita* in different length groups viz., 19.00 – 28.70 cm, 28.70 – 31.60cm, 31.60 – 36.80 cm, 36.80 – 39.70 cm, 39.70 – 49.50 cm. were found to be 35.036 \pm 0.1540, 39.407 \pm 0.1766, 41.651 \pm 0.0898, 35.865 \pm 0.3635, 38.538 \pm 0.4487 respectively (table – 1).

The haematocrit value in female *Labeo rohita* in different length groups viz., 19.00 – 28.70 cm, 28.70 – 31.60 cm, 31.60 – 36.50 cm, 36.50 – 39.50 cm, 39.50 – 49.25 cm were found to be 33.726 ± 0.1339 , 37.445 ± 0.2336 , 39.827 ± 0.0539 , 34.814 ± 0.3111 , 36.145 ± 0.3297 respectively (Table – 2).

The haematocrit value increased with increasing length. In male *Labeo rohita* haematocrit value increased from mean length 23.875 ± 0.4618 to 34.227 ± 0.2651 but decreased in fishes in the mean length 38.250 ± 0.1682 and again increased in the mean length 44.625 ± 0.4624 .

In female *Labeo rohita* the haematocrit value also increased with increasing length. The haematocrit value in female *Labeo rohita* increased from mean length 23.875 ± 0.4618 to 34.0500 ± 0.2062 but decreased in fishes in the mean length 37.950 ± 0.1607 and again increased in the mean length 44.375 ± 0.4621 .

In the present study high haematocrit value in the above mentioned length and weight groups of fish was due to high physical activity. This is in agreement with results from other fish species. *Tilapia zilli*, Ezzat et al, (1973); *Cyprinus carpio*, Fourie and Hatting (1976); *Cyprinion macrostomus*, Al – Mehdi and Khan (1984); *Amphiprourus cuchia*, Banarjee (1986); *Barbus xanthoptrus*, Hameed et al. (1985); *Anguilla anguilla*, Johnson et al (1974); *Clarius batrachus*, Joshi et al (1977); *Sarotheriodon mossambica*, Chaudhary et al (1986).

Comparative ht value in male and female sex of *Labeo rohita* in Table – 3 clearly showed higher ht value in male as compared to females.

The difference in haematocrit value between the two sexes might be genetically determined and also due to the higher metabolic rates of males compared to females. My finding was as per findings of Fouri & Hattingh (1976); Raizada et al (1983); Chaudhary et al. 1986, and Jawed et al, 2004.



References :

1. Al – Mehdi and Khan (1984) Fresh water carp. *Cyprinon macrostomus* from northern Iraq. *Envirn. Ecol.* ,(3): 222-226.
2. Ayoade and Ikulala (2007) Length weight relationship condition factor and stomach contents of *Hemichromis bimaculatus* , *Sarotherodon melanotheron* and *Chromidotilapia guentheri* (perciformes : Cichlidae) in Eleiyele Lake, South western Nigeria. *Rev. Bio. Trop.*, 55 : 969-977.
3. Banerjee, S. and Banerjee M. (1988)- Erythrocyte related Blood parameters in Rita Rita (Hamilton) with special references to body length, sex and season. *Proc. 75th and Sc. Cong. Part 3 pp.* 50-51
4. Beyer, J.E. (1987)- On length- Weight relationships. Part 1. Computing the main weight of the fish of a given length fish Byte Manila, 5:11-13.
5. Chaudhari et al. (1986)- Size & sex Related variation of some blood parameters of *sarotheriodon mossambica*, *Environ, Ecol.*, 4(i): 61-63
6. Ezzat et al. (1973)- Studies on the blood characteristics of *Tilapia Zilli* (gervasis). 1. Blood cells. *J. Fish Biol.*, 6: 1-12.
7. Fourie and Hatting (1976)- A seasonal study of the haemobiology of carp, *cyprinus carpio* from A locality in the Transvaal, South Africa. *Zool. Afri.* 11(i): 75-80.
8. Hameed et al. (1985)- The relationship between age and Blood constituents of *Barbus Xanthoptrus* (Heckel) in the different seasons from Sadat Al Hindia, Iraq, *J. Biol. Res. Center*, 17: 77-86.
9. Hosseini et al. (2009)- Feeding and Spawning of *sphyraena jello* in The North-West of Persian Gulf. *J. fish. Aquatic Sci.*, A : 57-62
10. Jawed et al. 2004- The relationship between haematocrit and biological parameters of the Indian shad *Tenualosa illisha* (family clupeidae). *Animal Biodiversity and conservation* 272: 47-52.
11. Johanson et al. (1974)- Metabolic and haematological studies on the yellow and silver phases of the European ell. *Anguilla Anguilla L. (iii) haematology Comp. Biochem, Physiol.* 47 B : 593-599.
12. Lawson, E.O., (2011)- Length-Weight relationships and Fecundity estimates in mudskipper, *Periophthalmus Papillo* (Bloch and Schneider 1801) caught from the mangrove swamps of Lagos lagoon, Nigeria. *J. fish, Aquat sci.*, 6 : 264-271.
13. Mortuza, M. g. and T. Rahman (2006)- Length-Weight relationship, condition factor and sex – ratio of fresh water fish, *rhinomugil corsula* (Hamilton) 9 *Mugliformes : Mugillidae*) from Rajshahi, Bangladesh. *J. Bio. Sci.*, 14 : 139-141.
14. Raizada et al. (1983)- Monthly Variations in The haematocrit values (PCV) in a teleost, *cirrhinus mrigala* (Ham) *comp. Physiol. R. Col.* 8 : 190-196.