

PHYTOREMEDIATION POTENTIAL OF METHYLENE BLUE AND METHYL ORANGE DYE (AZO DYE) BYE *EICHHORNIA CRASSIPES* PLANT.

CHIRAG SHAH¹, LINZ- BUOY GEORGE²

¹Departemnt Of Environmental science, School of science, Gujarat University, Ahmedabad, Gujarat, India.

²Department of Zoology, School of science, Gujarat University, Ahmedabad, Gujarat, India.

Abstract : Phytoremediation is defined as the use of plants and their associated microorganisms to remove harmless pollutants from contaminated sites. It is a promising approach in the treatment of dye wastewater due its cost-effectiveness. The aim of this study was to determine the effectiveness of water hyacinth (*Eichhornia crassipes*) in removing color. Water hyacinth was used to treat 50 mg/L of methylene blue (MB) and 50 mg/L of methyl orange (MO) for 20 days under ambient temperature (30±1 °C). The pH of the synthetic dye wastewater was observed throughout 20 days. Results obtained showed that percentage of color removal was higher for MB compared to MO which were 97.12% and 67.00% respectively. The relative growth of *Eichhornia crassipes* in MB and MO were also being determined. The cell structure of *Eichhornia crassipes* (leaf, shoot and root) before and after the plants were exposed to dye wastewater was analysed using light microscope.

Index Terms—*Eichhornia crassipes*, methylene blue, methyl orange, phytoremediation

I. Introduction

Dyes can give color to water bodies even when they exist in small amount. They are widely used in various industries such as textile, plastic, paper and rubber industries [1]. Among these industries, textile industry ranks first in the usage of dyes for coloration of fiber. The textile wastewater is well known to contain strong color, large amount of suspended solids, high fluctuation in pH, high temperature, high COD concentration and other organic contents [2]. Due to the usage of dye and pigments during the dyeing process, the strong color and turbidity of the textile wastewater effluents caused many problems because of its negative visual impact [3]. Some of the dyes are toxic and carcinogenic in nature. Methylene blue (MB) is a basic dye which is used extensively in the dyeing and printing of cotton, silk etc [4]. MB can have various harmful effects. The high concentration of this dye in contact with the eye can cause corneal injury in human beings. Doses in the range 500 mg can lead to anemia, dizziness, headache, abdominal pain, nausea, profuse sweating and mental confusion [5]. On the other hand, methyl orange (MO) is an example of azo dye. Azo dyes are the most versatile and the largest group of dyes. They are considered to be toxic and mutagenic to living organisms [6]. Their discharge into a water environment can cause serious health problems as well as acute and chronic effects on aquatic life [7]. Thus, the impact and treatment of the dye to the environment has been studied by many researchers in the past decades. Many treatment methods have been developed for the removal of dyes from wastewater, including adsorption, oxidation processes, microbiological or enzymatic decomposition, and etcetera [8]. However, phytoremediation proves to be quite an efficient method in comparison to these methods [8]. Phytoremediation is the use of plants for environmental cleanup. It is popular because it has a lot of advantages such as cost-effectiveness, aesthetic advantages, long-term applicability and it can be directly applied at the polluted sites where other treatment methods are too expensive [9]. In addition, plants offer protection against wind and water erosion, preventing contaminants from spreading. It is easier to manage because of its autotrophic system that has large biomass and require little nutrient input [9]. Plants that are fast growing, with deep and fibrous roots, such as grasses, are very useful in phytoremediation [9], [10]. There are five types of phytoremediation techniques, namely phytoextraction, phytostabilization, phototransformation, phytodegradation, and rhizofiltration [11]. Water hyacinth (*Eichhornia crassipes*) is a free floating aquatic plant that originated from tropical South America and is now widespread in all tropical climates. It is listed as one of the most productive plants and also one of the world's worst aquatic plants. Due to its fast growth rate, large biomass production, high tolerance to pollution and its heavy metal and nutrient absorption capacities, *Eichhornia crassipes* has a potential to cleanup various types of wastewaters [12].

II. METHODOLOGY

A. Preliminary Study

Eichhornia crassipes were collected from Tasik Harapan, Universiti Sains Malaysia. All the plants were washed thoroughly with tap water to remove any dirt and soil particles adhered to the plants. The plants were then grown in basins using tap water for one week to allow the plants to adapt to the new environmental conditions. After one week, the plants were introduced to 6 L of MB and 6 L of MO with different concentrations (50 mg/L, 100 mg/L, 150 mg/L, 200 mg/L) for 7 days, to determine the most effective concentration for color removal. The conditions of the plants were observed throughout the experimental works.

B. Experimental Study

Based on the preliminary study, concentrations of 50 mg/L of MB and 50 mg/L MO were chosen and prepared to grow the plants. The plants were wiped dry to remove excess water. Plants of almost the same size, which weighted between 210 g to 240 g and height within 15 cm to 20 cm were put into the basins which contained 6 L of 50 mg/L MB and 6 L of 50 mg/L MO dye aqueous solution. Experiments were carried out in triplicates with one control set each. The plants were left to grow for 20 days in open air and exposed to sufficient sunlight. Any decrease in volume of solution in each basin was added with deionized water to the mark to ensure their volume were maintained at 6 L to counter water loss due to evapotranspiration. 90 mL of samples were taken every 2 days.

The absorbance of MB and MO were measured by using spectrophotometer with λ_{max} 665 nm and 465 nm respectively. Equation 1 was used to calculate the percentage of dye removal.

$$\text{Percentage of dye removal (\%)} = [(C_o - C_f) / C_f] \times 100 \quad (1)$$

where, C_o is the initial dye concentration (mg/L) and C_f is the final dye concentration (mg/L).

The plant growth assessment was done by monitoring the wet weight and length of the plants before and after the experiment. The relative growth rate of plant was calculated as in (2) [13],

$$\text{relative growth rate (day}^{-1}\text{)} = (\ln w_2 - \ln w_1) / t \quad (2)$$

where, w_1 is the wet weight of *Eichhornia crassipes* before exposure to dye contaminant, w_2 is the final wet weight of *Eichhornia crassipes* after exposure to dye contaminant, and t is the duration (day).

III. RESULTS AND DISCUSSION

A. Variation of pH

The pH can affect the efficiency of color removal. It has been reported that *Eichhornia crassipes* can grow within a pH range of 4.4 to 10.0 in different water stream [14]. Therefore, the pH value of MB and MO must be well-monitored. Fig. 1(a) and 1(b) show the pH readings of MB and MO for 20 days. The pH value of MB and its control set were within the range of pH 4 to pH 6 throughout the experiment works. However, the MO and its control set fluctuated between pH 5 and pH 8. The fluctuation of the pH value is due to the free diffusion of the carbon dioxide with the atmosphere and the absorption of nutrient by the plants [15].




Fig. 1. pH reading of (a) MB and (b) MO for 20 days.

B. Color Removal of MB and MO

The percentages of color removal were calculated for MB and MO remediated by *Eichhornia crassipes*. Fig. 2(a) and 2(b) show the percentages of the color removal of MB and MO by *Eichhornia crassipes* after 20 days. From Fig. 1(a) and 1(b), the percentage of color removal of MB is higher than the percentage of color removal of MO which is 97.12 % and 67.00% respectively. This is because MO has a higher molecular weight and the structure of MO contains azo bond (-N=N-) as compared to MB which has a lower molecular weight. Many researchers have claimed that the rate of color removal is also dependent on the dye class rather than the molecular features [16]. The color removal of the control set is due to the photodegradation of dye molecules as the basins were exposed to sunlight. Photodegradation of dye occurs even in the presence of ultraviolet light only [17].




Fig. 2. Color removal of (a) MB and (b) MO for 20 days.

C. Relative Growth Rate of *Eichhornia crassipes*

Relative growth rate of the plant was investigated to determine the ability of the plants to grow and survive in the contaminated wastewater. Table I shows the relative growth rate of *Eichhornia crassipes* in MB and MO dye aqueous solution. The results show that all of the plants had increased in wet weight after exposure to the dye contaminants. The wet weight increases from 223.1 g to 319.8 g and from 241.9 g to 387.5 g in MB and MO dye aqueous solution respectively. The relative growth rate of *Eichhornia crassipes* in MO dye aqueous solution is higher than it in MB which is 0.03 day⁻¹ and 0.02 day⁻¹ respectively. This indicates that plants grew better in MO dye aqueous solution than in MB dye aqueous solution.

TABLE I: THE RELATIVE GROWTH OF *EICHHORNIA CRASSIPES* IN MB AND MO DYE AQUEOUS SOLUTION

Dye	Wet weight of plant (g)		Relative Growth (day ⁻¹)
	Before	After	
MB	223.1	319.8	0.02
MO	241.9	387.5	0.03

D. Average Length of *Eichhornia crassipes* after 20 Days

Table II shows the average length of the plant before and after 20 days in MB and MO dye aqueous solution. The results showed that all the plants had an increase in their length after 20 days. The average increase are 1.6 cm and 3.1 cm when the plants exposed to MB and MO dyes aqueous solution respectively. However, the plants in MB and MO dye aqueous solution started to show wilting symptoms after 18 days of the experimental works. Wilting is a biochemical changes that prevent the healthy tissues of the plants from damage by surrounding the damaged area with a high concentration of phenolic compounds [18]. The overall results indicated that *Eichhornia crassipes* has a potential to be used in phytoremediation since it is highly tolerant to the pollution and can be used in wastewater treatment in ponds [8]. Aqueous solution and methyl orange dye aqueous solution showed wilting symptoms after 18 days of exposure. The observation of the plant cells using light microscope analysis proved that the roots of *Eichhornia crassipes* can absorb the dyes. Thus, it can be concluded that *Eichhornia crassipes* is a suitable plant to be used in the phytoremediation of dye wastewater and has a potential to be used in water bodies contaminated with dyes.

REFERENCES

- [1] S. M. Kanawade and R. Gaikwad, "Removal of methylene blue from effluent by using activated carbon and water hyacinth as adsorbent," *Int. J. Chem. Eng. & Appl.*, vol. 2, no. 5, pp. 317-319, October 2011.
- [2] S. H. Lin and C. L. Lai, "Kinetic characteristics of textile wastewater ozonation in fluidized and fixed activated carbon beds," *Water Res.*, vol. 34, no. 3, pp. 763-772, February 2000.
- [3] I. Sengil, M. Ö zacar, and B. Ö MüRLü, "Decolorization of CI reactive red 124 using the electrocoagulation method," *Chem. Biochem. Eng. Quarterly*, vol. 18, no.4, pp. 391-401, 2004.
- [4] K. Dutta, S. Mukhopadhyay, S. Bhattacharjee, and B. Chaudhuri, "Chemical oxidation of methylene blue using a fenton-like reaction," *J. Hazard. Mater.*, vol. 84, no. 1, pp. 57-71, June 2001.
- [5] V. Ponnusami, S. Vikram, and S. N. Srivastava, "Guava (psidium guajava) leaf powder: Novel adsorbent for removal of methylene blue from aqueous solutions," *J. Hazard. Mater.*, vol. 152, no. 1, pp. 276-286, March 2008.
- [6] A. Stolz, "Basic and applied aspects in the microbial degradation of azo dyes," *Appl Microbiol. Biotechnol.*, vol. 56, no. 1-2, pp. 69-80, July 2001.
- [7] R. V. Khandare, A. N. Kabra, D. P. Tamboli, and S. P. Govindwar, "The role of aster amellus linn. in the degradation of a sulfonated azo dye remazol red: A phytoremediation strategy," *Chemosphere*, vol. 82, no. 8, pp. 1147-1154, February 2011.

- [8] V. Muthunarayanan, M. Santhiya, V. Swabna, and A. Geetha, "Phytodegradation of textile dyes by water hyacinth (*eichhornia crassipes*) from aqueous dye solutions," *Int. J. Env. Sci.*, vol. 1, no. 7, pp. 1702-1717, July 2011.
- [9] C. Cluis, "Junk-greedy Greens: Phytoremediation as a new option for soil decontamination," *BioTeach. J.*, vol. 2, pp. 61-67, 2004.
- [10] R. V. Khandare, A. N. Kabra, D. P. Tamboli, and S. P. Govindwar, "The role of aster amellus linn. in the degradation of a sulfonated azo dye remazol red: a phytoremediation strategy," *Chemosphere*, vol. 82. no. 8, pp. 1147-1154, February 2011.
- [11] M. Vidali, "Bioremediation. An overview," *Pure Appl. Chem.*, vol. 73, pp. 1163-1172, August 2001.
- [12] A. Malik, "Environmental challenge vis a vis opportunity: the case of water hyacinth," *Env. Int.*, vol. 33, no. 1, pp. 122-138, January 2007.
- [13] A. Khataee, A. Movafeghi, S. Torbati, S. S. Lisar, and M. Zarei, "Phytoremediation potential of duckweed (*lemna minor* L.) in degradation of ci acid blue 92: artificial neural network modeling," *Ecotox Environ. Safety*, vol. 80, pp. 291-298, June 2012.
- [14] A. El-Gendy, N. Biswas, and J. Bewtra, "Growth of water hyacinth in municipal landfill leachate with different pH," *Environ. Tech.*, vol. 25, no. 7, pp. 833-840, July 2004.
- [15] W. T. Haller and D. Sutton, "Effect of pH and high phosphorus concentrations on growth of water hyacinth," *Hyacinth Control Journal*, vol.11, pp. 59-61, 1973.
- [16] C. I. Pearce, J. R. Lloyd, and J. T. Guthrie, "The removal of colour from textile wastewater using whole bacterial cells: A review," *Dyes and Pigments*, vol. 58, no. 3, pp. 179-196, September 2003.
- [17] N. Daneshvar, D. Salari, and A. R. Khataee, "photocatalytic degradation of azo dye acid red 14 in water on ZnO as an alternative catalyst to TiO₂," *J. Photochem. Photobio. A: Chem.*, vol. 162, no. 2-3, pp. 317-322, March 2004.
- [18] S. Nilratnisakorn, P. Thiravetyan, and W. Nakbanpote, "Synthetic reactive dye wastewater treatment by narrow-leaved cattails.

