

# IMPACT ON FISH DIVERSITY IN LOWER GANGETIC DELTA DUE TO PRAWN SEED COLLECTION

<sup>1</sup>Maria Fernandes, <sup>2</sup>Goutam Sengupta, <sup>3</sup>Nabonita Pal & <sup>4</sup>Abhijit Mitra

<sup>1,2</sup>School of Management, Techno India University, West Bengal, Kolkata-700091, India

<sup>3</sup>Department of Oceanography, Techno India University, West Bengal, Kolkata-700091, India

<sup>4</sup>Department of Marine Science, University of Calcutta, 35 B.C. Road, Kolkata 700019, India

**Abstract:** West Bengal is a maritime state in the north-eastern part of Indian sub-continent adjacent to Bangladesh, with a coastal area of 10,158.22 sq. km. The coastal zone comprises of the famous mangrove ecosystem of Indian Sundarbans, which can be divided into various transects on the basis of salinity. The practice of wild harvest of tiger prawn seeds in the present geographical locale is the greatest ecological threat in terms of magnitude as a large variety of fish juveniles found damaged during the screening process. The present paper deals with the juvenile finfish count in three different salinity segments of the coastal ecosystem with particular emphasis on their spatial and temporal variations for a span of ten years (January, 2009 to December, 2018). This data bank has been generated from the wasted finfish juvenile samples that are thrown away after collection and subsequent segregation of tiger prawn seeds from the coastal waters. The ten years data revealed a sequential uniformity in all the selected stations, the diversity being maximum in the high saline zone and minimum in the upper stretch of the estuary where the salinity of the aquatic phase is lowest. Some conservation measures, in the form of alternative livelihood schemes have also been suggested at the end that can divert the tiger prawn seed collectors from this destructive practice encouraging positive conservation of fish germplasm in and around the Indian Sundarbans mangrove ecosystem.

**Keywords - Indian Sundarban, prawn seed collection, women shrimp fry catcher**

## I. INTRODUCTION

Ichthyoplankton and fish juveniles are the major biotic components of the world ocean [1] and play a vital role in regulating the aquatic food chain. They are widely distributed in the brackish water system. Mangrove dominated coastal zone owing to the presence of mangrove vegetation derived nutrients and congenial environmental conditions for their growth and survival. It is reported that the mangrove ecosystem serves as the nursery of a large spectrum of fish juveniles [2,3,4,5]. This unique ecosystem forms an ideal ecological reservoir for the pelagic community because the production of leaf litter and detrital matter from mangrove plants fulfills the nutritional requirements of prawn juveniles, crab megalops and juveniles of several species of osteichthyes. Due to this fact, the mangrove ecosystem is recognized as the world's most potential nursery and breeding ground [4].

The Indian Sundarbans lie at the apex of the Bay of Bengal and sustain the famous mangrove ecosystem of the tropics. The islands, its associated waters, creeks and estuaries of the deltaic lobe are the nurseries for many coastal and oceanic species of fish that support important subsistence and commercial fisheries in the northern Bay of Bengal. The diverse vegetation of the Sundarbans forests includes about 36 true mangrove species that provide a unique mix of habitats for wildlife, which includes tiger, deer, wild boar, monkey, turtles, terrapins, dolphins, estuarine crocodiles and several species of birds, most of which are migratory. This region is considered as one of the most backward regions in the maritime state of West Bengal and is home to about 4.2 million people, of which 56% are landless. Literacy rates a below 35%; most communities do not have electricity or safe drinking water and agricultural productivity is lower than the state average. Inadequate infrastructure, poor communication facilities, lack of access to clean drinking water, health and education services with a fragile, limited natural resource base that have contributed to a low level of development and high poverty existence in the region.

A diminishing natural resource base is threatening the ecological integrity of Indian Sundarbans and the livelihood of the inhabitants. At present, the major environmental-cum-socioeconomic issue in this mangrove ecosystem is the proliferation of the shrimp farms. Due to the absence of any hatchery in the entire state, the supply of tiger prawn seeds to these farms are done through the wild collection from the estuarine and coastal waters. Thousands of women and children collect wild tiger prawn seeds by employing nets of a particular mesh size to haul in the drifting community, irrespective of the tides, and throw away the major portion of the haul (containing the juveniles of finfish and shellfish) after sorting out the post larvae (or seeds) of tiger prawn. This practice results in a great loss of pelagic biodiversity (in terms of stock) which might cause an adverse effect on the delicate ecological balance of the system [6,7]. A negative impact also becomes obvious in the prospect of demersal fisheries of the state of West Bengal due to this indiscriminate destruction of the various species of fishes at their juvenile stages. With this background, the present paper aims to highlight the magnitude of finfish juvenile diversity loss in different seasons of the year during 2009-2018. This occupation has spread its root to such an extent that about 95% of the coastal population living below poverty line are engaged with this destructive activity. It is very difficult to implement a total ban on this activity, as these huge masses will face

extreme economic marginalization. So, our present paper suggests few alternative livelihood schemes, which may establish a harmony between the biological diversity and socio-economic profile of the area.

## II. DESCRIPTION OF THE STUDY AREA

West Bengal being a maritime state in the northeast part of the Indian sub-continent adjacent to Bangladesh. The coastal zone of this state spreads over an area of 10,158.22 sq. km. and is restricted within the latitude 21°30'N to 22°30'N and longitude 87°25'E to 89°10'E. The river Harinbari or Heronbhanga of the Indian Sundarbans (India- Bangladesh border) is the eastern most border while the New Digha coast of Orissa-West Bengal border constitutes the western boundary of coastal West Bengal. With considerable degree of marine characteristics in the major portion of the ecosystem, the important morphotypes of coastal West Bengal are sandflats, coastal dunes, beaches, mudflats, estuaries, creeks, inlets and mangrove flats. There is a drastic variation of salinity and dilution factor between different horizontal transects of the ecosystem. The dilution factor, salinity and pH reveal significant variations at the same time in different locations of the coastal zone [3]. The recent emergence of Haldia industrial complex and various fish landing stations in the coastal zone has also opened the gateway of input of various categories of wastes in this ecosystem.

Geographically, the study area encompasses three major districts of the state of West Bengal namely, 24 Parganas (North), 24 Parganas (South) and Medinipur (East). The Indian Sundarbans declared as the World Heritage Site by UNESCO in the year of 1989 fall within the North and South 24 Parganas districts. Three sampling stations were selected in and around this deltaic ecosystem namely, Diamond Harbour (station 1), Sagar light house (station 2) and Junput (station 3). Each of these sampling stations is markedly different from the other with respect to aquatic salinity due to their location and proximity to the Bay of Bengal.

Diamond Harbour (station 1) is situated in the low saline upper stretch of Hooghly estuary, just outside the northern boundary of Indian Sundarbans. The station is very near to the Haldia port-cum-industrial complex. Salinity of surface water is minimum around the station owing to its location far away from the Bay of Bengal in the extreme upstream region and also due to huge fresh water discharge from the Hooghly river, which is perennial in nature. The station has three mangrove species along with few mangrove associates like *Porteratia coarctata* (salt marsh grass) and seaweeds (*Enteromorpha* sp.).

Sagar light house (station 2) is situated in the south western tip of the Sagar Island and falls in the western sector of Indian Sundarbans. The station has rich mangrove vegetation and extensive mudflats. Although there are no industrial activities in this station, but the presence of sizeable number of shrimp farms (presently carrying on shrimp culture by traditional method with a very low stocking density of prawn seeds) has enriched the surrounding water with nutrients and organic load.

Junput (Station 3) is situated in the Medinipur (East) district of coastal West Bengal, which is noted for its high aquatic salinity owing to its proximity to the Bay of Bengal. The extremely high salinity has posed an inhibitory effect on the growth and survival of mangroves in this station [8]. Existence of salt pans has made the soil of the area hyper saline in nature. Although the station has no industry around its vicinity, but the presence of Digha tourist center and Shankarpur fishing harbour close to the station has multiplied the anthropogenic pressure around the zone.

In all these stations, tiger prawn seeds collection is a very common scene in which the coastal and estuarine waters are screened in search of target species (seeds of *Penaeus monodon*) by using nets of very fine mesh size. This activity traps a varied spectrum of several finfish juveniles (non-target species), which are subsequently thrown away thus destroyed.

## III. MATERIALS AND METHODS

Finfish juvenile samples were collected monthly during January 2009 to December 2018 from tiger prawn seed collectors. Collection of 10 nets were randomly mixed and a constant weight of 10 gm were preserved in 4% formaldehyde. The samples were identified from Zoological Survey of India and its count was done using hand lens. Triplicate counts were done to derive the maximum accuracy level and the mean value of the species count, was finally used to enumerate the community structure of finfish juvenile in the study area through computation of Shannon Weiner Species Diversity Index [9] as per the AAMSTAT software developed by BOOLEAN LOGIC PRIVATE LIMITED in 2005.

## IV. RESULT AND DISCUSSION

### A. Root Cause of Tiger Prawn Seed Collection

1. **Mushrooming of shrimp culture activities:** The aquatic sub-subsystem of coastal West Bengal is highly congenial for the culture of tiger prawn not only because of the aquatic salinity, tidal flushing (which favours the process of water exchange), but also because of the rich phytoplankton diversity (102 species documented so far) in the optimum level of aquatic nutrients [10]. About 33 000 ha in North 24 Parganas and 12 000 ha in South 24 Parganas districts have been devoted for shrimp culture activities, which have a high demand for tiger prawn seeds for economic viability. This

demand is mitigated through wild harvest of PL<sub>20</sub> (larval stage 20 days from the hatching date) from estuaries, coastal wetland and brackish water creeks [11,12,13].

- Absence of hatchery:** Currently, there are few operational shrimp nurseries in West Bengal, but no ideal hatchery in real sense (i.e. spawning of adult to produce nauplii). Low salinity is the biggest problem for shrimp hatchery in West Bengal coast, particularly in Sundarbans. In the absence of any hatchery, the demand for tiger prawn seeds has become very acute and the entrepreneurs have no option, but to depend on wild harvest of tiger prawn seeds.
- Socio-economic profile:** The socio-economic profile of coastal West Bengal is not very encouraging, particularly, in Indian Sundarbans, it is highly distressed. Recently, the population has touched the figure of 4.2 million, which is predominated by farming and fishing community. More than 90% of the farming community is either small or marginal farmers and about 50% survives below the poverty level. This sizable chunk of population engage themselves in screening the estuarine water from dawn to dusk in search of the target species to earn immediate cash to them. The price of tiger prawn seeds varies from Rs. 200 per 1000 to Rs. 800 per 1000 depending on the season and demand [14].

## B. Community Structure of Finfish Juvenile

The coastal zone of West Bengal is one of the most biologically productive, taxonomically diverse and aesthetically celebrated ecotone of the country, which is sustaining the famous mangrove ecosystem of Indian Sundarbans [15]. The pelagic system around the coastal zone is highly productive due to considerable concentrations of nutrients derived from anthropogenic origin, run-off from the adjacent landmasses, litters and detritus of mangrove vegetation, etc.

**Table 1:** Monthly average Standing Stock/fish juvenile count (N) of finfish juvenile species (H) of three selected sampling stations during Jan'94 to Dec'03

Year	Monthly average Standing Stock (N) of finfish species								
	Stn 1			Stn 2			Stn 3		
	Prm	Mon	Pom	Prm	Mon	Pom	Prm	Mon	Pom
2009	392	161	245	1598	134	522	1593	260	767
2010	321	133	291	1315	122	422	1303	213	625
2011	200	82	183	1273	99	416	1290	213	631
2012	195	79	175	1279	99	413	1229	201	596
2013	215	89	197	1090	84	361	1055	172	651
2014	217	89	198	716	56	233	679	123	333
2015	224	93	204	639	51	209	601	103	291
2016	221	92	201	545	42	180	521	89	255
2017	225	93	204	910	178	442	934	153	440
2018	389	38	67	936	73	304	1020	79	367

Table 1 exhibits the monthly average of Standing Stock (N) and Shannon Weiner Species Diversity Index (H) of finfish juvenile species in the three sampling stations. Members of genus *Coilia* sp., *Thryssa* sp., *Mugil* sp., *Tenualosa* sp., *Liza* sp., *Scatophagus* sp., *Stolephorus* sp., *Cynoglossus* sp., *Sillago* sp. are dominant (in terms of standing stock or biomass) in all the stations (Table 2) and in all the months of the study period, which suggests the wide range of tolerance of this genus in the present geographical locale. The study clearly depicts the regulation of finfish juveniles by aquatic salinity. Similar observations were also recorded by earlier workers in this study area [16,17,18,19,20]. The regulatory role of salinity on finfish juvenile species composition and abundance was also confirmed in a South African case study [21]. The diversity of finfish juvenile is highest at station 3 (Junput) which is followed by station 2 (Sagar light house) and station 1 (Diamond Harbour). The highest values of H and N at Junput confirm the positive role of aquatic salinity in maintaining the fish germplasm in the study area. Diamond Harbour is an extremely low saline zone in the upstream area with an average aquatic salinity of 1.7‰ during premonsoon, 0‰ during monsoon and 1.3‰ during postmonsoon [22] and hence chance of getting stenohaline fish juvenile species is rare. Irrespective of significant variation of species number and diversity (H) amongst the selected stations, the present geographical locale is an abode of several species of finfish and shellfish and nursery of fish juveniles [16]. Destruction of this juvenile stock may therefore create an adverse impact on the pelagic and demersal fish production status of the state in near future [6].

**Table 2:** Distribution of finfish juvenile species in 10 gm composite wasted sample collected from three selected sampling stations during January, 2009 to December, 2018

Sl. No	Species	Station 1	Station 2	Station 3
1.	<i>Coilia</i> sp.	+	+	+
2.	<i>Thryssa hamiltonii</i> (Gray)	+	+	+

3.	<i>Thryssa baelama</i>	+	+	+
4.	<i>Torquigener oblongus</i>	+	+	+
5.	<i>Rhinomugil corsula</i>	+	+	+
6.	<i>Mugil cephalus</i>	+	+	+
7.	<i>Sillaginopsis panijus</i>	+	+	+
8.	<i>Sillago sihama</i>	+	+	+
9.	<i>Sillago soringa</i>	+	+	+
10.	<i>Zenarchopterus dispar</i>	+	+	+
11.	<i>Glossogobius quiris</i>	+	+	+
12.	<i>Pseudapocryptes lanceolatus</i>	+	+	+
13.	<i>Eupleurogrammus glossodon</i>	+	+	+
14.	<i>Pseudorhombus sp.</i>	+	+	+
15.	<i>Pisodonophis boro</i>	+	+	+
16.	<i>Tenualosa ilisha</i>	+	+	+
17.	<i>Cynoglossus arel</i>	+	+	+
18.	<i>Cynoglossus sp.</i>	+	+	+
19.	<i>Leiognathus blochii.</i>	-	+	+
20.	<i>Leiognathus equalus</i>	+	+	+
21.	<i>Hilsa sp.</i>	+	+	+
22.	<i>Scatophagus argus</i>	+	+	+
23.	<i>Liza parsia</i>	+	+	+
24.	<i>Liza tade</i>	+	+	+
25.	<i>Stolephorus commersonii</i>	+	+	+
26.	<i>Stolephorus baganensis</i>	+	+	+
27.	<i>Stolephorus kammalensis</i>	+	+	+
28.	<i>Lutjanus johni</i>	+	+	+
29.	<i>Setipinna taty</i>	+	+	+
30.	<i>Lagocephalus lunaris</i>	-	+	+
31.	<i>Escualosa thoracata</i>	-	+	+
32.	<i>Epinephelus tauvina</i>	-	+	+
33.	<i>Epinephelus coioides</i>	-	+	+
34.	<i>Sphyraena sp.</i>	-	+	+
35.	<i>Pomadasys sp.</i>	-	+	+
36.	<i>Sardinella longiceps</i>	-	+	+
37.	<i>Periophthalmus sp.</i>	-	+	+
38.	<i>Macroglyphus sp.</i>	-	+	+
39.	<i>Tetraron cutcutia</i>	-	+	+
40.	<i>Bregmaceros meclllandii</i>	-	+	+
41.	<i>Ichthyocampus carce</i>	-	+	+
42.	<i>Stigmatogobius sp.</i>	-	+	+
43.	<i>Channa sp.</i>	-	+	-
44.	<i>Kurtus indicus</i>	-	+	-
45.	<i>Harpodon nehereus</i>	-	+	-
46.	<i>Moringua raitaborua</i>	-	+	-
47.	<i>Suggrundus rodricensis</i>	-	+	-

### C. Conservation Measures

The wasted juvenile species of osteichthyes have considerable total economic value and hence the destructive operation needs an immediate halt through few suggestive measures highlighted below:

- Development of alternative livelihood (like piggery, campbell culture, kroiler rearing, sewing, apiculture, fish feed manufacturing, handicrafts, spice grinding, skill development etc.) to defray the prawn seed collectors from the present destructive operation. Such livelihood schemes have already been implemented by several NGOs of Indian Sundarbans since 2005 and considerable success has been obtained and inclusion of WB. Govt. schemes like Tourism, Home stay, Education etc.

- Setting up of tiger prawn seed bank (nursery) in the high saline zone of Sundarbans and linking the same with adjacent maritime states like Orissa and Andhra Pradesh.
- Increase of awareness programme amongst the prawn seed collectors. Several Academic Institutes and reputed NGOs of the state have initiated this task through several projects financed by Ministry of Environment & Forests, Govt. of India, Dept. of Science & Technology, Govt. of India, Dept. of Environment, Govt. of West Bengal etc.
- Encouraging the people for canal excavation to harvest the rain water which can help in production of huge quantum of brackish water fishes.
- Manufacturing of bio-fertilizers from decayed plants in mangroves.
- Promotion and subsequent scaling up of non-conventional pilot projects like seaweed culture, oyster culture [23], ornamental fishery, bio-fertilizer preparation, herbal drinks from mangrove associate floral species, food products from seaweeds and mangrove associate etc. Such small scale projects have already been initiated by research scholars of Techno India University, West Bengal [24-27].

Again, a new concept of conservation is soon going to crystallize in the Indian Sundarbans regions in the form of 'Rural Technology Park'. In this unit, apart from awareness generation programme through adult education, some experimental ponds will be designed to monitor the survival rate of fish juveniles which are thrown away and wasted during wild tiger prawn seed collection. It is expected that such approach may reduce the mortality percentage of fish species in their juvenile stage [28].

#### Acknowledgement

The authors gratefully acknowledge the scientists of Zoological Survey of India for identifying the fish juveniles and Dr. Uttam Kumar Panja, Joint Director, Fisheries Department, Govt. of West Bengal for overall domain knowledge on the problem.

#### REFERENCES

- [1] Verlecar, X.N. and Parulekar, A.H. 2001. Primary Productivity. In: The Indian Ocean- A Perspective, Volume 2. (Ed. Rabin Sengupta & Ehrlich Desa). Oxford and IBH Publishers Co. Pvt. Ltd. Chapter 10, pp. 397-415.
- [2] Banerjee, K., Mitra, A., Bhattacharyya, D.P. and Choudhury, A. 2002. Role of nutrients on phytoplankton diversity in the northeast coast of the Bay of Bengal. In: Ecology and Ethology of Aquatic Biota. (Ed. Arvind Kumar ) Daya Publishing House, New Delhi, Chapter 6, 102-109.
- [3] Mitra, A. 2000. The northeast coast of the bay of Bengal and deltaic Sundarbans. In: Seas at the Millennium- An Environmental Evaluation, Chapter 62 (Ed. Charles Sheppard), University of Warwick, Country, U.K., Elsevier Science, 143-157.
- [4] Mitra, A. and Pal, S. 2002. In: The Oscillating Mangrove Ecosystem and the Indian Sundarbans (Eds. Col. Shakti Banerjee & Farida Tampal). Published by WWF-India, West Bengal State Office.
- [5] Mitra, A., Choudhury, A and Zamaddar, Y.A. 1992. Effects of heavy metals on benthic molluscan communities in Hooghly estuary. Proceedings of the Zoological Society, 45,481-496.
- [6] Trivedi, S., Mitra, A. Choudhury, A., Gupta, A., B. Singh and A. Choudhuri. 1994. A case study of the loss of Bio-diversity during Prawn Seed Collection from the Hooghly Estuary, India. Proceedings of the National Convention of Environment of India- Challenges for the 21<sup>st</sup> century; Institution of Public Health Engineers, India, pp. T-V/16-T-V/24.
- [7] Nageswara Rao, C.A. 2004. Faunal Diversity: Estuarine Ecosystem, EPTRI-ENVIS Newsletter, 10(3).
- [8] Mitra, A., Banerjee, K. and Bhattacharyya, D.P., 2004. In: The Other Face of Mangroves, 1<sup>st</sup> ed. Dept. of Environment, GoWB, India.
- [9] Shannon, C.E. and Weiner, V. 1949. In: The Mathematical Theory of Communication. University of Illionis Press, Urbana. 117pp.
- [10] Mitra, A., Banerjee, K and Gangopadhyay, A. 2004. In: Introduction to Marine Plankton. Daya Publishing House, New Delhi, India.
- [11] Banerjee, K., Das, S. and Mitra, A. 2005. Temporal loss of ichthyoplankton due to wild harvest of prawn seeds in Indian Sundarbans, Seshiyana (In press)
- [12] Banerjee, K., Bhattacharyya, D.P. and Mitra, A. 2005. Tiger prawn seed catch in coastal waters: Estimation of ichthyoplankton community damage, SIOS.(In press)
- [13] Mitra, A., Banerjee, K., Chakraborty, R, Banerjee, A., Mehta, N. and Berg, H. 2005. Study on the water quality of the shrimp culture ponds in Indian Sundarbans. Indian Science Cruiser.(In press)
- [14] Mitra, A. and Bhattacharyya, D.P. 2003. Environmental issues of shrimp farming in mangrove ecosystems, Journal of Indian Ocean Studies, 11 (1).
- [15] Bhattacharya, M., Banerjee, K., Roy, A., Mitra, A and Bhattacharyya, D.P. 1999. Preliminary study on the impact of oil and grease on finfish juveniles around Haldia Port-cum-industrial complex. Journal of the Inland Fisheries Society of India, 31(2), 36-38.

- [16] Bhattacharyya, D.P., Mahapatra, B., Banerjee, K. and Mitra, A. 2003. Application of Statistics in Marine Biology- a right choice is required, *Sea Explorers*, 6, 13-20.
- [17] Panja, U. 2004. Shrimp seed collection and Ecological crop loss in coastal West Bengal, *World Environment Day Special, WWF-India, West Bengal*.
- [18] Mitra, A., Niyogi, S. and Choudhury, A. 1998. Impact of wild harvest of prawn seeds on the community structure of juvenile fish in the coastal part of West Bengal. In: *Living Resources of India's Exclusive Economic Zone* (Editors: S.Z.Qasim and G.S. Roonwal), Omega Scientific Publisher, 86-96.
- [19] Niyogi, S, Mitra, A., Saha, S. and Choudhury, A. 1998. Inter-relationship between diversity of prawn juveniles and mangroves in the coastal zone of West Bengal. In: *Living Resources of India's Exclusive Economic Zone* (Editors: S.Z.Qasim and G.S. Roonwal), Omega Scientific Publisher, 97-102.
- [20] Mitra, A., Sasmal, S.K., Choudhury, A. and Bhattacharyya, D.P. 2000. Seasonal variations of ichthyoplankton in mangrove ecosystem of Indian Sundarbans, *Indian Journal of Environment and Ecoplanning*, 3(3), 535-538.
- [21] Whitfield, A.K.. 1999. Ichthyofaunal assemblages in estuaries: A South African case study, *Reviews in Fish Biology and Fisheries*, 9,151-186.
- [22] Mitra, A., Banerjee, K., Raha, A.K., Ghosh, S. and Bhattacharyya, D.P. 2005. Status of shark community in coastal West Bengal. *Indian Science Cruiser*, 19(2), 1-9.
- [23] Mitra, A. 2013. In: *Sensitivity of Mangrove ecosystem to changing Climate*. Springer. DOI: 10.1007/978-; 81-322-1509-7, 323.
- [24] Pramanick, P., Zaman, S. and Mitra, A. 2014. Processing of fruits with special reference to *S. apetala* fruit jelly preparation, *International Journal of Universal Pharmacy and Bio Sciences*, 3(5), 36-49.
- [25] Biswas, S., Pal, N., Biswas, P., Zaman, S. and Mitra, A. 2018. Nutritional status of food products developed from *Salicornia brachiata*, *International Journal of Pharmacy and Biological Sciences*, 8(3), 546-551.
- [26] Pal, N., Biswas, S., Biswas, P., Zaman, S. and Mitra, A. 2018. *Suaeda* sp. based food products: A unique avenue for alternative livelihood, *Journal of Emerging Technologies and Innovative Research*, 5(9), 774-778.
- [27] Biswas, S., Pal, N., Fazli, P., Zaman, S. and Mitra, A. 2019. Samosa and kachuri from mangrove associate species: An innovative utilization of coastal flora, *International Journal for Research in Applied Science & Engineering Technology*, 7(1), 73-79.
- [28] Mitra, A. and Banerjee, K. In: *Living Resources of the Seas: Focus Sundarbans*, Published by WWF-India. (Ed: Col S. R. Banerjee), Canning Field Office, 24 Parganas(S), W.B. 96pp.

