

EFFECT OF VAM AND PHOSPHORUS LEVELS ON YIELD, QUALITY, NUTRIENT UPTAKE AND POST HARVEST NUTRIENT STATUS OF SUNFLOWER

*Kalaiyarasan.C¹., S.Jawahar¹, M.V.Sriramachandrasekharan², K.Suseendran¹, R.Ramesh¹, S.Ramesh¹ and R.Kanagarajan³

¹Department of Agronomy, ²Department of Soil Science and Agricultural Chemistry

³Department of Entomology, Faculty of Agriculture, Annamalai University, Annamalai Nagar – 608 002, Tamil Nadu, India.

Abstract

The field experiment was conducted at Experimental Farm, Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Tamil Nadu during July – October 2015 to study the effect of mycorrhizal inoculation (VAM) and different levels of phosphorus on yield, quality, nutrient uptake and post harvest nutrient status of hybrid sunflower cv. sunbred. The experiment was conducted by factorial randomized block design with two replications. The treatments consisted of 4 levels of P₂O₅ (0, 25, 50, 75 and 100 kg/ha) applied in the presence or absence of VAM inoculates. The results of experiment revealed that mycorrhizal inoculated plants produced maximum seed and stalk yield, oil content, oil yield, protein content, nutrient uptake than non mycorrhizal inoculation, nutrient availability were maximum in non mycorrhizal inoculation. Among the various phosphorus levels tried, P₂O₅ at 100 kg ha⁻¹ recorded maximum values for yield, quality, nutrient uptake while P₂O₅ at 25 kg ha⁻¹ registered maximum NPK availability over other levels. With regard to interaction effect, mycorrhizal inoculation with P₂O₅ @ 100 kg ha⁻¹ recorded maximum values for yield, quality and nutrient uptake of sunflower but it was on par with mycorrhizal inoculation with P₂O₅ @ 75 kg ha⁻¹. Non-mycorrhizal inoculation with P₂O₅ @ 25 kg ha⁻¹ recorded maximum values for post harvest nutrient status of Sunflower. Lowest values of yield, quality and nutrient uptake were recorded by non-mycorrhizal inoculation with P₂O₅ @ 25 kg ha⁻¹. Mycorrhizal inoculation with P₂O₅ @ 100 kg ha⁻¹ recorded minimum values for post harvest nutrient status of Sunflower.

Keywords : Sunflower, VAM, yield, quality, NPK uptake & availability.

Introduction

In India, oil seed crops constitute the second largest agricultural produce, next only to food grains and they contribute 5% to GNP. Sunflower oil has excellent nutritional properties, and has a relatively high concentration of linoleic acid (Seiler, 2007). It is also a wealthy source of vitamins A and D. Mycorrhiza has symbiotic association between the soil fungi and roots of higher plants (Smith *et al.*, 2010). These fungi enhance the plant growth through making availability of mineral nutrients such as P, Zn and Cu (Phiriet *et al.*, 2003). These glomeromycotan fungi bank on their plant host for carbon in return for which fungus improves nutrition especially phosphate nutrition (Smith and Read, 1997). Phosphorus is an essential plant macronutrient which is required to build important molecules such as nucleic acids and phospholipids, and plays vital role during energy transfer in processes like NADPH, ATP and regulation of enzymatic and metabolic reactions. P is an essential plant nutrient required for higher and sustained productivity of oil from sunflower. Its influence on seed yield, oil yield and oil quality has been well established (Bahl and Toor, 1999). Phosphorus moves by diffusion in soil and is taken up by plants through root interception. The true phosphate levels available to plant around the soil are used to be very low.

As phosphorus typically constitutes around 30-80% of total P in the soil as organic P but still the availability of 80-99% for uptake in plant is scarce because of different factors like adsorption, precipitation or conversion into organic forms. The conceivable role of AM fungi in terms of their ability in phosphate nutrition has been gaining much importance in recent years (Karandoshov, 2005). In general, external supply of nutrient in terms of costly fertilizers to crops by farmers as of low economy may lead to low yield and supply to common society. Hence, the present study was taken up to find out the effect of mycorrhizal inoculation and phosphorus on yield, quality, nutrient uptake and post harvest nutrient status of hybrid sunflower.

Materials and methods

Field experiment was conducted during July – October 2015 at the Experimental Farm, Department of Agronomy, Faculty of Agriculture, Annamalai University. The experimental soil was clay loam with pH 8.1, OC 5.0 g kg⁻¹, available N (235 kg ha⁻¹), P (22.1 kg ha⁻¹) and K (356 kg ha⁻¹). The experiment consisted of eight treatments and was laid out in factorial randomized block design with two replications. M₁-Non inoculated and M₂-inoculated *Glomus intraradices* were tried along with different phosphorus levels (P₁-25, P₂-50, P₃-75 and P₄-100 kg P₂O₅ ha⁻¹) through SSP. Recommended dose of 60:60 kg of NK ha⁻¹ was applied in the form of Urea, and MOP respectively. Half the dose of N and entire dose of K were applied basally. P was applied as per the treatments. The remaining quantity of N was applied at 30 DAS. The mycorrhizal inoculum was applied near

the root zone of sunflower. 2 gm of VAM was applied per plant by placement method. At harvest seed and stalk yield was recorded. Oil content was analysed by using commercial Nuclear Magnetic Resonance Spectrometer (NMRS) method and oil yield was calculated by multiplying seed yield with oil content. N content of seed was analysed and multiplied with 6.25 to get crude protein. Seed was analysed for N, P and K content and corresponding nutrient uptake was computed. Post harvest soil was analyzed for available N, P and K.

Results and discussion

Yield

Mycorrhizal inoculated plants showed significant influence on yield (Table 1). Mycorrhizal inoculation recorded the maximum seed yield (1845 kg ha⁻¹) and stalk yield (4069 kg ha⁻¹) than non-mycorrhizal inoculation. The seed yield increase in this treatment was 28 % than non mycorrhizal inoculation. The yield increase might be due to increased growth, which resulted in increased rate of photosynthesis and stomatal conductance which can be due to more absorption of nutrients and ultimately resulted in increased seed and stalk yield. These findings are conformed to the report of Tahereh Vaseghmanesh *et al.* (2013) and Naser Hajiketabiet *et al.* (2014).

Phosphorus levels significantly influenced the yield (Table 1). Among the different levels, P₂O₅ at 100 kg ha⁻¹ produced maximum seed yield (2048 kg ha⁻¹) and stalk yield (4379 kg ha⁻¹). Phosphorus improves plants photosynthesis, resulting in higher assimilates production and consequently higher grain yield. The lowest value for yield was recorded in the treatment P₂O₅ at 25 kg ha⁻¹. This result is in conformity with the findings of Babaeiet *et al.* (2012).

The interaction effect between the mycorrhizal inoculation and phosphorus was not significant (Table 2). The treatment combination of mycorrhizal inoculation along with P₂O₅ at 100 kg ha⁻¹ recorded higher values for yield but it was on par with mycorrhizal inoculation along with P₂O₅ at 75 kg ha⁻¹. This might be due to availability of phosphorus and other nutrients at both vegetative and productive stages. Similar findings were earlier reported by Vaseghmanesh *et al.* (2013). The lowest values for seed and stalk yield were recorded under the treatment combination of non mycorrhizal inoculation with 25 kg P₂O₅ ha⁻¹. This might be due to the absence of mycorrhiza resulted in reduced growth and yield attributing characters and seed and stalk yield. Similar findings were earlier reported by Ultra Jret *et al.* (2007) and Mostafa Heidari and Vahid Karami (2014).

Quality characters

Mycorrhizal inoculated plants significantly influenced the quality characters viz., oil content, oil yield and crude protein content (Table 1). Mycorrhizal inoculation recorded the highest oil content (42.54 %), oil yield (790 Kg ha⁻¹) and crude protein content (17.64%) than non-mycorrhizal inoculation. Increased oil content and oil yield in this might be due to increased P availability in the plants because of the P absorption and availability. Similar findings were earlier reported by Martin *et al.* (2012) and Abdallah *et al.* (2013). The reason for increase in crude protein content with the inoculation of mycorrhiza might be ascribed to the fact that the protein content had a higher degree of positive relationship with the mycorrhizal plants which have assimilated greater amounts of P and translocated to grains which assisted in enrichment of protein. Similar finding was earlier reported by Karami (2011). Minimum oil content, oil yield and protein content was noticed in the treatment M₁ (non-mycorrhizal inoculation). This might be due to lesser availability and uptake of nutrients for oil synthesis in the sunflower seeds. Similar findings were earlier reported by Mostafa Heidari and Vahid Karami (2014).

Among the different levels of phosphorus, application of P₂O₅ at 100 kg ha⁻¹ significantly recorded highest oil content (43.49 %), oil yield (891 Kg ha⁻¹) and crude protein content (18.30 %). Phosphorus is known to play an important role in carbohydrate metabolism and helps in conversion of carbohydrate into oil. The lowest values for quality characters were recorded in the treatment P₂O₅ at 25 kg ha⁻¹. Similar finding was earlier reported by Hemalatha *et al.* (2013) and Rajendran and Veeraputhiran (2001).

The interaction effect between the mycorrhizal inoculation and phosphorus was not significant. The treatment combination of mycorrhizal inoculation along with P₂O₅ at 100 kg ha⁻¹ recorded maximum values for quality characters but it was on par with mycorrhizal inoculation along with P₂O₅ at 75 kg ha⁻¹. This might be due to mycorrhizal inoculation and phosphorus application and its effect by enhancing photosynthesis activity which, in turn, resulted in seed formation, an increase in oil content, oil yield of hybrid sunflower. Similar result was earlier reported by Babaeiet *et al.* (2012). The lowest values for quality characters were recorded in the treatment combination of non mycorrhizal inoculation with 25 kg P₂O₅ ha⁻¹ which could be due to inadequate availability of nutrients.

Nutrient uptake

Mycorrhizal inoculated plants significantly influenced the nutrient uptake of N, P and K (Table 1). Mycorrhizal inoculation recorded the highest N uptake (79.5 kg ha⁻¹), P uptake (22.0 kg ha⁻¹) and K uptake (70.5 kg ha⁻¹), than non-mycorrhizal inoculation. The association of AMF with sunflower plants exerted a stimulating effect on P and K uptake by the plants (Apolino Jose Nogueira da Silva *et al.*, 2015).

Phosphorus levels significantly influenced the uptake of N, P and K (Table 1). Among the different levels, P₂O₅ at 100 kg ha⁻¹ recorded maximum N uptake (81.4 kg ha⁻¹), P uptake (23.52 kg ha⁻¹) and K uptake (72.5 kg ha⁻¹). Increased uptake of nutrients at higher doses of phosphorus might have resulted in initial build up of plants due to vigorous growth and high photosynthetic rate which led to better uptake throughout the crop growth period. The lowest nutrient uptake of N, P and K was recorded in the treatment P₂O₅ at 25 kg ha⁻¹. This result is in conformity with the findings of Rajendran and Veeraputhiran (2001) and Madhavi *et al.* (2015).

The interaction effect between the mycorrhizal inoculation and phosphorus was not significant (Table 2). The treatment combination of mycorrhizal inoculation along with P₂O₅ at 100 kg ha⁻¹ recorded higher uptake of N, P and K but it was on par with mycorrhizal inoculation along with P₂O₅ at 75 kg ha⁻¹. Similar finding was earlier reported by Kumaresan *et al.* (2003). The lowest nutrient uptake of N, P and K were recorded under the treatment combination of non mycorrhizal inoculation with 25 kg P₂O₅ ha⁻¹. Similar finding was earlier reported by Kumaresan *et al.* (2003).

Post harvest soil available nutrients

Non-mycorrhizal inoculation recorded the highest available N(214.5 kg ha⁻¹), P (18.18kg ha⁻¹) and K (321.8kg ha⁻¹). This might be due to poor uptake of N, P and K under this treatment. The least soil available N, P and K status was observed under mycorrhizal inoculation.

Among the phosphorus levels, 25 kg P₂O₅ ha⁻¹ recorded the highest post harvest N (219.6kg ha⁻¹), P (18.67kg ha⁻¹) and K (326.9kg ha⁻¹). The lowest values for N, P and K availability were recorded under P₂O₅ at 100 kg ha⁻¹.

The interaction effect between mycorrhizal inoculation and phosphorus was significant in both the crops (Table 2). The maximum amount of N, P and K was available in non-mycorrhizal inoculation with P₂O₅ @ 25 kg ha⁻¹. The least value was recorded under mycorrhizal inoculation with 100 kg P₂O₅ ha⁻¹ due to maximum uptake of N, P and K Similar findings were earlier reported by Choudhary *et al.* (2011).



Table 1. Effect of mycorrhizal inoculation (VAM) and phosphorus levels on yield, quality characters, nutrient uptake and post harvest nutrient status of sunflower

Treatments	Yield		Quality characters			Nutrient uptake (Kg ha ⁻¹)			Post harvest nutrient status (Kg ha ⁻¹)		
	Seed yield (kg ha ⁻¹)	Stalk yield (kg ha ⁻¹)	Oil content (%)	Oil yield (Kg ha ⁻¹)	Crude protein content (%)	N	P	K	N	P	K
VAM											
M1	1438	3570	40.29	586	15.43	75.5	17.9	66.5	214.5	18.18	321.8
M2	1845	4069	42.54	790	17.64	79.5	22.0	70.5	202.8	17.05	310.6
SEd	28.46	20.75	0.16	12.35	0.06	0.37	0.15	0.27	0.57	0.06	0.61
CD(P=0.05)	61.05	44.51	0.34	26.49	0.13	0.78	0.33	0.57	1.23	0.13	1.31
Phosphorus levels (kg ha⁻¹)											
P1	1197	3179	39.12	470	14.27	73.4	15.95	64.2	219.6	18.67	326.9
P2	1477	3589	40.58	603	15.96	76.1	18.44	67.1	213.6	18.10	320.7
P3	1843	4129	42.47	786	17.61	79.4	21.93	70.4	203.5	17.04	310.4
P4	2048	4379	43.49	891	18.30	81.4	23.52	72.5	197.9	16.65	307.1
SEd	40.25	29.35	0.22	17.46	0.09	0.52	0.22	0.38	0.81	0.08	0.86
CD(P=0.05)	86.34	62.95	0.48	37.46	0.19	1.11	0.47	0.81	1.74	0.18	1.85

Table 2. Interaction effect between VAM and phosphorus levels on yield, quality nutrient uptake and post harvest nutrient status of sunflower

Treatments	Seed yield (Kg ha ⁻¹)	Stalk yield (Kg ha ⁻¹)	Oil content (%)	Oil yield (kg ha ⁻¹)	Crude Protein (%)	Nutrient Uptake (kg ha ⁻¹)			Nutrient availability (kg ha ⁻¹)		
						N	P	K	N	P	K
M₁P₁	1019	2946	38.15	388.6	13.15	71.7	13.9	62.2	223.9	18.9	331.5
M₁P₂	1219	3271	39.18	477.7	14.28	73.4	15.7	64.3	221.10	18.7	327.4
M₁P₃	1571	3795	41.05	645.0	16.52	76.8	19.5	68.07	211.6	17.8	316.9
M₁P₄	1942	4268	42.78	831.0	17.78	80.3	22.5	71.5	201.5	17.2	311.3
M₂P₁	1376	3412	40.09	552.0	15.40	75.1	18.0	66.2	215.3	18.3	322.3
M₂P₂	1735	3908	41.99	729.0	17.65	78.5	21.2	69.9	206.1	17.6	314.0
M₂P₃	2114	4464	43.89	927.8	18.70	81.9	24.4	72.7	195.5	16.2	303.8
M₂P₄	2153	4491	44.19	952.0	18.82	82.5	24.6	73.4	194.4	16.09	302.8
SEd	56.9	41.5	0.31	24.70	0.13	0.73	0.31	0.53	1.15	0.12	1.22
CD (P = 0.05)	122.1	89.03	0.68	52.97	0.27	1.57	0.66	1.15	2.46	0.26	2.61

References

- Abdallah, M.M., A.A. Abd El-Monem, R.A. Hassanein and H.M.S. El-Bassiouny. 2013. Response of sunflower plant to the application of certain vitamins and arbuscularmycorrhiza under different water regimes. **Aust. J. Basic & Appl. Sci.**, **7**(2): 915-932.
- Antunes, P. M., K. Schneider, D. Hillis and J.N. Klironomos. 2007. Can the arbuscularmycorrhizal fungus *Glomus intraradices* actively mobilize P from rock phosphates?. **Pedobiologia**, **51**: 281-286.
- Apolino Jose Nogueira da Silva, Ricardo Alencar da Silva, Juliana da Silva Santos, Jordan Carlos Silva de Medeiros, Fabiola Gomes de Carvalho, Valeria Nogueira da Silva, Cosme Jales de Oliveira, Afranio César de Araujo, Luiz Eduardo Souza Fernandes da Silva and Julio Gomes Junior. 2015. Soil chemical properties and growth of sunflower (*Helianthus annuus*L.) as affected by the application of organic fertilizers and inoculation with arbuscularmycorrhizal fungi. **R. Bras. Ci. Solo.**, **39**: 151-161.
- Babaei, M., M.R. Ardakani, F. Rejali, A.H.S. Rad, F. Golzardi and S. Mafakheri. 2012. Response of agronomical traits of sunflower (*Helianthus annuus*L.) to co-inoculation with *Glomus intraradices* and *Pseudomonas fluorescens* under different phosphorus levels. **Ann. Biol. Resh.**, **3**(8): 4195-4199.
- Bahl, G.S., G.S. Toor. 1999. Efficiency of P utilization by sunflower grown on residual P fertility. **Bioresour. Technol.**, **67**: 97-100.
- Choudhary, S.K., M.K. Jat, S.R. Sharma and P. Singh. 2011. Effect of INM on soil nutrient and yield in groundnut field of semi-arid area of Rajasthan. **Legume Research – An International Journal**, **34**(4): 283-287.
- Gandhi, A.P., K. Jehan and V. Gupta. 2008. Studies on production of defatted sunflower meal with low polyphenol and phytate contents and its nutritional profile. **Asian Food J.**, **15**: 97-100.
- Hemalatha, S., V. Praveen Rao, J. Padmaja and K. Suresh. 2013. An overview on role of phosphorus and water deficits on growth, yield and quality of groundnut (*Arachis hypogaea*L.). **International Journal of Applied Biology and Pharmaceutical Technology**, **4**(3): 188-201.
- Karami, V. 2011. Effect of different strains of mycorrhizae on quantitative and qualitative characteristics of sunflower in drought stress condition. **MSc. Thesis** of agronomy, University Of Zabol, Iran (In Persian).
- Kumaresan, M., V.S. Shanmugasundharam, M. Govindasamy and T.N. Balasubramaniam. 2003. Influence of phosphorus management on soil available phosphorus and uptake of NPK in a maize-sunflower-cowpea cropping system. **Indian Journal of Agricultural Research**, **37**(1): 34-38.
- Madhavi, A., P. Surendra Babu, and P. Venkata Reddy. 2015. Requirement of phosphorus and its use efficiency by sunflower in high phosphorus soils. **Journal of Progressive Agriculture**, **6**(1): 27-30.
- Martin, S.L., S.J. Mooney, A.J. Dickinson, H.M. West. 2012. The effects of simultaneous root colonisation by three *Glomus* species on soil pore characteristics. **Soil Biology and Biochemistry**, **49**: 167-173.
- Mostafa Heidari and Vahid Karami. 2014. Effects of different mycorrhiza species on grain yield, nutrient uptake and oil content of sunflower under water stress. **Journal of the Saudi Society of Agricultural Sciences**, **13**: 9-13.
- Naser Hajiketabi, Reza Zarghami, Amir Hasan Omid, Mohsen Tarigh Al- Eslami. 2014. Effect of Drought stress and use of Vesicular Arbuscular Mycorrhizal (VAM) fungi (*Glomus mosseae* and *Glomus intraradices*) on Grain yield, Biological yield and Harvest Index on safflower (*Carthamus tinctorius* L.). **The 1st international conference on new ideas in agriculture Islamic Azad Univ., Isfahan, Iran.**
- Phiri, S., I.M. Rao, E. Barrios, B.R. Singh. 2003. Plant growth, Mycorrhizal association, nutrient uptake and Phosphorus dynamics in a volcanic ash soil in Colombia as affected by the establishment of *Tithonia diversifolia*. **Journal of Sustainable Agriculture**, **21**: 41-49.
- Rajendran, K. and R. Veeraputhiran. 2001. Phosphorus nutrients in sunflower - a review. **Agricultural Reviews**, **22**(1): 68-70.
- Seiler, G.J. 2007. Wild annual *Helianthus anomalus* and *H. deserticola* for improving oil content and quality in sunflower. **Indian Crop Prod.**, **25**: 95-100.
- Smith, E.S., E. Facelli, S. Pope, F.A. Smith. 2010. Plant performance in stressful environments: Interpreting new and established knowledge of the roles of arbuscular mycorrhizas. **Plant Soil**, **326**: 3-20.
- Tahereh Vaseghmanesh, Khodabakhsh Panahi Kordlaghari, Ghazanfar Mohamadi Neia and Abdolsamad Kelidari. 2013. The response of yield components of sunflower to mycorrhiza inoculation and phosphorus fertilizer. **Annals of Biological Research**, **4**(3): 101-104.
- Vaseghmanesh, T., K.P. Kordlaghari, G.M. Neia, A. Kelidari. 2013. The response of yield components of sunflower to mycorrhizal inoculation and phosphorus fertilizer. **Ann. Biol. Res.**, **4**: 101-104.
- Vijay Kumar. 2005. Influence of phosphorus solubilising biofertilizers on seed yield and quality in seed parent of sunflower hybrid KBSH-44. **M. Sc. (Agri.) Thesis**, Univ. Agric. Sci., Bangalore.
- Zubillaga, M.M., J.P. Aristi and R.S. Lavado. 2002. Effect of phosphorus and nitrogen fertilization on sunflower (*Helianthus annuus*L.) nitrogen uptake and yield. **J. Argon. Crop Sci.**, **188**: 267-274.