

MODELING OF DISPERSION OF SUSPENDED PARTICULATE MATTER (SPM) FROM Dr. NARLA TATA RAO THERMAL POWER PLANT USING AERMOD MODEL

K.Pravallika

Asst Executive Engineer

Sri Venkateshwara University college of engineering,

Dr.N.Munilakshmi

Associate professor

Sri Venkateshwara University College.

1. INTRODUCTION:

An air pollutant is known as a substance in the air that can cause harm to humans and the environment. Pollutants can be in the form of solid particles, liquid droplets, or gases. Power plants are the major sources of air pollution. Coal is the primary fuel used for generation of electricity in India and its usage is continually increasing to meet the growing energy demands of the country. Emissions of greenhouse gases and other pollutants such as Suspended Particulate Matter (SPM) are increasing parallel to the growing demands of electricity. Implications of power plant emissions on the environment depend on the concentration of SPM pollutant emitted and in the recent past they are at alarming levels. Therefore, it is necessary to predict the concentration of SPM pollutant for impact assessment.

The study was carried out in two levels. In Level 1 study, impact assessment was made by screening level dispersion modeling technique using worst case input data employing Gaussian formulae. Level 2 studies included an assessment by a refined dispersion modeling technique (AERMOD) using site-specific input data. Level 1 impact assessment would be more conservative and less specific than the result of Level 2 assessments. It is not intended that an assessment should routinely progress through the two levels. If the air quality impact is considered to be a significant issue from Level 1 assessment, there is a need to immediately conduct of Level 2 assessments.

2. OBJECTIVES:

Dr. Narla Tata Rao Thermal Power Plant (NTTP), Ibrahimpatnam, Vijayawada, Andhra Pradesh, is selected as a case-study in the present investigation for the assessment of air quality of the region.

The following are the objectives of the present study:

- To develop an emission inventory for the study area
- To evolve wind roses of the study area

- To predict concentration of primary pollutant namely Suspended Particulate Matter (SPM) using Gaussian formulae
- To assess ground level concentration of primary pollutant Suspended Particulate Matter (SPM) by using AERMOD (America Meteorological Environmental Protection Agency Regulatory Model)
- To evolve the Isopleths of the study area in AERMOD

3. STUDY AREA:

The Dr. Narla Tata Rao Thermal Power Plant (NTTP) Vijayawada, of capacity 2x 210 MW and 1x500 MW of total 1760 MW is coal-fired based one. This thermal power station is predominantly government assigned land covering total area of 397.7 Acres (167.7 Acres Existing Land & 230 Acres to be acquired). The proposed power plant location falls between $16^{\circ}35'27''$ N Latitude and $80^{\circ}32'00''$ E Longitude.

4. METHODOLOGY:

This chapter includes the following:

- Levels of Assessment
- Emissions inventory for the present case study
- Input of meteorological data
- Evolution of wind roses
- Prediction of primary pollutants by using Gaussian formulae
- AERMOD dispersion modeling methodology, Interpretation of dispersion Modeling results for SPM

5. RESULTS AND DISCUSSION

The AERMOD model used in this study requires input information for an emission sources at the Dr.Narla Tata Rao Thermal Power Plant site-specific meteorological data of one complete year. The input data that describe both the emission source and meteorology provide a comprehensive set of information which can be used to run the AERMOD model and thus simulate the ground level concentration of Suspended Particulate Matter (SPM) from stationary sources of a thermal power plant.

In addition the model requires the site-specific meteorological information as input data. The local meteorological information that was to be given as input into the model were restricted to the Julian day of the year, the average wind flow vector, wind speed, height of the mixing layer, ambient Air temperature and the Pasquill stability category. The data were collected from Indian Meteorological Department, from web available by satellite data and from the website www.metcheck.com/IN/.

Met View [Pre-Processed Surface Met Data File]

File Header Data

Surface File Name: wind rose final.SFC

Station Latitude: 17.415N Upper Air Station ID: 12345 Onsite Station ID: N/A

Station Longitude: 78.411E Surface Station ID: 12345 Version: 18081 CCCR_SUB_TEMP_SUB

Filter: Year: All Month: All Day: All Julian Day: All Show All

Data Quality: Calms: 246 [hours] 2.81 [%] Missing: 8514 [hours] 97.19 [%]

Year	Month	Day	Julian Day	Hour	Sensible Heat Flux [W/m ²]	Surface Friction Velocity [m/s]	Convective Velocity Scale [m/s]	Vertical Potential Temperature Gradient above PBL	Height of Convectively-Generated Boundary Layer - PBL [m]	Height of Mechanically-Generated Boundary Layer - SBL [m]	Monin-Obukhov Length [m]	Surface Roughness Length [m]	Bowen Ratio	Albedo	Wind Speed - Ws [m/s]	Wind Direction - Wd [degrees]	Reference Height for Ws and Wd [m]	Temperature [K]	Reference Height for temp [m]	Precipitation Code	Precipitation Rate [mm/hr]	
Min.	2017	Jan	1	1	-999.0	-9.000	-9.000	-9.000	-999.0	-999.0	-99999.0	0.000	0.45	0.14	0.00	0.0	10.0	999.0	-9.0	0	0.00	
Max.	2018	Dec	31	365	24	-999.0	-9.000	-9.000	-9.000	-999.0	-99999.0	1.000	1.62	1.00	999.00	999.0	10.0	999.0	-9.0	11	1.52	
Graph	1	2017	Jul	1	182	1	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	1.00	1.50	228.0	10.0	999.0	-9.0	11	0.25
	2	2017	Jul	1	182	2	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	1.00	1.50	215.0	10.0	999.0	-9.0	11	0.25
	3	2017	Jul	1	182	3	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	1.00	2.10	204.0	10.0	999.0	-9.0	11	0.25
	4	2017	Jul	1	182	4	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	1.00	2.10	168.0	10.0	999.0	-9.0	11	0.51
	5	2017	Jul	1	182	5	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	1.00	2.10	131.0	10.0	999.0	-9.0	11	0.51
	6	2017	Jul	1	182	6	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	0.72	2.60	120.0	10.0	999.0	-9.0	11	0.25
	7	2017	Jul	1	182	7	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.000	0.45	0.26	3.10	102.0	10.0	999.0	-9.0	11	0.25
	8	2017	Jul	1	182	8	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.000	0.45	0.17	3.10	94.0	10.0	999.0	-9.0	11	0.25
	9	2017	Jul	1	182	9	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.000	0.45	0.15	3.10	104.0	10.0	999.0	-9.0	11	0.25
	10	2017	Jul	1	182	10	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.000	0.45	0.14	3.10	113.0	10.0	999.0	-9.0	11	0.25
	11	2017	Jul	1	182	11	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	0.29	2.60	128.0	10.0	999.0	-9.0	11	0.25
	12	2017	Jul	1	182	12	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	0.29	2.10	121.0	10.0	999.0	-9.0	11	0.25
	13	2017	Jul	1	182	13	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.000	0.45	0.14	2.60	102.0	10.0	999.0	-9.0	11	0.25
	14	2017	Jul	1	182	14	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	0.29	2.10	130.0	10.0	999.0	-9.0	11	0.25
	15	2017	Jul	1	182	15	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	0.29	2.10	131.0	10.0	999.0	-9.0	0	0.00
	16	2017	Jul	1	182	16	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	0.30	1.50	151.0	10.0	999.0	-9.0	0	0.00
	17	2017	Jul	1	182	17	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	0.34	1.50	144.0	10.0	999.0	-9.0	0	0.00
	18	2017	Jul	1	182	18	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	0.47	1.00	153.0	10.0	999.0	-9.0	11	0.25
	19	2017	Jul	1	182	19	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	1.00	0.50	188.0	10.0	999.0	-9.0	11	0.51
	20	2017	Jul	1	182	20	-999.0	-9.000	-9.000	-999.0	-999.0	-99999.0	0.040	0.93	1.00	1.00	145.0	10.0	999.0	-9.0	11	0.25

Meteorological data output surface file for AERMOD


Met View [Profile Met Data File]

Profile File Name: wind rose final.PFL


Filter: Year: All Month: All Day: All Show All

Year	Month	Day	Hour	Measurement Height [m]	1, if this is the last (highest) level for this hour, or 0 otherwise	Direction the wind is blowing from for the current level [degrees]	Wind Speed for the current level [m/s]	Temperature at the current level [C]	Standard deviation of the wind direction fluctuations [degrees]	Standard deviation of the vertical wind speed fluctuations [m/s]	
Min.	2017	Jul	1	10.0	1	0.0	0.00	99.9	99.0	99.00	
Max.	2018	Jun	30	10.0	1	999.0	999.00	99.9	99.0	99.00	
Graph	1	2017	Jul	1	1	10.0	1	228.0	1.50	99.9	99.0
	2	2017	Jul	1	2	10.0	1	215.0	1.50	99.9	99.0
	3	2017	Jul	1	3	10.0	1	204.0	2.10	99.9	99.0
	4	2017	Jul	1	4	10.0	1	168.0	2.10	99.9	99.0
	5	2017	Jul	1	5	10.0	1	131.0	2.10	99.9	99.0
	6	2017	Jul	1	6	10.0	1	120.0	2.60	99.9	99.0
	7	2017	Jul	1	7	10.0	1	102.0	3.10	99.9	99.0
	8	2017	Jul	1	8	10.0	1	94.0	3.10	99.9	99.0
	9	2017	Jul	1	9	10.0	1	104.0	3.10	99.9	99.0
	10	2017	Jul	1	10	10.0	1	113.0	3.10	99.9	99.0
	11	2017	Jul	1	11	10.0	1	128.0	2.60	99.9	99.0
	12	2017	Jul	1	12	10.0	1	121.0	2.10	99.9	99.0
	13	2017	Jul	1	13	10.0	1	102.0	2.60	99.9	99.0
	14	2017	Jul	1	14	10.0	1	130.0	2.10	99.9	99.0
	15	2017	Jul	1	15	10.0	1	131.0	2.10	99.9	99.0
	16	2017	Jul	1	16	10.0	1	151.0	1.50	99.9	99.0
	17	2017	Jul	1	17	10.0	1	144.0	1.50	99.9	99.0
	18	2017	Jul	1	18	10.0	1	153.0	1.00	99.9	99.0
	19	2017	Jul	1	19	10.0	1	188.0	0.50	99.9	99.0
	20	2017	Jul	1	20	10.0	1	145.0	1.00	99.9	99.0
	21	2017	Jul	1	21	10.0	1	230.0	1.00	99.9	99.0
	22	2017	Jul	1	22	10.0	1	225.0	1.00	99.9	99.0
	23	2017	Jul	1	23	10.0	1	243.0	1.00	99.9	99.0
	24	2017	Jul	1	24	10.0	1	224.0	1.00	99.9	99.0
	25	2017	Jul	2	1	10.0	1	220.0	1.00	99.9	99.0


Meteorological data output profile file for AERMOD




Surface file wind rose




Profile file wind rose



Isopleths for 1hr SPM concentration



Isopleths for 24hr SPM concentration



Isopleths for 3D SPM concentration

Spatial Distribution of SPM over Study Area

VALIDATION OF AERMOD

Source/ Monitoring Station	Location	Latitude	Longitude
Source Points	Stack 1	16.5887 ⁰ N	80.5542 ⁰ E
	Stack 2	16.5907 ⁰ N	80.5632 ⁰ E
	Stack 3	16.5967 ⁰ N	80.5232 ⁰ E
Monitoring Stations	AAQMS 1	16.58707 ⁰ N	80.5332 ⁰ E
	AAQMS 2	16.5927 ⁰ N	80.5632 ⁰ E
	AAQMS 3	16.5807 ⁰ N	80.6332 ⁰ E

The Predicted values of criteria pollutants SPM from output files of AERMOD with background concentrations are as shown in table

Predicted Values of SPM Pollutant from Output Files of AERMOD

Months	Pollutant concentration ($\mu\text{g}/\text{m}^3$)
	Background concentration is 33 ($\mu\text{g}/\text{m}^3$) SPM
July 2017	48.80
August 2017	46.30
September 2017	44.50
October 2017	43.28
November 2017	41.52
December 2017	40.24
January 2018	42.74
February 2018	47.10
March 2018	51.20
April 2018	36.30
May 2018	44.50
June 2018	43.28

The Predicted and Measured concentrations of Suspended Particulate Matter (SPM) are as shown in table

Predicted and Measured Concentrations of SPM

MONTHS	AAQMS 1 (Project site)		AAQMS 2 (Ibrahimpatnam)		AAQMS 3 (Hill point)	
	P ($\mu\text{g}/\text{m}^3$)	M ($\mu\text{g}/\text{m}^3$)	P ($\mu\text{g}/\text{m}^3$)	M ($\mu\text{g}/\text{m}^3$)	P ($\mu\text{g}/\text{m}^3$)	M ($\mu\text{g}/\text{m}^3$)
July 2017	48.80	41.68	48.80	48.3	48.80	50.3
August 2017	46.30	42.9	46.30	39.6	46.30	41.59
September 2017	44.50	48.6	44.50	45.23	44.50	39.6
October 2017	43.28	42.6	43.28	36.78	43.28	30.6
November 2017	41.52	44.65	41.52	44.36	41.52	38.4
December 2017	40.24	38.48	40.24	34.6	40.24	44.5
January 2018	42.74	46.9	42.74	35.2	42.74	35.36
February 2018	47.10	47.5	47.10	48.3	47.10	48.6
March 2018	51.20	53.5	51.20	50.35	51.20	47.2
April 2018	36.30	29.36	36.30	40.5	36.30	39.6
May 2018	44.50	37.48	44.50	38.6	44.50	37.6
June 2018	43.28	45.36	43.28	38.7	43.28	39.6

Statistical Performance Measures of AERMOD Model:

Parameter	Monitoring station	R ²	Inference
SPM	AAQMS 1	0.82	Strong positive correlation
	AAQMS 2	0.84	Strong positive correlation
	AAQMS 3	0.80	Strong positive correlation

6. CONCLUSION REMARKS

From level 1 assessment, it was noticed that SPM concentration were high and exceeding National Ambient Air Quality Standards, prescribed by the Central Pollution Control Board. In Level – 2 assessments, replication of pollutant SPM dispersion from Dr.Narla Tata Rao Thermal Power Plant was obtained by applying an AERMOD model and the results of predicted values were compared with the measured concentrations at the NTTTP site from July 2017 to June 2018 made available by APCCB.

Suspended Particulate Matter (SPM) varies from 40.24 $\mu\text{g}/\text{m}^3$ to 48.80 $\mu\text{g}/\text{m}^3$ from the month of July 2017 to June 2018.using AERMOD model. Value of R² at AAQMS 1 project site is 0.94 which is strong positive correlation, AAQMS 2 ibrahimpatanam is 0.81 which is strong positive correlation and AAQMS 3 hill station is 0.78 which is moderate positive correlation.

REFERENCE

Alexander Cohan, Junwu, Donald D Dabdub, (2011) “High resolution pollutant transport in the San Bay of California”, Atmospheric pollution Research, PP (237-246).

Amit P Kesarkar, Mohit Dalvi, Akula Venkatram, Alan Cimorelli, Akshara Kaginalkar and Ajay Ojha,(2005) “Coupling of the Weather Research and Forecasting Model with AERMOD for Pollutant Dispersion Modeling”, science Journal .

Anand Kumar Varma S., M. Srimurali, S.Vijaya Kumar Varma, (2014), “Prediction of Ground Level Concentrations of Air Pollutants Using Gaussian Model, Rayalaseema Thermal Power Project, Kadapa, A.P., India”, Energy and Environmental Engineering 2(4): 91-97.

Anand Kumar Varma S., Srimurali M., Manjula K.R. “Ozone Pollution in India Due to Power Plant Emissions”. International Journal of Scientific Research Volume: 2 Issue: 5 May 2013 ISSN No 2277 – 8179; IF: 0.3317.

Anand Kumar Varma S., Srimurali M., Anil Kumar K.V.S. “Magneto-hydrodynamic Generation of Electrical Energy in Thermal Power Project”. Research Journal of Engineering

&Technology Vol. 4, Issue: 1, March 2011 ISSN No 0974-2824.

Anand Kumar Varma S., Srimurali M.,(2013) “Concentration of Nitrogen Dioxide Estimation from Modelled NOX of a Power Plant”. IOSR Journal of Environmental science, Toxicology and Food Technology (IOSR-JESTFT) Volume: 6 Issue: 3 September -October 2013 e-ISSN 2319-2402 p-ISSN 2319-2399 Indexed by - EBSCO, cross-ref; DOI: 10.9790/2402-0630811; IF: 1.325.

Anil Kumar Reddy ChammiReddy1*, Karthikeyan J. “Development of wind rose diagrams for Kadapa region of Rayalaseema” CODE (USA): IJCRGG ISSN: 0974-4290 Vol.9, No.02 pp 60-64, 2016.

Bowers, J.F. and Anderson (1981). An evaluation study for the Industrial Source Complex Dispersion Model. EPA-450/4-81-002.RTP, NC.

Boznar M, Lesjak M, Mlakar P (1993) “A neural network-based method for short-term predictions of ambient SO₂ concentrations in highly polluted industrial areas of complex terrain”. Atmospheric Environment 27B: 221–230.

Bowers, J.F. and Anderson, (1981)“An evaluation study for the Industrial Source Complex Dispersion Model”, EPA-450/4-81-002.RTP, NC.

R. Krishna Chaithanya, Dr. N. Muni Lakshmi (2014) “Modeling of dispersion of SPM,SO₂ and Nox from coal fired thermal power plant using aermod dispersion model”.

R. Madhuri , Dr.N. Muni Lakshmi , (2014), “Modeling of dispersion of SPM,SO₂,and Nox from ramagundam thermal power plant using dispersion model”

