

SYNTHESIS OF SOME NEW 2-ISOTHIAZOLINE DERIVATIVES OF CHALCONES AND THEIR ANTIBACTERIAL ACTIVITY

Dr. S.M. Naik
I/C Principal

Naran lala College of Professional & Applied Sciences, Navsari.

ABSTRACT

The Isothiazoline derivatives can be synthesized from the chalcones by the action of hydroxyl amine hydrochloride, followed by the reaction with phosphorous penta sulphide. All these synthesized derivatives of Isothiazoline have been screened for antibacterial activity and characterized by spectral studies.

Keywords: Substituted aldehyde, Chalcones, 2-isothiazoline, 2-hydroxy-5-methyl-4,6-dichloroacetophenone Antibacterial activity, IR/NMR Spectroscopy.

INTRODUCTION

Chalcones have many pharmacological activity such as antimicrobial¹, antiviral², anti-inflammatory³, antifungal⁴, antispasmodics⁵ etc. Various chalcone derivatives can be synthesized by the condensation of aryl ketone with the various substituted aldehydes^{6,7}. Chalcone and its related heterocyclic derivatives such as isothiazoline, oxazine, pyrazoline, isoxazole, thiazine, pyrimidine, benzthiazepine, quinoxaline etc., shows antibacterial activity against various gram positive and gram negative bacteria⁸⁻¹². We report the reaction of 2-hydroxy-5-methyl-4,6-dichloroacetophenone with various substituted aromatic aldehydes to produced corresponding 2'-hydroxy-5'-methyl-4',6'-dichloro chalcones[1A-1J]., which on treatment with hydroxyl amine hydrochloride, gives 2-isoxazoline derivative[2A-2J] and finally reaction with phosphorous pentasulphide produced 2-isothiazoline derivatives[3A-3J]. The constitution of all compounds synthesized was characterized by elemental analysis, IR and H¹ NMR spectral study. The antibacterial activity of these synthesized compounds were also carried out.

MATERIAL AND METHODS

All melting points were taken in open capillary tubes and are incorrect. IR spectra in KBr were recorded on perkin-Elmer-377 spectrophotometer and H¹ NMR spectra were recorded on Varian NMR spectrophotometer. All compounds gave satisfactory elemental analysis.

General method for the synthesis of 2'-hydroxy-5'-methyl-4',6'-dichloro chalcones[1A-1J]

A mixture of 2-hydroxy-5-methyl-4,6-dichloroacetophenone (0.01 mole) and aryl aldehyde (0.01 mole) in ethanol (30 ml) was stirred and to it excess of 40% potassium hydroxide (25 ml) solution was added. The mixture was kept overnight at room temperature. The colour of the reaction mixture was change from yellow to orange. The content was then poured over crushed ice and acidified with hydrochloric acid (1:1). The solid separated was filtered, washed with distilled water, dried and crystallized by using ethanol, yield 60-70%.


General method for the synthesis of 3-(2'-hydroxy-5'-methyl-4',6'-dichlorophen-1'-yl)-5-(substituted phenyl)-2-isoxazoline[2A-2J]

A mixture of 2'-hydroxy-5'-methyl-4',6'-dichloro chalcone (0.01 mol) and hydroxylamine hydrochloride (0.02 mol) in ethanol (25 ml) was refluxed on water bath at 60-70°C for 4 hours. The reaction was cooled and acidified with glacial acetic acid. The solid separated was filtered and washed with water, dried and crystallized from ethanol (99%), yield 40-50%.

General method for the synthesis of 3-(2'-hydroxy-5'-methyl-4',6'-dichlorophen-1'-yl)-5-(substituted phenyl)-2-isothiazoline[3A-3J]

A mixture of 3-(2'-hydroxy-5'-methyl-4',6'-dichlorophen-1'-yl)-5-(substituted phenyl)-2-isoxazoline (0.01 mol) and phosphorous pentasulphide (0.01 mol), was taken in pyridine (20 ml). The reaction mixture was refluxed on water bath at 80-90°C for an hour. The reaction mixture was then cooled and diluted with water. The solid obtained was filtered, washed with water and crystallized from ethanol (99%), yield 40-60%.

REACTION SCHEME



Where R = 4-chlorophenyl, 4-hydroxyphenyl, Phenyl, 2,4-dichlorophenyl, 3-phenoxyphenyl, 2,6-dichlorophenyl, 3-nitrophenyl, 3,4,5-trimethoxyphenyl, 4-methoxyphenyl, 4-N,N-dimethylaminophenyl.

SCHEME
TABLE- 1

Characterization Table of 3-(2'-hydroxy-5'-methyl-4',6'-dichlorophen-1'-yl)-5-(substituted phenyl)-2-isothiazoline[3A-3J]

Compd. No.	R	Molecular formula	(M. wt.)	Yield (%)	M.P. °C.
3A	4-chlorophenyl	C ₁₆ H ₁₂ ONSCl ₃	371.79	39	146
3B	4-hydroxyphenyl	C ₁₆ H ₁₃ O ₂ NSCl ₂	353.35	40	122
3C	Phenyl	C ₁₆ H ₁₃ ONSCl ₂	337.35	42	133
3D	2,4-dichlorophenyl	C ₁₆ H ₁₁ ONSCl ₄	406.24	40	128
3E	3-phenoxyphenyl	C ₂₂ H ₁₇ O ₂ NSCl ₂	429.44	35	144
3F	2,6-dichlorophenyl	C ₁₆ H ₁₁ ONSCl ₄	406.24	54	154
3G	3-nitrophenyl	C ₁₆ H ₁₂ O ₃ N ₂ SCl ₂	382.35	33	134
3H	3,4,5-trimethoxyphenyl	C ₁₉ H ₁₉ O ₄ NSCl ₂	427.43	32	154
3I	4-methoxyphenyl	C ₁₇ H ₁₅ O ₂ NSCl ₂	367.37	40	179
3J	4-N,N-dimethylaminophenyl	C ₁₈ H ₁₈ ON ₂ SCl ₂	380.42	45	114

¹H NMR Spectroscopy

Nuclear magnetic resonance (NMR) spectroscopy is one of the latest physical methods which is used for the structure determination of organic compounds. PMR spectra of 2-isothiazoline derivatives were recorded on a varian spectrophotometer. Spectra were examined in CDCl₃ at room temperature using TMS as an internal standard.

TABLE-2

200 MHz PMR spectra of 3-(2'-hydroxy-5'-methyl-4',6'-dichlorophen-1'-yl)-5-(4''methoxyphenyl)-2-isothiazoline (Compound no. 3I)

Chemical shift	Relative Number of Protons	Assignment
2.30	3	-CH ₃
2.57	2	-CH ₂ of isothiazoline ring
3.82	3	-OCH ₃
4.32	1	-CH of isothiazoline ring
6.33	1	-OH
6.75-7.80	5	Ar-H

Infrared spectra

Infrared absorption was recorded using potassium bromide pellets method. The spectra were recorded using "Perkin-Elmer" spectrophotometer. The results are described in table no. 3.

TABLE-3

IR spectra of 3-(2'-hydroxy-5'-methyl-4',6'-dichlorophen-1'-yl)-5-(4''chlorophenyl)-2-isothiazoline (Compound no. 3A)

Position of absorption band (cm ⁻¹)	Intensity	Band and its mode of vibration	Functional group
780	S	C-Cl stretching	Chloro compound
845	S	N-S stretching	Isothiazoline ring
950	M	C-S stretching	Isothiazoline ring
1385	Sh	O-H bending	Ar-OH intramolecular
1470	S	C-H bending	-
1590	V	C=N stretching	Compound containing C=N group
2980	M	C-H stretching	-
3400	Sh	O-H stretching	Ar-OH group

S=strong, m=medium, b=broad, w=weak, sh=sharp, v=variable

Antibacterial activity

The synthesized compounds were screened for their antibacterial activity using *S.aureus*, *E. coli* by cup plate method using DMF as solvent. All the compounds shows mild activity against both bacteria in comparison with ampicillin and gentamycin. The results are describe in table no. 4.

Table-4

Compound No.	Zone of inhabitation in mm Antibacterial (24 hrs.)	
	S.aureus (+ve)	E.coli (-ve)
3A	9	11
3B	13	12
3C	10	8
3D	14	13
3E	N.A.	N.A.
3F	11	10
3G	14	12
3H	10	11
3I	12	17
3J	13	14
Standard Drugs:		
Ampicilin	18	-
Gentamycin	-	21

RESULTS AND DISCUSSION

2-isothiazoline derivatives shown medium activity against both bacteria i.e. *S.aureus* and *E.coli*. From the table no. 4, compound no. **3I** shown the maximum activity amongst all the compounds towards –ve bacteria i.e. *E.coli*. The compound no. **3D** and **3G** have shown good activity against *S.aureus* bacteria. The compound no. **3E** found inactive against both bacteria while rest of the compounds have shown the medium to poor activity against both bacteria. The activities of synthesized compounds are compared with standard antibiotics like Ampicillin and Gentamicin. It was observed that the synthesized compounds are less active than the standard one.

ACKNOWLEDGEMENT

The author is thankful to Department of Chemistry,

REFERENCES

1. Y. B. Vibhute; Indian J. Chem. Soc., **53**, 736 (1976); C.A. **86**, 106308 (1977).
2. K.G.Mallikarjun; E-Journal of Chemistry, **2(1)**, 58 -61 (2005).
3. Y. R. Prasad, A. Srinivasa Rao And R. Rambabu; Asian Journal of Chemistry, **21(2)**, 907-914 (2009).
4. C. K. Bradsher, F. C. Brown, and W. B. Blue; Am. Chem. Soc., **71**, 3570 (1949).
5. S. Oshiro, T. Nagura, Y. Sugihara, K. Okamoto, R.Ishida and K. Shintomi; Japanese Patent, **73** 19,569, 19,594 (1973); C.A. **78**, 147788_g, 147790_b, 147963_k (1973).
6. G. N. Vyas and N. M. Shah; Indian J. Chem. Soc., **28**, 75 (1951).
7. S. M. Naik and H. B. Naik, Orient. J. Chem., **14(1)** 167-168 (1998).
8. Chinna Rao Gantla, Y.Suresh, S.Harikrishana And S.P. Shrivastava; Oriental Journal of Chemistry, **25(1)**, 153-157 (2009).
9. V.M. Barot And S.D. Desai; Asian Journal Of Chemistry, **21(8)**, 6091-6094 (2009).
10. R. Kalirajan, S. U. Sivakumar, S. Jubie and B. Gowramma, B. Suresh, Int. J. Chem. Tech. Res., **1(1)**, 27-34 (2009).
11. K. G. Desai and K. R. Desai; Indian Journal of Chemistry, **46B**, 1179-1186 (2007).
12. M. Kaithwal, P. Garg and S. Ahmad, Orient. J. Chem., **25(4)**, 935-943 (2009).