

# Bi-domination in Graphs

V. Lavanya<sup>1</sup>Dr. D. S. T. Ramesh<sup>2</sup> and Dr. N. Meena<sup>3</sup>

<sup>1</sup>Research Scholar, Department of Mathematics, Nazareth Margoschis College at Pillaiyanmanai, Nazareth 628 617, Affiliated to Manonmaniam Sundaranar University, India.

<sup>2</sup>Associate Professor, Department of Mathematics, Nazareth Margoschis College at Pillaiyanmanai, Nazareth 628 617, Affiliated to Manonmaniam Sundaranar University, India.

<sup>3</sup> Associate Professor, Department of Mathematics, MDT Hindu College, Pettai, Tirunelveli  
627004 Affiliated to Manonmaniam Sundaranar University, India.

**Abstract:** Let  $G = (V, E)$  be a simple graph. A set  $S \subseteq V(G)$  is a bi-dominating set if  $S$  is a dominating set of  $G$  and every vertex in  $S$  dominates exactly two vertices in  $V-S$ . The bi-domination number  $\gamma_{bi}(G)$  of a graph  $G$  is the minimum cardinality of the minimal bi-dominating set. In this paper, bi-domination number for some standard graphs are determined. Bounds for bi-domination number are obtained.

**Key Words:** Domination number, Bi-Domination number and Upper bi-domination number.

**1.Introduction:** Let  $G(V,E)$  be a simple, connected graph where  $V$  is its vertex set and  $E$  is its edge set. The degree of any vertex  $v$  in  $G$  is the number of edges incident with  $v$  and is denoted by  $\deg v$ , the minimum degree of a graph is denoted by  $\delta(G)$  and the maximum degree of a graph  $G$  is denoted by  $\Delta(G)$ . A vertex of degree 0 is called an isolated vertex and a vertex of degree 1 is called a pendent vertex. A subdivision of an edge  $uv$  is obtained by replacing the edge  $uv$  with the edges  $uw$  and  $wv$  with a new vertex  $w$ . A subset  $S$  of  $V(G)$  is called a dominating set of  $G$  if every vertex in  $V - S$  is adjacent to at least one vertex in  $S$  [2-3]. The domination number  $\gamma(G)$  of  $G$  is the minimum cardinality of all dominating sets in  $G$ . In this paper, bi-domination number for some standard graphs are determined. Bounds for bi-domination number are obtained. For graph theoretic notations, Harary [1] is referred to.

**Definition 1.1:** Let  $G = (V, E)$  be a simple graph. Let  $|V(G)| = n$ . A vertex  $v$  is called a full degree vertex if  $\deg v = n - 1$ .

**Definition 1.2:** The graph  $B_{n,n}$ ,  $n \geq 2$  is a bistar obtained from two disjoint copies of  $K_{1,n}$  by joining the centre vertices by an edge. It has  $2n + 2$  vertices and  $2n + 1$  edges.

**Definition 1.3:** A spider is a tree on  $2n + 1$  vertices obtained by subdividing each edge of a star. One or more (but not all) of the edges from this subdivision are removed results a wounded spider.[5]

## 2. Main Results:

**Definition 2.1:** A set  $S \subseteq V(G)$  is a bi-dominating set if  $S$  is a dominating set of  $G$  and every vertex in  $S$  dominates exactly two vertices in  $V-S$ .

**Remark 2.2:** The bi-domination number  $\gamma_{bi}(G)$  of a graph  $G$  is the minimum cardinality of all minimal bi-dominating sets. The maximum cardinality of a bi-dominating set of  $G$  is called the upper bi-domination number of  $G$  and it is denoted by  $\Gamma_{bi}(G)$ .

**Example 2.3:** Consider the following graph  $G$  in figure 2.1




Figure 2.1

Let  $S_1 = \{v_1, v_3, v_6\}$ ,  $S_2 = \{v_1, v_3, v_7\}$ ,  $S_3 = \{v_2, v_4, v_5\}$ ,  $S_4 = \{v_2, v_4, v_6\}$  and  $S_5 = \{v_2, v_4, v_7\}$ , every vertex of the set  $S_i$ ,  $1 \leq i \leq 5$  dominate exactly two vertices of  $V - S_i$ . Hence  $S_i$ ,  $1 \leq i \leq 5$  are bi-dominating sets of  $G$ . Therefore  $\gamma_{bi}(G) \leq 3$ .  $\{v_2, v_5\}$  is the unique minimum dominating set,  $\gamma(G) = 2$ . It is not a bi-dominating set, since  $v_5$  dominates three vertices  $v_4, v_6$  and  $v_7$ . Therefore  $\gamma_{bi}(G) \geq 3$ . Hence  $\gamma_{bi}(G) = 3$ .

**Example 2.4:** Consider the following graph  $G$  in figure 2.2




Figure 2.2

Let  $S_1 = \{a, d\}$ ,  $S_2 = \{e, c, f\}$ ,  $S_3 = \{b, f, d\}$ . Every vertex of the set  $S_i$ ,  $1 \leq i \leq 3$  dominates exactly two vertices in  $V - S_i$ . Hence  $S_i$ ,  $1 \leq i \leq 3$  are bi-dominating sets of  $G$ . Hence  $\gamma_{bi}(G) \leq 2$ , Since there is no full degree vertex in  $G$ ,  $\gamma(G) \geq 2$ . Therefore  $\gamma_{bi}(G) \geq 2$ . Also it is verified that no set with four vertices is a bi-dominating set. Hence  $S_1$  is the minimum bi-dominating set and  $S_2$  and  $S_3$  are maximum bi-dominating sets. Hence  $\gamma_{bi}(G) = 2$  and

$$\Gamma_{bi}(G) = 3.$$

**Remark 2.5:** Any bi-dominating set does not contain pendent vertices.

**Example 2.6:** Graph  $G$  without bi-dominating set is given in the following figure 2.3




Figure 2.3

Suppose  $G$  has bi-dominating set  $S$ . Since there is no full degree vertex in  $G$ ,  $\gamma(G) \geq 2$ . Hence  $\gamma_{bi}(G) \geq 2$ . Since every vertex in  $S$  dominates exactly two vertices in  $V - S$ ,  $\gamma_{bi}(G) \leq 3$ . Since  $e$  does not belong to  $S$ ,  $d$  must belong to  $S$ . Let  $S_1 = \{a, d\}$ ,  $S_2 = \{c, d\}$  and  $S_3 = \{b, d\}$ . In  $S_1$ ,  $d$  dominates three vertices  $b, c$  and  $e$  of  $V - S_1$  of  $G$ . Hence  $S_1$  is not a bi-dominating set of  $G$ . In  $S_2$ ,  $c$  dominates only one vertex  $a$  of  $V - S_2$  of  $G$ . Hence  $S_2$  is not a bi-dominating set of  $G$ .

In  $S_3$ ,  $b$  dominates only one vertex  $a$  of  $V - S_3$  of  $G$ . Hence  $S_3$  is not a bi-dominating set of  $G$ . Therefore no set with two elements is a bi-dominating set. In the set  $S_4 = \{a, b, d\}$ , the vertices  $a$  and  $b$  do not dominate two elements of  $V - S_4$  of  $G$ . In the set  $S_5 = \{a, c, d\}$ , the vertices  $a$  and  $c$  do not dominate two elements of  $V - S_5$  of  $G$ . In the set  $S_6 = \{c, b, d\}$ , no element dominate two vertices of  $V - S_6$  of  $G$ . Therefore no sets with three elements is a bi-dominating set of  $G$ . Hence bi-dominating set does not exist.

**Observation 2.6:** For any connected graph  $G$  with  $p$  vertices,  $\gamma_{bi}(G) = 1$  if and only if  $G \cong P_3$  or  $K_3$ .

**Remark 2.7:**  $\gamma(G) \leq \gamma_{bi}(G)$ , since every bi-dominating set is a dominating set.

**Example 2.8:** Consider a graph  $G$  given in example 2.  $\gamma(G) = 2$  and  $\gamma_{bi}(G) = 3$ . Therefore  $\gamma(G) < \gamma_{bi}(G)$ .

**Example 2.9:** For the Graph  $G$  given in figure 2.4,  $\gamma_{bi}(G) = \gamma(G)$




Figure 2.4

There is no full degree vertex in  $G$ . Therefore  $\gamma(G) \geq 2$ . Let  $S_1 = \{v_1, v_4\}$ ,  $S_2 = \{v_2, v_5\}$  and  $S_3 = \{v_3, v_5\}$ , Every vertex of the set  $S_i$ ,  $1 \leq i \leq 3$  dominates exactly two vertices of  $V - S_i$ . Hence  $S_i$ ,  $1 \leq i \leq 3$  are bi-dominating sets of  $G$ . Therefore  $\gamma(G) \leq 2$  and  $\gamma_{bi}(G) \leq 2$ . Hence  $\gamma(G) = 2$ . Since  $G$  is not either  $P_3$  or  $K_3$ ,  $\gamma_{bi}(G) = 2$ . Hence  $\gamma_{bi}(G) = 2 = \gamma(G)$

**Theorem 2.10:** Let  $P$  be a path of length  $n$ , then  $\gamma_{bi}(P_n) = \lceil \frac{n}{3} \rceil$ ,  $n \geq 3$ ,  $n \neq 2, 4$ .

**Proof: case (i):** Let  $n = 3m$ ,  $m \geq 1$ .  $\{v_2, v_5, \dots, v_{3m-1}\}$  is the unique bi-dominating set of  $P_{3m}$ . Therefore  $\gamma_{bi}(P_n) = \frac{n}{3}$ .

**Case (ii):** Let  $n = 3m + 1$ ,  $m \geq 2$ .  $\{v_2, v_5, v_8, \dots, v_{3m-4}, v_{3m-2}, v_{3m}\}$  is a bi-dominating set of  $P_{3m+1}$ . Therefore  $\gamma_{bi}(P_n) = m + 1 = \lceil \frac{n}{3} \rceil$ .

**Case (iii):** Let  $n = 3m + 2$ ,  $m \geq 1$ .  $\{v_2, v_4, v_7, v_{10}, \dots, v_{3m+1}\}$  are some bi-dominating set of  $P_{3m+2}$ . Therefore  $\gamma_{bi}(P_n) = m + 1 = \lceil \frac{n}{3} \rceil$ .

**Case (iv):** when  $n = 2$ .  $\{v_1\}$  and  $\{v_2\}$  are dominating sets and it dominates exactly one vertex. Hence it is not a bi-dominating set.

**Case (v):**  $S_1 = \{v_1, v_4\}$ ,  $S_2 = \{v_1, v_3\}$ ,  $S_3 = \{v_2, v_3\}$  and  $S_4 = \{v_2, v_4\}$  are dominating sets of  $P_4$ . No vertices in  $S_1$  and  $S_3$  dominates exactly two vertices.  $V - S_1$  and  $V - S_3$  respectively  $v_1$  in  $S_2$  and  $v_4$  in  $S_4$  do not dominate exactly two vertices in  $V - S_2$  and  $V - S_4$ .

**Theorem 2.11:** Let  $C$  be a cycle of length  $n$ , then  $\gamma_{bi}(C_n) = \lceil \frac{n}{3} \rceil$ ,  $n \geq 3$ .

**Proof: Case (i):** Let  $n = 3m$ ,  $m \geq 1$ .  $\{v_2, v_5, v_8, \dots, v_{3m-1}\}$  is the bi-dominating set of  $C_{3m}$ . Therefore  $\gamma_{bi}(C_n) = \frac{n}{3}$ .

**Case (ii):** Let  $n = 3m + 1$ ,  $m \geq 2$ .  $\{v_2, v_5, v_8, \dots, v_{3m-4}, v_{3m-2}, v_{3m}\}$  is a bi-dominating set of  $C_{3m+1}$ . Therefore  $\gamma_{bi}(C_n) = m + 1 = \lceil \frac{n}{3} \rceil$ .

**Case (iii):** Let  $n = 3m + 2$ ,  $m \geq 1$ .  $\{v_2, v_4, v_7, v_{10}, \dots, v_{3m+1}\}$  are some bi-dominating set of  $C_{3m+2}$ . Therefore  $\gamma_{bi}(C_n) = m + 1 = \lceil \frac{n}{3} \rceil$ .

**Case (iv):** when  $n = 4$ . The antipodal vertices  $\{v_1, v_2\}$  or  $\{v_3, v_4\}$  is a bi-dominating set of  $C_4$ .

**Theorem 2.12:** Let  $G$  be a connected graph, If  $G = K_n$  then  $\gamma_{bi}(G) = n - 2$ .

**Proof:** Let  $G = K_n$  be a connected graph  $\{v_1, v_2, \dots, v_n\}$  be the vertex set of  $G$ . Since  $G$  is complete and degree of every vertex in  $G$  is  $n - 1$ , clearly by definition the vertices  $v_1, v_2, \dots, v_{n-2}$  dominate exactly two vertices. Hence  $\gamma_{bi}(G) = n - 2$ .

**Theorem 2.13:** Let  $G$  be a complete bi-partite graph  $K_{m,n}$  and  $m, n \geq 3$ . Then  $\gamma_{bi}(G = K_{m,n}) = m + n - 4$ .

**Proof:** Let  $U = \{u_1, u_2, \dots, u_m\}$  and  $W = \{w_1, w_2, \dots, w_n\}$  be a bi-partition of  $G$ . Every  $u_i \in U$  dominates all the vertices in  $W$ . As each vertex can dominate exactly two vertices except two  $w_i$ 's all other  $w_i$ 's are in  $S$ , the bi dominating set. Similarly all  $u_i$ 's except two are in  $S$ . Hence  $\gamma_{bi}(G) = m - 2 + n - 2 = m + n - 4$ .

**Theorem 2.14:** For a star  $G = K_{1,k}$ ,  $k \neq 2$  dominating set exist,  $\gamma(G) = 1$  but bi-dominating set does not exist. **Proof:** Let  $K_{1,k}$  be a star with  $k$  pendant vertices. Let  $v$  be a vertex with maximum degree and  $v_1, v_2, \dots, v_k$  be the pendant vertices of  $K_{1,k}$ . The vertex  $v$  can dominate  $v_1, v_2, \dots, v_k$  vertices and the vertices  $v_1, v_2, \dots, v_k$  dominate exactly one vertex  $v$ . Hence  $\{v\}$  is the minimum dominating set and  $\gamma(G) = 1$ . The vertex  $v$  dominates more than two vertices but not exactly two. Clearly  $\{v\}$  is a dominating set but not bi-dominating. Therefore bi-dominating set does not exist for  $K_{1,k}$ .

**Note 2.15:** If  $G = K_{1,2}$  then  $\gamma(G) = \gamma_{bi}(G) = 1$ .

**Proof:** Let  $G = K_{1,2}$ . Let  $v$  be a vertex with maximum degree 2 and  $v_1$  and  $v_2$  be the pendant vertices of  $K_{1,2}$ . Since degree of  $v$  is 2 and it dominates other vertices  $v_1$  and  $v_2$ . Hence  $\{v\}$  is the minimum dominating set and it dominates exactly two vertices and it also a bi-dominating set. Therefore  $\gamma(G) = \gamma_{bi}(G) = 1$ .

**Remark 2.16:** For any connected graph  $G$  with  $n$  vertices,  $1 \leq \gamma_{bi}(G) \leq n - 2$ .

**Theorem 2.18:** A bi-dominating set does not exist for a graph  $G$  in which a vertex adjacent with more than two pendent vertices.

**Proof:** Let  $u_1, u_2, \dots, u_m, m \geq 3$  be the pendent vertices adjacent with a vertex  $v$  in  $G$ . Suppose  $S$  is a bi-dominating set of  $G$ . Since pendent vertices do not belong to  $S$ , the vertex  $v$  must belong to  $S$  to dominate  $u_1, u_2, \dots, u_m, m \geq 3$ . Then  $v$  dominate more than two vertices of  $V - S$ , a contradiction. Hence bi-dominating set does not exist in  $G$ .

**Observation 2.19:** For a bistar,  $B_{m,n}, m$  and  $n \neq 2$  bi-dominating set does not exist.

**Theorem 2.20:** For  $B_{2,2}$  a bi-star,  $\gamma(G) = \gamma_{bi}(G) = 2$

**Proof:** Let  $u$  and  $v$  be the central vertices of  $B_{2,2}$ . Let  $u_1, u_2$  and  $v_1, v_2$  be the vertices adjacent with  $u$  and  $v$  respectively. Since  $\{u, v\}$  is the unique minimum dominating set of  $B_{m,n}$  and  $\gamma(G) = 2$ . Clearly  $u$  dominate exactly two vertices  $u_1$  and  $u_2$  and  $v$  dominate exactly two vertices  $v_1$  and  $v_2$ . Hence  $\{u, v\}$  is unique bi-dominating set and  $\gamma_{bi}(G) = 2$ . Therefore  $\gamma(G) = \gamma_{bi}(G) = 2$ .

**Theorem 2.21:** For a wounded spider with at least one leg, bi-dominating set does not exist.

**Proof:** Consider the following wounded spider  $K_{1,t}^*$ , where  $k$  edges are subdivided and  $k < t$ .




Figure 2.5

Let  $u$  be the central vertex. Take  $D = \{u_i\}, 1 \leq i \leq t$  is a unique minimum dominating set, since the vertices  $u_{k+1}, \dots, u_t$  has only one neighbourhood  $u$ . Hence the set  $D$  is dominating set but not a bi-dominating set. Therefore bi-dominating set does not exist for wounded spider.

**Remark 2.22:** Complement of  $S$  a bi-dominating set of a graph  $G$  need not be a bi-dominating set. Consider the following graph  $P_5$




Figure 2.6

Here  $\{v_2, v_4\}$  is the unique bi-dominating set of  $P_5$ .  $V - S$  is a dominating set but not a bi-dominating set of  $P_5$

**Theorem 2.23:** Let  $G$  be a connected graph with  $n$  edges. Then  $\gamma_{bi}(S(G)) = n$ .

**Proof:** Let  $G$  be a simple graph with  $n$  edges. Let  $H = S(G)$  and  $|E(H)| = 2n$ . Let the edges of  $G$  subdivided by the new vertices  $u_1, u_2, \dots, u_n$ .  $\{u_1, u_2, \dots, u_n\}$  is the unique bi-dominating set of  $H$ . Therefore  $\gamma_{bi}(H) = n$ .

**References:**

[1] Harary, F., Graph theory, Addison-Wesley Publishing Company Reading, Mass (1972).  
 [2] Teresa W. Haynes, Stephen T. Hedetniemi, Peter J. Slater, Fundamentals of Domination in graphs, Marcel Dekker, Inc (1998).  
 [3] Teresa W. Haynes, Routledge, Domination in graphs: Volume 2: Advanced Topics, (2017)  
 [4] John Adrian Bondy, Murty U.S.R., Graph Theory, Springer, 2008.  
 [5] Selvam Avadayappan<sup>1</sup>, M. Bhuvaneshwari<sup>2</sup>, R. Iswarya<sup>3</sup>, International Journal for Research in Applied Science and engineering Technology, Volume 4 Issue III, March 2016.