

Ranking Triangular Fuzzy Numbers Using Area of Rectangle At Different Level Of α -Cut for Fuzzy Transportation Problem

Ashok S. Mhaske

Department of Mathematics Dada Patil Mahavidhyalaya, Karjat-Ahmednagar, Maharashtra, India

Email: mhaske.math@gmail.com

Abstract:

The uncertainty plays a vital role in any branch of science and technology, engineering, medical and management. In proposed work new method ranking to order two fuzzy triangular numbers. This new ranking method is used to find the best approximate solution to the fuzzy transportation problem. The objective of the transportation problem is to control the amount to be transported from several origins to several destinations such that the total transportation cost is minimized. But in real-life situations, the available supply and forecast demand are often fuzzy because some information is incomplete or unavailable.

In this article, the crisp transportation problem is transformed into a fuzzy transportation problem by using triangular fuzzy numbers. To order any two triangular fuzzy numbers the new and simple method is invented which are based on the area of a rectangle at prominent places using α -cut method at $\alpha=0.5$. A computer program was written in Mat-lab which is given in this article to make calculation easier and simple. This article is organized as follows. In section first we introduced new ranking method to compare any two trapezoidal fuzzy numbers. In third section we extend this method to solve fuzzy transportation problem using numerical example, while paper is concluded in the last section.

Keywords: Ranking, Rectangle, Triangular, Fuzzy Number, Fuzzy Transportation

Introduction

In Artificial Intelligence Fuzzy Mathematics and fuzzy logic is used to process Natural language and widely used in decision making. Theoretically, fuzzy numbers cannot be compared but merely partially ordered. But, when fuzzy numbers are used in real-world applications, e.g., when we want to determine optimal solution of objective function or want to make the decision among alternatives, then order of fuzzy numbers becomes important. However, there are many methods of fuzzy numbers say qualitative, quantitative and based on α -cuts to the ordering relation between any two fuzzy numbers. However in fuzzy set theory there is no universally acceptable method to compare any two fuzzy numbers. The ranking of fuzzy number using α -weighted valuations was started by *Marcin Detyniecki and Ronald R Yager* [9] in year 2001. Later *P. Phani Bushan Rao and N. Ravi Shankar* [10] obtained ranking fuzzy numbers with a distance method using circumcenter of centroids and an index of modality. *D. Stephen Dinagar, K. Latha* [5] studied some types of type-2 triangular fuzzy matrices. Representation and ranking of fuzzy numbers

with heptagonal membership function using value and ambiguity index developed by *K. Rathi and S. Balamohan* [7]. The solution of fuzzy game problem using triangular fuzzy number studied by *R. Senthil Kumar, S. Kumaraghuru* [11]. *A. Rahmani, F. Hosseinzadeh Lotfi, M. Rostamy-Malkhalifeh, and T. Allahviranloo* [1] applied New Method for Defuzzification and ranking of fuzzy numbers based on the statistical beta distribution. A new approach for ranking fuzzy numbers developed by *M. Lotfi, S. Salahshour, F. Nasr Esfahani, A. Jafarnejad* [8]. Ordering of generalised trapezoidal fuzzy numbers based on area method using euler line of centroids introduced by *A. Hari Ganesh and M. Suresh* [10].

Basic Definitions

Triangular Fuzzy Number:

A triangular fuzzy number \bar{A} or simply triangular number represented with three points as follows (a_1, a_2, a_3) holds the following conditions

- i) a_1 to a_2 is increasing function
- ii) a_2 to a_3 is decreasing function
- iii) $a_1 \leq a_2 \leq a_3$

Its membership function is defined as follows

$$\mu_A(x) = \begin{cases} \frac{(x-a_1)}{(a_2-a_1)} & a_1 \leq x < a_2 \\ 1 & x = a_2 \\ \frac{(a_3-x)}{(a_3-a_2)} & a_2 < x \leq a_3 \end{cases}$$




Figure 1: Triangular fuzzy number $[a_1, a_2, a_3]$

α - Cut for triangular fuzzy number:

For any $\alpha \in [0, 1]$ from

$$\frac{a_1^\alpha - a_1}{a_2 - a_1} = \alpha, \quad \frac{a_3 - a_3^\alpha}{a_3 - a_2} = \alpha$$

$$a_1^\alpha = (a_2 - a_1) \alpha + a_1, \quad a_3^\alpha = - (a_3 - a_2) \alpha + a_3$$

$$\text{Thus } \bar{A}_\alpha = [a_1^\alpha, a_3^\alpha] = [(a_2 - a_1) \alpha + a_1, - (a_3 - a_2) \alpha + a_3]$$

Operations on triangular fuzzy number:

Addition, Subtraction and Multiplication of any two triangular fuzzy numbers are also triangular fuzzy number. Suppose triangular fuzzy numbers \bar{A} and \bar{B} are defined as,

$$\bar{A} = (a_1, a_2, a_3) \text{ and } \bar{B} = (b_1, b_2, b_3)$$

i) Addition $\bar{A} (+) \bar{B} = (a_1, a_2, a_3) + (b_1, b_2, b_3)$
 $= (a_1 + b_1, a_2 + b_2, a_3 + b_3)$

ii) Subtraction $\bar{A} (-) \bar{B} = (a_1, a_2, a_3) - (b_1, b_2, b_3)$
 $= (a_1 - b_1, a_2 - b_2, a_3 - b_3)$

iii) Symmetric image: $(-\bar{A}) = (-a_3, -a_2, -a_1)$

iv) Multiplication:

$$\bar{A} \times \bar{B} = (\min(a_1b_1, a_1b_3, a_3b_1, a_3b_3), a_2b_2, \max(a_1b_1, a_1b_3, a_3b_1, a_3b_3)).$$

Proposed Ordering (Ranking) of two Triangular fuzzy numbers:

The proposed method is based on the area of Trapezium, area of rectangle and centroid of triangle formed due to α -cut taken at values of $\alpha = 0.5$ respectively of triangular fuzzy number. The calculation of the proposed area presented from the point of view of analytical geometry. Given triangular fuzzy number $[G \ H \ I]$, is divided area of rectangle (CDKJ) using $\alpha = 0.5$. Suppose $\bar{X} = (x_1, x_2, x_3)$ and $\bar{Y} = (y_1, y_2, y_3)$ be given any two triangular fuzzy number. Area = Length (JK) \times width (CJ). The width CJ is equal to value of $\alpha = 0.5$. i.e CJ = 0.5. To determine the value of JK, from Fig.(3) we can observe that length JK is α -cut value at 0.5. Therefore by using α -cut method the value of $J = (H - G) \alpha + H$ and value of $K = - (I - H) \alpha + I$.

Suppose the area of Rectangle of Triangle for triangular fuzzy number \bar{X} and \bar{Y} due to α - cut at $\alpha = 0.5$ are A_1, A_2, A_3 and B_1, B_2, B_3 Respectively. If $\frac{(A_1 + A_2 + A_3)}{3} \leq \frac{(B_1 + B_2 + B_3)}{3}$ then we can say that fuzzy numbers $\bar{X} \leq \bar{Y}$.

The aim for selecting the proposed area of reference is that the plane figures Rectangle are balancing points of each plane figure, of triangular fuzzy number.



Figure 2: α -cut of Triangular fuzzy number at 0.5

Figure 3: Area of Rectangle CDMN

Mat-lab Code:

```

clc %clear the previous output
format short
fprintf('Enter first triangular Fuzzy Number');
a1=input("");
a2=input("");
a3=input("");
A=[a1 a2 a3] %display the first triangular number

```

```

alp=0.5;
fprintf('Enter second triangular Fuzzy Number');
b1=input("");
b2=input("");
b3=input("");
B=[b1 b2 b3] %display the second triangular number
A1_alpha=[(a2-a1)*alp+a1, -(a3-a2)*alp+a3]
B1_alpha=[(b2-b1)*alp+b1, -(b3-b2)*alp+b3]
A1_area=(alp-0.2)*(A1_alpha(2)-A1_alpha(1)) %area of reactangle for first fuzzy number
B1_area=(alp-0.2)*(B1_alpha(2)-B1_alpha(1)) %area of reactangle for second fuzzy number
area1=A1_area %Summation of three value obtained at 0.5 for first fuzzy number
area2=B1_area %Summation of three value obtained at 0.5 for second fuzzy number
if(abs(area1-area2)<0.000001) %if condition to check order of two fuzzy number
fprintf('A=B\n');
else if(area1-area2>0)
fprintf('B<A\n');
else
fprintf('A<B\n');
end
end
    
```

Comparison of Various Ranking Method:

Here we compare proposed method of order triangular fuzzy number with others by using example. Suppose $\bar{A} = (0.1, 0.2, 0.3)$ and $\bar{B} = (0.3, 0.4, 0.9)$ be any given triangular fuzzy numbers that we want to order

Table 1: Comparison of existing method of ranking two fuzzy numbers with proposed method

Name Ranking Method		\bar{A}	\bar{B}	Ranking Order
New Method	$\alpha = 0.5$	0.2332	0.5455	
Yager	-	0.200	0.500	$\bar{A} \leq \bar{B}$
Chen	$\beta = 0.5$	-0.200	0.000	$\bar{A} \leq \bar{B}$
	$\beta = 1$	-0.200	0.000	$\bar{A} \leq \bar{B}$

Defuzzification:

If $\bar{x} = (a, b, c)$ be any given triangular fuzzy number then we use mean and center method to Defuzzify i.e.

$x = \frac{a+b+c}{3}$ and method $x = b$ respectively.

If $(-a, 0, a)$ is given fuzzy triangular number then its crisp value is zero.

Method for solving fuzzy transportation problem:

In this section, the crisp transportation problem is converted into fuzzy transportation problem by using Triangular, Pentagonal, and Heptagonal Fuzzy numbers. Three methods are used to obtain the fuzzy transportation cost Fuzzy North West Corner Method (FNWCM), Fuzzy Matrix Minima Method (FMMM) and Fuzzy Vogel's Approximation Method (FVAM).

Fuzzy Matrix Minima Method (FLCM):

Suppose there are m factories and n warehouses then transportation problem is usually represented in tabular form

Table 2: Fuzzy transportation problem

Destination Origin	\bar{D}_1	\bar{D}_2	\bar{D}_3		\bar{D}_n	Supply
\bar{O}_1	\bar{C}_{11}	\bar{C}_{12}	\bar{C}_{13}		\bar{C}_{1n}	\bar{A}_1
\bar{O}_2	\bar{C}_{21}	\bar{C}_{22}	\bar{C}_{23}		\bar{C}_{2n}	\bar{A}_2
\bar{O}_m	\bar{C}_{m1}	\bar{C}_{m2}	\bar{C}_{m3}		\bar{C}_{mn}	\bar{A}_m
Demand	\bar{B}_1	\bar{B}_2	\bar{B}_3		\bar{B}_n	$\sum_{i=1}^n \bar{B}_i = \sum_{j=1}^m \bar{A}_j$

Step 1: Decide the smallest fuzzy cost in fuzzy transportation table. Let it be \bar{c}_{ij} . Find $\bar{x}_{ij} = \text{minimum}(\bar{A}_i, \bar{B}_j)$. The following three conditions may arise:

Condition (i): If $\text{minimum}(\bar{A}_i, \bar{B}_j) = \bar{A}_i$ then allocate $\bar{x}_{ij} = \bar{A}_i$ in the NWC of $m \times n$ fuzzy transportation table. Ignore the i^{th} row to obtain a new fuzzy transportation table of order $(m-1) \times n$. Replace \bar{B}_j by $\bar{B}_j - \bar{A}_i$ in the obtained fuzzy transportation table. Go to Step 2.

Condition (ii): If $\text{minimum}(\bar{A}_i, \bar{B}_j) = \bar{B}_j$ then allocate $\bar{x}_{ij} = \bar{B}_j$ in the NWC of $m \times n$ fuzzy transportation table. Ignore the j^{th} row to obtain a new fuzzy transportation table of order $m \times (n-1)$. Replace \bar{A}_i by $\bar{A}_i - \bar{B}_j$ in the obtained fuzzy transportation table. Go to Step 2.

Condition (iii): If $\bar{A}_i = \bar{B}_j$ then either follow condition (i) or condition (ii) but not both simultaneously. Go to Step 2.

Step 2: Repeat Step 1 for the obtained fuzzy transportation table, until the fuzzy transportation table is reduced into a fuzzy transportation table of order 1×1 .

Step 3: Assign all \bar{x}_{ij} in the ij^{th} cell of the given fuzzy transportation table.

Step 4: The obtained IFBFS and initial fuzzy transportation cost are \bar{x}_{ij} and $\sum_{i=1}^m \sum_{j=1}^n \bar{x}_{ij} \bar{c}_{ij}$ respectively.

Numerical example:

Consider the following crisp transportation problem

Table 3: Crisp transportation problem

Origin \ Destination	\bar{c}_1	\bar{c}_2	\bar{c}_3	\bar{c}_4	Supply
\bar{o}_1	6	3	5	4	22
\bar{o}_2	5	9	2	7	15
\bar{o}_3	5	7	8	6	8
Demand	7	12	17	9	

Minimum Transportation Cost = 150

Table 4: FLCM for triangular fuzzy number

Origin \ Destination	\bar{d}_1	\bar{d}_2	\bar{d}_3	\bar{d}_4	Supply
\bar{o}_1	[5,6,7]	[2,3,4] (11,12,13)	[4,5,6] (-2,1,4)	[3,4,5] (8,9,10)	[21,22,23]
\bar{o}_2	[4,5,6]	[8,9,10]	[1,2,3] (14,15,16)	[6,7,8]	[14,15,16]
\bar{o}_3	[4,5,6] (6,7,8)	[6,7,8]	[7,8,9] (-2,1,4)	[5,6,7]	[6,8,10]
Demand	[6,7,8]	[11,12,13]	[16,17,18]	[8,9,10]	

Minimum Transportation Cost = [22, 36, 52] + [-8, 5, 24] + [24, 36, 50] + [14, 30, 48] + [24, 35, 48] + [-14, 8, 36] = [62, 150, 258] = **156.66**

Final computational value of fuzzy triangular number is **[-9, 0, 9] = 0**

Conclusion and Future Work

Proposed method in this article is simple to order any two triangular fuzzy numbers. This method can order normal and non-normal fuzzy triangular numbers. The mat-lab code is used to obtain to get exact ordering of triangular fuzzy number. This method gives a correct ranking order to the problems for decision making problem under uncertainty since it is easy to calculation and gives acceptable result. In this article the proposed method is applied to fuzzy transportation problem. The crisp transportation problem is converted into a fuzzy transportation problem by using various triangular fuzzy numbers and solved by

using FLCM (Fuzzy Least Cost Method) methods. Through the numerical example, we can conclude that minimum fuzzy transportation cost obtained from this method is optimum.

Bibliography:

- [1]. A Rahmani, F Hosseinzadeh Lotfi, M Rostamy-Malkhalifeh, T Allahviranloo. "A New Method for Defuzzification and Ranking of Fuzzy Numbers Based on the Statistical Beta Distribution." Hindawi Publishing Corporation Advances in Fuzzy Systems 2016 (2016): 1-9.
- [2]. Ashok S Mhaske, K L Bondar. "Fuzzy Transportation by Using Monte Carlo method." Advances in Fuzzy Mathematics 12 (2017): 111-127.
- [3]. Ashok S Mhaske, K L Bondar. "Fuzzy Database and Fuzzy Logic for Fetal Growth Condition." Asian Journal of Fuzzy and Applied Mathematics 03 . 03 (2015): 95-104.
- [4]. Ashok Sahebrao Mhaske, Kirankumar Laxmanrao Bondar. "Fuzzy Transportation Problem by Using Triangular, Pentagonal and Heptagonal Fuzzy Numbers With Lagrange's Polynomial to Approximate Fuzzy Cost for Nonagon and Hendecagon." International Journal of Fuzzy System Applications 9.1 (2020): 112-129.
- [5]. D. Stephen Dinagar, K. Latha. "SOME TYPES OF TYPE-2 TRIANGULAR FUZZY MATRICES." International Journal of Pure and Applied Mathematics 82 (2013): 21-32.
- [6]. K L Bondar, Ashok Mhaske. "Fuzzy Transportation Problem with Error by Using Lagrange's Polynomial." The Journal of Fuzzy Mathematics 24 (2016): 825-832.
- [7]. K. Rathi, S. Balamohan. "Representation and Ranking of Fuzzy Numbers with Heptagonal Membership Function Using value and Ambiguity Index." Applied Mathematical Sciences 8 (2014): 4309 - 4321.
- [8]. M Lotfi, S Salahshour, F Nasr Esfahani, A Jafarnejad. "A new approach for ranking fuzzy numbers." Journal of Fuzzy Set Valued Analysis 2016 (2016): 102-109.
- [9]. MARCIN DETYNIECKI, RONALD R. YAGER. "RANKING FUZZY NUMBERS USING α -WEIGHTED VALUATIONS." International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems 8 (2001): 573-592.
- [10]. P Phani Bushan Rao, N Ravi Shanka. "Ranking Fuzzy Numbers with a Distance Method using Circumcenter of Centroids and an Index of Modality." Hindawi Publishing Corporation Advances in Fuzzy Systems 2011 (2011): 1-8.
- [11]. R. Senthil Kumar, S. Kumaraghuru. "Solution of Fuzzy Game Problem Using Triangular Fuzzy Number." IJISSET - International Journal of Innovative Science, Engineering & Technology 2.2 (2015).