

CURATIVE ROLE OF CAFFEINE ON COPPER INDUCED HISTOPATOLOGICAL ASSESSMENT OF HEPATOPANCREATIC CELLS IN FRESH WATER BIVALVE, *LEMELLIDENS CONSOBRINUS*

P. R. Mahajan

Head, Department of Zoology, Sardar Vallabhabhai Patel Arts and Science College,

Ainpur, Tal - Raver, Dist – Jalgaon . 425509

Email : prmahajan1971@gmail.com

ABSTRACT

The present study describes antioxidant effect of caffeine (1,3,7-Trimethylxanthine) against copper sulphate induced histopathological alteration on hepatopancreas in an experimental model, the fresh water bivalve, *Lamellidens consobrinus*. The effect of CuSO₄ and caffeine were studied under three groups. Group A bivalves was maintained as control, Group B was exposed to chronic LC_{50/10} doses of copper sulphate (2.12 ppm) for 18 days, while group C bivalves were exposed to respective chronic concentrations of heavy metals with 5 ml/lit caffeine. The changes are observed at 6, 12 and 18 days. In hepatopancreatic structure showed more disintegration of basement membrane due to damaged epithelial cells, disruption of hepatic tubules, and increase in internal luminal area in bivalves exposed to copper sulphate as compared to those exposed in copper sulphate along with caffeine. The probable antioxidant role of caffeine is discussed in the paper.

Key words - *Lamellidens consobrinus*, Copper sulphate, Hepatopancreas, Histopathological alterations, Caffeine

INTRODUCTION

Bivalves can take up contaminants from sediments, suspended particulate materials, water column and also food sources (Laffon et al, 2006 and Livingstone, 1993). They have been widely used for many years as bioindicator organisms in monitoring of chemical pollutants and biomonitoring in aquatic ecosystems. This is particularly due to their sedentary nature or immobility, filter-feeding activity, low metabolism, contact with sediments, suitable size for biochemical analysis, wide distribution in marine, estuarine and freshwater environments, practicality in collection, ability to bioaccumulate pollutants and high tolerance to chemical exposure due to a remarkably active immune system (Emmanouil, Kyriotakis, Kungolos, & Machera, 2008; Gupta & Singh, 2011)

Copper sulphate is frequently used as an algacide in commercial and recreational fish ponds to control growth of phytoplankton and filamentous algae, and to control certain fish diseases (Tucker, 1990). Consequently, treating plankton with copper compounds might lead to copper bioaccumulation reaching a toxic stage in fish. The toxic effect of copper is correlated to its aptitude for catalyzing oxidative reactions, leading to the production of reactive oxygen species (Lopes et al, 2001). Contamination of fresh water with a wide range of pollutants has become a matter of concern over last few decades (Vutukuru, 2005). Heavy metals have devastating effects on ecological balance of

the recent environment and a diversity of aquatic organisms (Farombi et.al, 2007). Heavy metals are very difficult to remove from body; the damage of tissues caused by heavy metals may be recovered. Various antioxidants are used for recovery or reduce the damage of tissue due to heavy metals. Vitamin C and E are common antioxidants in the diet.

The main constituent of coffee is caffeine. The caffeine molecule is a bitter alkaloid, which contributes to both acidity as well as the bitter properties of coffee. Caffeine is found to have antioxidant activity. This activity of caffeine can protect the damage of tissues, biochemicals and genetic materials of organisms from the heavy metal generated free oxygen radicals. Dissolved heavy metal ions are positively charged and caffeine contains uncharged and negatively charged groups. The metal ions might be taken out of solution by binding to negatively charged groups of caffeine in the coffee granules. The molecules of caffeine being small (Mole wt.193.), its chelate with heavy metal can be easily excreted out by the biological system. This property of caffeine indicates that caffeine can have the capacity to remove the heavy metals from the living organism, and prevent the damage of tissues. Coffee is the most widely consumed natural beverage by the people around the world. It contains caffeine as major bioactive constituent along with caffeic acid, chlorogenic acid, kahweol palmitate, and cafestol palmitate as trace amounts. Caffeine acts as neurostimulator and exerts protective effect against genotoxic/carcinogenic activity of environmental chemicals in *in vitro* and *in vivo* assay system (Ferguson, 1994; Abraham, 1984; Aeschbacher and Jaccaud, 1990; Stavric, 1992). The aim of our study was to determine the histological effect in the hepatopancreatic cells of freshwater bivalves, *Lamellidens consobrinus* after heavy metal stress.

MATERIALS AND METHODS

Chronic study

Lamellidens consobrinus are the fresh water bivalves occurring abundantly in Hartala pond at Hartala, Dist-Jalgaon of Maharashtra. For acute studies LC₅₀ (lethal concentration with 50% mortality) was calculated for 96 hours (APHA, 2005) by probit regression analysis (Finney, 1971). Water was replaced every day and fresh toxicant was added after water renewal for 96/10 hours. Bivalves were monitored at regular intervals and dead bivalves were removed from aquaria. A total of 30 mussels were collected and divided into three groups of 10 each. The first group A was consisted by control bivalves while the second B group was exposed to LC_{50/10} concentration of 96 hrs. CuSO₄ (2.12 ppm) and group C bivalves exposed to LC_{50/10} concentration of 96 hrs. CuSO₄ (2.12 ppm) along with 5 mg/lit. Caffeine upto 18 days.

Histological studies


The experimental bivalves from A, B and C group were dissected after 6, 12, and 18 days. The hepatopancreas from all experimental group were removed and fixed in Bouin's fluid for 24 hours; washed and dehydrated in alcohol grades, cleared in toluene and embedded in paraffin wax (58-60°C). Serial sections of 5 µ thickness were cut and stained with Mallory's triple stain. The stained sections were examined under light microscope for histopathological impact of heavy metal salts. Hepatopancreas of bivalves from all groups of control and exposed were screened and data is presented and compared.

Tissue processing

The removed wet tissue of digestive gland from all groups were homogenate in blender with M/150 phosphate buffer at 1-4°C and centrifuge. Stir sediment with cold phosphate buffer and allow standing in the cold with shaking occasional then repeating the extraction once or twice and using the supernatant for assay of catalase.

OBSERVATION AND RESULTS

The accumulation of heavy metals in the body of the organism leads to the formation of histological lesions in the organs. Chronic exposure of *Lamellidens consobrinus* to copper sulphate causes severe histological lesions in the hepatopancreas as compared to those exposed to LC_{50/10} concentration of CuSO₄ with 5 mg/lit. Caffeine and control group of bivalves.



Normal histological structure of hepatopancreas









Fig-B/6,B/12 and B/18

Fig-C/6,C/12 and C/18

Exposed to LC50/10 conc. of CuSO₄

LC50/10 conc. of CuSO₄+5mg/lit.caffeine

CT- Control tissue, BL – Blood lacunae, CC – Calciferous cell, AC – Absorptive cell, L – Lumen, BM –Basement membrane

Histological study shows different types of cells in hepatic lobules, the typical liver cells or digestive cells (columnar epithelial cells) and secretory cells, absorptive cells, blood lacunae. The cells rest on basement membrane. The lumen present inside the lobules. The lobules of the gland are bound together by the thin connective tissue layer as shown in fig. A. The deformity in the cytoarchitecture of hepatopancreas due to heavy metal stress definitely reflect in its metabolic activity. The histological changes in the hepatopancreas of *Lamellidens consobrinus* exposed to LC_{50/10} concentration, 2.12 ppm for 6 days, 12 days and 18 days of CuSO₄, show in fig. B/6, B/12 and B/18. The deformity in the histologic structure of digestive gland is more in 18 days exposure group of bivalves as compared to those bivalves exposed to LC_{50/10} concentration of CuSO₄ with 5mg/lit. caffeine and control group.

DISCUSSION

The alterations observed in histo-architecture in the various tissues are tissue specific and time dependent. When *Channa punctatus* was exposed to mercuric chloride for a prolonged span of 30 days profound histological changes in the liver were observed which induced necrosis, vacuolation and degeneration of hepatocytes (Sastry and Gupta, 1978). The histological techniques are the promising area of research in aquatic toxicology as it gives the real picture of the effects imposed and the involvement of the xenobiotics in either disturbing or destroying the vital organs of living organisms. Many workers have reported the degenerative changes in selected tissues of the animals in response to pollution by various toxicants (Shaikh, *et al.*, 2010; Andhale, *et al.*, 2011). Victor *et al.* (1990) observed histopathological changes in the hepatopancreas of *P. hydrodromous* in response to cythion resulting in reduction in the height of tubular epithelium, enlargement of lumen, vacuolation and atrophy. The histopathological changes indicated that the animals were not able to digest and store food properly.

The exposure of the bivalves to LC_{50/10} concentrations of heavy metals (CuSO₄) caused varying degrees of histological alterations in the organs examined hepatopancreas. In the hepatopancreas, there was a prevalence of hepatocellular foci of cellular alterations (FCA) in bivalves exposed to LC_{50/10} concentrations of copper sulphate. Peripheral thickening and inflammation of hepatic tubules of the hepatocytes were also observed in exposed bivalves. Less damage in histological structure in bivalves those exposed to LC_{50/10} concentration of copper sulphate with 5 mg/lit. caffeine. Over all study indicates that, caffeine has a protective and curative role in the heavy metal induced alterations. The damages in the histological structure of hepatopancreas are less in presence of caffeine and the recovery from the heavy metal induced damage in the structure of hepatopancreas was faster in presence of caffeine.

Caffeine has antioxidant activity. This activity of caffeine can protect tissues from heavy metal induced free radical damage of the tissues. Shomer (1994) studied caffeine stimulation of malignant hyperthermia-susceptible sarcoplasmic reticulum Ca²⁺ release, and suggested that caffeine sensitivity of malignant hyperthermia-susceptible (MHS) skeletal muscle fiber bundles is due to an altered caffeine sensitivity of the MHS calcium ion release channel protein. Chung Fung- Lung (1999) reported that caffeine when given in drinking water at a concentration identical to that found in 2% tea was able to inhibit lung tumours induced by 4-(Methylnitrosoamino)-1-(3-pyridyl)-1 butanol (NNK). Hosta *et al.*; (2001) has studied hepatocarcinogenesis inhibition by caffeine in AGI rats treated with 2-

acetylaminofluorene and has showed that caffeine inhibited hepatocarcinogenesis induced by 2-acetylaminofluorene. Break of DNA strands in hepatopancreas of mussels exposed to resin acids was reported by Gravato et al. (2005).

The results of the present study indicated the positive role in detoxification of heavy metals by caffeine. Its antioxidant property, capacity to chelate the heavy metal and induced increase in urine formation may be the cause of hepatoprotection.

ACKNOWLEDGEMENT

The authors are thankful to the Principal, Dhanaji Nan Mahavidyalaya, Faizpur for providing the laboratory facility to carry out the work.

REFERENCES

1. **Abraham SK.(1989).** Inhibition of *in vivo* genotoxicity by coffee. *Food Chem Toxicol*, 27:787-92.
2. **Aeschbacher HU, Jaccaud E(1990).** Inhibition by coffee of nitrosourea-mediated DNA damage in mice. *Food Chem Toxicol*, 28:633-7.
3. **Andhale, A.V., Bhosale, P.A. and Zambare, S.P.(2011):** Histopathological study of nickel induced alterations in the fresh water bivalve, *Lammellidens marginalis*, *Journal of Experimental Sciences* Vol. 2, Issue 4: 01-03.
3. **Chung Fung-Lung (1999):** The prevention of lung cancer induced by a tobacco-specific carcinogen in rodents by green and black tea. *Proc. Soc. Exp. Biol. Med.* 220 -244.
- 4 **Emmanouil, C., Kypriotakis, S., Kungolos, A., & Machera, K. (2008):** Effects of Thiram or MCPA acid on mussel gill DNA. Discussion Paper Series, 14(24), 455e468.
5. **Finney, D.J. (1971):** Probit Analysis, 3rd edition, London and New York, *Cambridge University Press*, pp. 333.
6. **Farombi, E. O., Adelowo, O. A. and Ajimoko, Y. R.(2007):** Biomarker of oxidative stress and heavy metal levels as induced by environmental pollution in African cat fish *Clarias gariepinus* from Nigeria Ogun river. *Int. J. Environ. Res. Public Health*, 4 (2): 158-165.
7. **Ferguson LR (1994).** Antimutagens as cancer chemopreventive agent in the diet. *Mutat Res*, 307:395-410.
8. **Gravato, C., Oliveira, M. and Santos, M.A.(2005):** Oxidative stress and genotoxic responses to resin acids in Mediterranean mussels. *Ecotoxicol. Environ. Saf.*, 61(2): 221-229.
9. **Gupta, S. K., & Singh, J. (2011):** Evaluation of mollusc as sensitive indicator of heavy metal pollution in aquatic system: a review. *IIOAB Journal*, 2,49e57.
10. **Hosaka, S; Kawa, S; Aoki, Y; Tanaka, E; Yoshizawak; Karasawa, Y; Hosaka, N. and Kiyosawak (2001):** Hepatopantocarcinogenesis inhibition by caffeine in ACI rats treated with 2-acetylaminofluorene, *food chem Toxicol*, (39): 557.
11. **Laffon, B., Rabade, T., Pasaro, E., & Mendez, J. (2006):** Monitoring of the impact of Prestige oil spill on *Mytilus galloprovincialis* from Galician coast. *Environment International*, 32, 342e348.
12. **Lopes, P.A, Pinheiro, T, Santos, M.C, Mathias, M, Collares-Pereira, M.J, et al. (2001):** Response of antioxidant enzymes in freshwater fish populations (*Leuciscus alburnoides* complex) to inorganic pollutants exposure. *Sci Total Environ* 280: 153-163.
13. **Sastry, K.V. and Gupta, P.K. (1978):** Effect of mercuric chloride on the digestive system of *Channa punctatus*, A histopathological study, *Environ. Res.* (16): 270-278.
14. **Shaikh, F.I, Ustad, I.R. and Ansari, N.T. [2010]:** Effect of heavy metal on the ovary of freshwater crab, *Barytelphusa cunicularis* [Westwood]. *The Bioscan*, 5[2]: 335-338.
15. **Shomer, N.H; and Nickelson, J.R. (1994):** Caffeine stimulation of malignant hyperthermia susceptible sarcoplasmic reticulum Ca^{2+} release, *American Journal of Physiology*, Vol. 267(5): PC 1253.

- 16. Stavric B(1989):** An update on research with coffee/caffeine (1989-1990).Food Chem Toxicol 1992;30:533-55.
- 17.Tucker, C.S and Robinson, E.H. (1990):** Channel Catfish Farming Handbook. Van-Nostrand-Reinhold, New York, USA.
- 18.Vutukuru, S. S. (2005):** Acute effect of Hexavalent chromium on survival, oxygen consumption, haematological parameter and some biochemical profiles of the Indian Major carp, Labeo rohita. Int. J. Environ. Res. Public Health, 2 (3): 456-462

