

# “Awareness on Active Immunization”: A survey

Ankit<sup>1</sup>, S. Bose\*, Pooja Sharma<sup>2</sup>

\*School of Pharmaceutical Sciences, Lovely Professional University, Punjab  
2 Abhilashi College of Pharmacy, Nerchowk, Mandi

## Abstract

Vaccination is one of the most financially savvy general wellbeing mediations and to a great extent liable for decrease of under-5 death rate. In any case, antibody preventable sicknesses (VPDs) are as yet liable for more than 5 lakhs passings every year in India. This underlines the need of further improvement. There are various reasons why India falls behind its a lot less created neighbors in inoculation rates. They incorporate colossal populace with moderately high development rate, topographical decent variety and some hard to arrive at populaces, absence of mindfulness with respect to immunization, lacking conveyance of wellbeing administrations, insufficient checking, absence of miniaturized scale arranging and general absence of between sectoral coordination, and feeble VPD observation framework. Since it gives direct and viable security against preventable dreariness and mortality. It has been a significant benefactor in the decay of under-5 death rate from ~ 233 to ~63 (per 1000) in most recent five decades in India. This underlines the requirement for additional improvement. India, alongside many creating nations, is lingering behind in adequate inclusion of Routine Immunization (RI). Lack of mindfulness is one extraordinary hindrance to accomplish penny percent vaccination inclusion. A later report in 225 towns of Uttar Pradesh validated the way that absence of mindfulness is the one of the principle purpose behind incomplete vaccination/non-inoculation. Subsequently, the interest for antibodies likewise endures. Low levels of training contrarily sway wellbeing looking for conduct.

**Key words: Immunization, Vaccines, Awareness.**

## 1. Introduction

Immunology is a science that looks at the structure and capacity of the insusceptible framework. It starts from medication and early investigations on the reasons for insusceptibility to ailment. From early works, plainly crude man thought about sickness and its assaults. One finds in the Babylonian Epic of Gilgamesh (2000 B.C. – Mesopotamian saint) records of the nearness of epidemic and infection. In other, later compositions from old administrations of old Egypt, one finds much more portrayals of illness. Further, one can even recognize the malady of which they talked. Review that, back then, ailment and disease was discipline rendering because of "terrible deeds" or "underhandedness considerations". Indeed, even the old confirmation is loaded up with plague that God fashioned upon the individuals who "crossed" him. From these works, it is similarly evident that man realized that once he had been harrowed with sickness, on the off chance that he endure, he was ordinarily not ready to contract it once more. [1]

### 1.1. Oral Polio Vaccine

Worldwide poliomyelitis destruction was proposed in 1988 by the World Health Organization to its part states. The key arrangement built up the exercises required for polio annihilation, affirmation for districts, OPV discontinuance stage and post-OPV stage. OPV is the antibody of decision for the poliomyelitis annihilation program since it prompts both a foundational and mucosal insusceptible reaction. [2]

### 1.2. DPT Vaccine

The antibody expands the insurance that individuals should as of now have from their standard youth inoculations. All young people ought to get one "promoter" portion of Tdap at 14 to 16 years old. [3, 7]

### 1.3. BCG Vaccine

BCG is an antibody against tuberculosis (TB). It is named after the two individuals who found the antibody (Bacillus Calmett-Guerin). It utilizes a live TB germ that has been adjusted in the research center with the

goal that it won't cause illness in a sound individual. BCG inoculation decreases the danger of TB in individuals who are not effectively contaminated with TB. The immunization doesn't generally forestall TB infection. BCG inoculation isn't prescribed for grown-ups on the grounds that the advantage is substantially less. Insurance against TB will just beginning 6 two months after the immunization has been given and goes on for around 10 years. Revaccination isn't prescribed. [4, 8]

#### 1.4. MMR Vaccine (Measles, Mumps, Rubella Vaccine)

Measles, mumps, and rubella are not kidding sicknesses. Before immunizations they were exceptionally normal, particularly among kids. The danger of MMR immunization causing genuine damage, or passing, is very little. Getting MMR antibody is a lot more secure than getting measles, mumps or rubella. The vast majority who get MMR antibody don't have any difficult issues with it. [9]

#### 1.5. Hepatitis Vaccine

"Hepatitis" signifies aggravation of the liver. The Hepatitis B infection (HBV) is an infection transmitted in blood, semen and salivation. It was found in 1965 in Australia (it was initially named the Australia Antigen after its revelation in Indigenous Australians). Right now, somewhere in the range of 90,000 and 160,000 individuals in Australia are incessantly contaminated with HBV. [5, 6]

## 2. Survey

### 2.1. Questionnaire for Awareness on Active Immunization

1. Have you ever hear about vaccination?

- a. Yes
- b. No
- c. Don't know

2. From where did you hear about vaccination?

- a. Television program
- b. Media
- c. Internate
- d. Radio
- e. From maternity hospital

3 .Are you aware about vaccination of your child?

- a. Yes
- b. No

4. What is the Current age of your child.\_\_\_\_\_?

5. At what age you have started vaccination of your baby?

- i. 60 days

ii. 120 days

iii. 180 days

iv. 240 days

6. Has your baby immunized by OPV?

i. Agree

ii. Disagree

iii. Don't know

7. Has your baby immunized by BCG?

i. Agree

ii. Disagree

iii. Don't know

8. Has your baby immunized by DPT?

i. Agree

ii. Disagree

iii. Don't know

9. Has your baby immunized by hepatitis-B vaccine?

i. Agree

ii. Disagree

iii. Don't know

10. Has your baby immunized by measles vaccine?

i. Agree

ii. Disagree


iii. Don't know

11. Has your baby immunized by Hib –vaccine?

i. Agree

ii. Disagree

iii. Don't know



12. Has your baby given Vitamin-A solution?

- i. Agree
- ii. Disagree
- iii. Don't know

Form completed by: \_\_\_\_\_

date: \_\_\_\_\_

Form received by: \_\_\_\_\_

date: \_\_\_\_\_


## 4. Result and Discussion

### 4.1. Immunisation status

Survey includes responses of 50 families. Nearly all parents responded that their children were up to date with immunization. On the basis of response submitted by different families it was found that 60% of them were aware of immunization, 30% were not aware and 10% were not sure regarding the same as represented in table 1 and figure 2.

**Table no. 1; Children's immunization status**

Immunisation Status	No. Of Persons	Percentage
Aware	30	60
Not aware	15	30
Not sure	5	10



**Fig no.2; Children's immunization status**


### 4.2. Sources of Immunisation

According to the survey main information source of immunization is television. Most of the parents got the information regarding immunization from television and maternity hospitals. As per the survey 40% of the

parents got the information from television, 30% got the information from maternity hospitals, 16% informed by media, 12% were gathered information from internet and 2% were heard from radio as discussed in table 2 and figure 10.

**Table no.2; Sources of Immunisation information**

Source of information	No. of Persons	Percentage
Television	20	40
Maternity hospital	15	30
Media	8	16
Internet	6	12
Radio	1	2



**Fig no.2; Sources of Immunisation information**


**4.3. Awareness on vaccines**

From the survey it was found that most of the parents were aware of Oral Polio vaccine, whereas, the parents were not aware of Hib vaccines. According to the survey 100% parents were aware of OPV, 86% were aware of BCG vaccine, 74% were aware of DPT and Vitamin A solution, 20% were of Hepatitis B vaccine, 34% were aware of Measles and 8% were aware of Hib vaccine as per the data given in table no 3 and figure 3.

**Table no. 3; Awareness on vaccines**

Vaccines	Aware (%)	Not Aware (%)	Not Sure (%)
Oral polio vaccines	100	0	0
BCG	86	8	6
DPT	74	20	6

Vitamin A solution	74	10	16
Hepatitis B vaccine	40	30	30
Measles	34	50	16
Hib vaccines	8	26	66



**Fig no 13; Awareness on vaccines**

## 5. Conclusion

Depending on the survey report it is concluded that, many parents are aware for vaccination of their children. Mostly they got the information on immunization from Television. From our survey, it is concluded that almost all the parents are aware of Oral Polio vaccination. Few of the parents (40%) still are not aware of immunization due to lack of information provided to them. A Low literacy level of parents is the matter of discuss. Some of them don't know about the diseases for which their child is being immunized. Also many parents don't know the timings of vaccination.

According to the survey report it was found that most of the parents are aware of Oral Polio vaccine, whereas they are not aware of some of the vaccines like Hib vaccines, Measles, Hepatitis B etc. As per the study it is concluded that still awareness is required for the proper immunization or vaccination among the parent for their children to complete eradication of disease caused due to lack of vaccination.

## REFERENCES

- [1]. Abdul Nasser Kaadan , Mohammed Nour Alsayyed Ali., 'The History of Immunology and Vaccines'. Doc, 8-9
- [2]. Ashtiyani, Amoozandeh, Rhazes., Diagnostic Differentiation of smallpox and Measles, Iranian Red Crescent Medical Journal, 2005; 43 (RR-1): 1-38.
- [3]. CDC, Tetanus surveillance—United States, 1998–2000. MMWR 2003;55-56.
- [4]. Houston S, et al., The effectiveness of Bacillus Calmette- Guérin (BCG) vaccination against Tuberculosis. Am J Epidemiol 1990;131:340-34.
- [5]. Jamison DT et al., eds. Disease control priorities in developing countries, 2nd ed. New York,

Oxford University Press, 2006;105-108.

[6]. Hepatitis A vaccines, Weekly Epidemiological Record, 2000; 75(5):38–40.

[7]. CDC, Diphtheria, tetanus, and pertussis, Recommendations for vaccine use and other preventive measures. Recommendations of the Immunization Practices Advisory Committee (ACIP). MMWR 1991;40 (No. RR-10):1–15.

[8]. CDC, The role of BCG vaccine in the prevention and control of tuberculosis in the United States. MMWR 1996;45(RR-4):1-18.

[9]. Measles vaccines, Weekly Epidemiological Record, 2004; 130–135.

