

P₄ - Decomposition in Boolean Function Graph $B(\overline{K_p}, L(G), \text{NINC})$ of a graph

S. Muthammai

Principal(Retd), Alagappa Government Arts college, Karaikudi, Tamil Nadu, India.

S. Dhanalakshmi

Government Arts College for Women (Autonomous), Pudukkottai, Tamil Nadu, India.

Abstract

For any graph G , let $V(G)$ and $E(G)$ denote the vertex set and edge set of G respectively. The Boolean function graph $B(\overline{K_p}, L(G), \text{NINC})$ of G is a graph with vertex set $V(G) \cup E(G)$ and two vertices in $B(\overline{K_p}, L(G), \text{NINC})$ are adjacent if and only if they correspond to two adjacent edges of G or to a vertex and an edge not incident to it in G . For brevity, this graph is denoted by $B_2(G)$. In this paper, P_4 -decomposition of Boolean Function Graph $B(\overline{K_p}, L(G), \text{NINC})$ of some standard graphs and corona graphs are obtained.

Keywords: Boolean Function graph, Edge Domination Number, Decomposition

1. Introduction

Graphs discussed in this paper are undirected and simple graphs. For a graph G , let $V(G)$ and $E(G)$ denote its vertex set and edge set respectively. A graph with p vertices and q edges is denoted by $G(p, q)$. A subset $F \subseteq E(G)$ is called an edge dominating set of G , if every edge not in F is adjacent to some edge in F . The edge domination number $\gamma'(G)$ of G is the minimum cardinality taken over all edge dominating sets of G . The corona $G_1 \circ G_2$ of two graphs G_1 and G_2 is defined as the graph obtained by taking one copy of G_1 (which has p_1 vertices) and p_1 copies of G_2 , and then joining the i^{th} vertex of G_1 to every vertex of in the i^{th} copy of G_2 . For any graph G , $G \circ K_1$ is denoted by G^+ .

A decomposition of a graph G is a family of edge-disjoint subgraphs $\{G_1, G_2, \dots, G_k\}$ such that $E(G) = E(G_1) \cup E(G_2) \cup \dots \cup E(G_k)$. If each G_i is isomorphic to H , for some subgraph H of G , then the decomposition is called a H -decomposition of G . In particular, a P_4 -decomposition of a graph G is a partition of the edge set of G into paths of length 3. In this case, G is said to be P_4 -decomposable. Several authors studied various types of decomposition by imposing conditions on G_i in the decomposition. Heinrich, Liu and Yu[3] proved that a connected 4-regular graph admits a P_4 -decomposition if and only if $|E(G)| \equiv 0 \pmod{3}$. Sunil Kumar[10] proved that a complete r - partite graph is P_4 -decomposable if and only if its size is a multiple of 3. P.Chithra devi and J. Paulraj Joseph [1] gave a necessary and sufficient condition for the decomposition of the total graph of standard graphs and corona of graphs into paths on three edges. Janakiraman et al., introduced the concept of Boolean function graphs [4 - 6]. For a real x , $\lfloor x \rfloor$ denotes the greatest integer less than or equal to x . For any graph G , let $V(G)$ and $E(G)$ denote the vertex set and edge set of G respectively. The Boolean function graph $B(\overline{K_p}, L(G), \text{NINC})$ of G is a graph with vertex set $V(G) \cup E(G)$ and two vertices in $B(\overline{K_p}, L(G), \text{NINC})$ are adjacent if and only if they correspond to two adjacent edges of G or to a vertex and an edge not incident to it in G . For brevity, this graph is denoted by $B_2(G)$.

In this paper, P_4 -decomposition of Boolean Function Graph $B(\overline{K_p}, L(G), \text{NINC})$ of some standard graphs are obtained.

2. Prior Results

Observation 2.1. [4]

Let G be a graph with p vertices and q edges.

1. $L(G)$ is an induced subgraph of $B_2(G)$ and the subgraph of $B_2(G)$ induced by vertices of G in $B_2(G)$ is totally disconnected.
2. Number of vertices in $B_2(G)$ is $p + q$ and if $d_i = \deg_G(v_i)$, $v_i \in V(G)$, then the number of edges in $B_2(G)$ is $q(p - 3) + \frac{1}{2} \sum_{1 \leq i \leq p} d_i^2$.
3. The degree of a vertex of G in $B_2(G)$ is q and the degree of a vertex e' of $L(G)$ in $B_2(G)$ is $\deg_{L(G)}(e') + p - 2$. Also if $d^*(e')$ is the degree of a vertex e' of $L(G)$ in $B_2(G)$, then $0 \leq d^*(e') \leq p + q - 3$.

3. Main Results

In the following, P_4 -decomposition of $B_2(P_n)$, $B_2(C_n)$, $B_2(K_{1,n})$ and corona graphs are found.

Theorem 3.1

- (i) For $n \geq 6$ and $n \equiv 0 \pmod{3}$, the graph $B_2(P_n) - (2n - 3)K_2$ is P_4 -decomposable.
- (ii) For $n \geq 7$ and $n \equiv 1 \pmod{3}$, the graph $B_2(P_n) - 2K_2$ is P_4 -decomposable.
- (iii) For $n \geq 5$ and $n \equiv 2 \pmod{3}$, the graph $B_2(P_n) - (n+1)K_2$ is P_4 -decomposable.

Proof: Let $V(P_n) = \{v_1, v_2, \dots, v_n\}$ and $e_i = (v_i, v_{i+1})$, $i = 1, 2, \dots, n-1$, be the edges of P_n . Then $v_1, v_2, \dots, v_n, e_1, e_2, \dots, e_{n-1} \in V(B_2(P_n))$. $B_2(P_n)$ has $2n-1$ vertices and $n^2 - 2n$ edges. It is to be noted that, in all the sets the suffix i in v_i is integer modulo n , and j in e_j is integer modulo $n-1$, $v_0 = v_n$ and $e_0 = e_{n-1}$.

Let $F = \bigcup_{i=1}^{n-1} F_i$, where $F_1 = \{(v_1, e_i), i = 2, 3, \dots, n-1\}$,

$F_i = \{(v_i, e_j), 1 \leq j \leq n-1 \text{ and } j \neq i-1, i\}$, $i = 2, 3, \dots, n-2$.

$F_{n-1} = \{(v_n, e_j), j = 1, 2, \dots, n-2\}$. $E(B_2(P_n)) = E(L(P_n)) \cup F = E(P_{n-1}) \cup F$.

Case 1. $n \equiv 0 \pmod{3}$.

Then the edge set of $B_2(P_n)$ can be decomposed into $((n^2 - 4n + 3)/3)P_4$ and $(2n - 3)K_2$. The edge set of $(2n - 3)K_2$ is given by the set $\{(v_1, e_{n-1})\} \cup \{\bigcup_{i=1}^{n-2} (v_i, e_i) (e_i, e_{i+1})\}$. The edge sets of $((n^2 - 4n + 3)/3) P_4$ are given by the edge sets $A^{(1)}$, where $A^{(1)} = \bigcup_{i=1}^{n-1} (\bigcup_{j=1}^{(n-3)/3} A_{j,i}^{(1)})$ and $A_{j,i}^{(1)} = \{(v_i, e_{i+j}), (e_{i+j}, v_{i+3j}), (v_{i+3j}, e_{i+j-1})\}$. Here, $\langle A^{(1)} \rangle \cong ((n-1)(n-3)/3) P_4$. Therefore, $B_2(P_n) - (2n - 3)K_2$ is P_4 -decomposable.

Case 2. $n \equiv 1 \pmod{3}$.

Then the edge set of $B_2(P_n)$ can be decomposed into $((n^2 - 2n - 2)/3)P_4$ and $2K_2$. The edge set of $2K_2$ is given by the edge set $\{(v_1, e_{n-1}), (v_{(2n+1)/3}, e_1)\}$. The edge sets of $((n^2 - 2n - 2)/3)P_4$ are given by the edge sets $A^{(2)}$ and $A^{(3)}$, where $A^{(2)} = \bigcup_{i=1}^{n-1} (\bigcup_{j=1}^{(n-4)/3} A_{j,i}^{(2)})$, where $A_{j,i}^{(2)} = \{(v_i, e_{i+j}), (e_{i+j}, v_{i+3j}), (v_{i+3j}, e_{i+j-1})\}$ and $A^{(3)} = \bigcup_{i=1}^{n-2} A_i^{(3)}$, $A_i^{(3)} = \{(v_n, e_i), (e_i, e_{i+1}), (e_{i+1}, v_{i+(2n+1)/3})\}$. Here, $\langle A^{(2)} \rangle \cong ((n-1)(n-4)/3) P_4$ and $\langle A^{(3)} \rangle \cong (n-2) P_4$. Therefore, $B_1(P_n) - 2K_2$ is P_4 -decomposable.

Case 3. $n \equiv 2 \pmod{3}$

Then the edge set of $B_2(P_n)$ can be decomposed into $((n^2 - 3n - 1)/3) P_4$ and $(n+1)K_2$. The edge set of $(n+1)K_2$ is given by the set $\{(v_{n-4}, e_1), (v_1, e_{n-1})\} \cup (\bigcup_{i=1}^{n-1} \{(v_i, e_{i+1})\})$. The edge sets of $((n^2 - 3n - 1)/3) P_4$ are given by the edge sets $A^{(4)}$ and $A^{(5)}$, where

$A^{(4)} = \bigcup_{i=1}^{n-1} (\bigcup_{j=1}^{(n-5)/3} A_{j,i}^{(4)})$ and $A_{j,i}^{(4)} = \{(v_i, e_{i+j+1}), (e_{i+j+1}, v_{i+2j+2}), (v_{i+2j+2}, e_{i+j-1})\}$ and

$A^{(5)} = \bigcup_{i=1}^{n-2} A_i^{(5)}$, $A_i^{(5)} = \{(v_n, e_i), (e_i, e_{i+1}), (e_{i+1}, v_{i+(n-4)})\}$. Here, $\langle A^{(4)} \rangle \cong ((n-1)((n-5)/3) P_4$ and $\langle A^{(5)} \rangle \cong (n-2) P_4$. Therefore, $B_2(P_n) - (n+1)K_2$ is P_4 -decomposable.

Theorem 3.2

- (i) For $n \geq 6$ and $n \equiv 0 \pmod{3}$, the graph $B_2(C_n) - 2nK_2$ is P_4 -decomposable.
- (ii) For $n \geq 4$ and $n \equiv 1 \pmod{3}$, the graph $B_2(C_n)$ is P_4 -decomposable.
- (iii) For $n \geq 5$ and $n \equiv 2 \pmod{3}$, the graph $B_2(C_n) - nK_2$ is P_4 -decomposable.

Proof: Let $V(C_n) = \{v_1, v_2, \dots, v_n\}$ and $e_i = (v_i, v_{i+1})$, $i = 1, 2, \dots, n-1$, be the edges of C_n . Then $v_1, v_2, \dots, v_n, e_1, e_2, \dots, e_{n-1}, e_n \in V(B_2(C_n))$. $B_2(C_n)$ has $2n$ vertices and $n^2 - n$ edges. It is to be noted that, in all the sets the suffices i in v_j and j in e_j is integer modulo n , $v_0 = v_n$ and $e_0 = e_n$. Let $F_n = \bigcup_{j=1}^n F_j$, where $F_j = \{(v_i, e_{i+j}), i = 1, 2, \dots, n-1\}$.

$E(B_2(C_n)) = E(L(C_n)) \cup F = E(C_n) \cup F$.

Case 1. $n \equiv 0 \pmod{3}$.

Then the edge set of $B_2(C_n)$ can be decomposed into $((n^2 - 3n)/3) P_4$'s and $2nK_2$. The edge set of $2nK_2$ is given by the set $\{\bigcup_{i=1}^n (v_i, e_{i+1}) \cup (e_i, e_{i+1})\}$. The edge sets of $((n^2 - 3n)/3)P_4$ are given by the edge sets $B^{(1)}$, where $B^{(1)} = \bigcup_{i=1}^{n-1} (\bigcup_{j=1}^{(n-3)/3} B_{j,i}^{(1)})$, where $B_{j,i}^{(1)} = \{(v_i, e_{i+j+1}), (e_{i+j+1}, v_{i+3j+1}), (v_{i+3j+1}, e_{i+j})\}$. Here, $\langle B^{(1)} \rangle \cong ((n)(n-3)/3)P_4$. Therefore, $B_2(C_n) - 2nK_2$ is P_4 -decomposable.

Case 2. $n \equiv 1 \pmod{3}$.

Then the edge set of $B_2(C_n)$ can be decomposed into $((n^2 - n)/3) P_4$. The edge sets $((n^2 - n)/3)P_4$ are given by the edge sets $B^{(2)}$ and $B^{(3)}$, where $B^{(2)} = \bigcup_{i=1}^n (\bigcup_{j=1}^{(n-4)/3} B_{j,i}^{(2)})$, where $B_{j,i}^{(2)} = \{(e_{i+j+1}, v_i), (v_i, e_{i+2j+2}), (e_{i+2j+2}, v_{i+j+1})\}$ and $B^{(3)} = \bigcup_{i=1}^n B_i^{(3)}$, $B_i^{(3)} = \{(e_{i+1}, v_i), (v_i, e_{i+2}), (e_{i+2}, e_{i+1})\}$. Here, $\langle B^{(2)} \rangle \cong ((n(n-4)/3) P_4$ and $\langle B^{(3)} \rangle \cong nP_4$. Therefore, $B_2(C_n)$ is P_4 -decomposable.

Case 3. $n \equiv 2 \pmod{3}$

Then the edge set of $B_2(C_n)$ can be decomposed into $((n^2 - 2n)/3) P_4$ and nK_2 .

The edge set of nK_2 is given by the set $\bigcup_{i=1}^n \{(e_i, e_{i+1})\}$. The edge sets of $((n^2 - 2n)/3)P_4$ are

given by the edge set $B^{(4)}$, where $B^{(4)} = \bigcup_{i=1}^n (\bigcup_{j=1}^{(n-2)/3} B_{j,i}^{(4)})$ and

$B_{j,i}^{(4)} = \{(e_{i+2j-1}, v_i), (v_i, e_{i+2j}), (e_{i+2j}, v_{i+3j+1})\}$. Here, $\langle B^{(4)} \rangle \cong ((n^2 - 2n)/3)P_4$.

Therefore, $B_2(C_n) - nK_2$ is P_4 -decomposable.

Theorem 3.3

For $n \geq 3$, the graph $B_2(K_{1,n})$ is P_4 -decomposable.

Proof: Let $V(K_{1,n}) = \{v, v_1, v_2, \dots, v_n\}$, where v is the central vertex and $e_i = (v, v_i)$, $i = 1, 2, \dots, n$ be the edges of $K_{1,n}$. Then $v, v_1, v_2, \dots, v_n, e_1, e_2, \dots, e_n \in V(B_2(K_{1,n}))$. $B_2(K_{1,n})$ has $2n+1$ vertices and $(3n^2 - 3n)/2$ edges. In all the sets defined below, the suffices are integers modulo n and $v_0 = v_n, e_0 = e_n$.

Let $F = \bigcup_{j=1}^n F_j$, where $F_i = \{(v_i, e_j), \text{ for all } j, 1 \leq j \leq n \text{ and } j \neq i\}$.

$E(B_1(K_{1,n})) = E(L(K_{1,n})) \cup F = E(K_n) \cup F$.

Case 1. n is even

Then the edge set of $B_2(K_{1,n})$ can be decomposed into $((n^2 - n)/2) P_4$.

Let $C^{(1)} = \bigcup_{i=1}^n (\bigcup_{j=1}^{(n-2)/2} C_{j,i}^{(1)})$, where $C_{j,i}^{(1)} = \{(e_{i+2j-1}, v_i), (v_i, e_{i+2j}), (e_{i+2j}, e_{i+j})\}$ and

$C^{(2)} = \bigcup_{i=1}^{n/2} C_i^{(2)}$, where $C_i^{(2)} = \{(v_i, e_{n+i-1}), (e_{n+i-1}, e_{n/2+i-1}), (e_{n/2+i-1}, v_{n/2+2-i})\}$.

Here, $\langle C^{(1)} \rangle \cong ((n-2)/2) P_4$, $\langle C^{(2)} \rangle \cong (n/2) P_4$. Hence, $B_2(K_{1,n})$ is P_4 -decomposable.

Case 2. n is odd.

The edge set of $((n^2 - n)/2) P_4$ is given by the set $C^{(3)}$, where

$C^{(3)} = \bigcup_{i=1}^n (\bigcup_{j=1}^{(n-1)/2} C_{j,i}^{(3)})$, where $C_{j,i}^{(3)} = \{(e_{i+2j-1}, v_i), (v_i, e_{i+2j}), (e_{i+2j}, e_{i+j})\}$,

$\langle C^{(3)} \rangle \cong n ((n-1)/2) P_4$. Therefore, $B_2(K_{1,n})$ is P_4 -decomposable.

In the following, P_4 -decomposition of Boolean function graph of corona of P_n , C_n and $K_{1,n}$ are found.

Theorem 3.4

(i) For $n \geq 3$ and $n \equiv 0 \pmod{3}$, the graph $B_2(P_n^+) - (n+1)K_2$ is P_4 -decomposable.

(ii) For $n \geq 4$ and $n \equiv 1 \pmod{3}$, the graph $B_2(P_n^+) - 2nK_2$ is P_4 -decomposable.

(iii) For $n \geq 5$ and $n \equiv 2 \pmod{3}$, the graph $B_2(P_n^+) - (3n-1)K_2$ is P_4 -decomposable.

Proof: Let $V(P_n^+) = \{v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n\}$, where v_1, v_2, \dots, v_n are the vertices of P_n in P_n^+ and u_1, u_2, \dots, u_n are the pendant vertices of P_n^+ and $e_i = (v_i, v_{i+1})$, $i = 1, 2, \dots, n-1$ and $f_i = (v_i, u_i)$, $i = 1, 2, \dots, n$ be the edges of P_n^+ . Then $v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n, e_1, e_2, \dots, e_{n-1}, f_1, f_2, \dots, f_n \in V(B_1(P_n^+))$. $L(P_n^+)$ is an induced subgraph of $B_2(P_n^+)$. $B_2(P_n^+)$ has $4n-1$ vertices and $4n^2 - 3n - 2$ edges.

Let $F_k = \{(v_i, e_j) \mid \text{for all } i, 1 \leq i \leq n, j \equiv (i+k) \pmod{(n-1)} \text{ and } e_0 = e_{n-1}\}$ and $F = \bigcup_{k=1}^{n-3} F_k$.

Let $H_k = \{(v_i, f_j), (u_i, f_j) \mid \text{for all } i, 1 \leq i \leq n, j \equiv (i+k) \pmod{n} \text{ and } f_0 = f_n\}$ and $H = \bigcup_{k=1}^{n-1} H_k$ and $|H| = 2n(n-1)$.

Let $J_k = \{(u_k, e_j) \mid \text{for all } i, 1 \leq i \leq n-1\}$, $J = \bigcup_{k=1}^n J_k$ and $|J| = n(n-1)$.

$E(B_2(P_n^+)) = L(P_n^+) \cup (F \cup H \cup J)$

In all the sets the suffix j in e_j are integers modulo $n-1$ and suffices k in u_k, v_k and f_k , are integers modulo n , $u_0 = u_n, v_0 = v_n, f_0 = f_n, e_0 = e_{n-1}$.

Case 1. $n \equiv 0 \pmod{3}$.

Then the edge set of $B_2(P_n^+)$ can be decomposed into $((4n^2 - 4n - 3)/3) P_4$ and $(n+1) K_2$.

The edges set $(n+1)K_2$ is given by the set $\{\bigcup_{i=1}^n \{(e_i, f_{i+1})\} \cup \{(v_1, e_{n-1}), (e_1, f_1)\}\}$. The edge

sets of $((4n^2 - 4n - 3)/3) P_4$'s are given by the edge sets $D^{(1)}, D^{(2)}$, and $D^{(3)}$, where

$D^{(1)} = \bigcup_{i=1}^n (\bigcup_{j=1}^{(n-1)/3} D_{j,i}^{(1)})$, where $D_{j,i}^{(1)} = \{(v_i, f_{i+j}), (f_{i+j}, u_i), (u_i, e_{i+j-1})\}$ and

$D^{(2)} = \bigcup_{i=1}^{n-1} (\bigcup_{j=1}^{(n-3)/3} D_{j,i}^{(2)})$, where $D_{j,i}^{(2)} = \{(e_{i+2j+1}, v_i), (v_i, e_{i+2j}), (e_{i+2j}, v_{i+3j+1})\}$ and

$D^{(3)} = \{(v_n, e_i), (e_i, e_{i+1}), (e_{i+1}, v_{i+(n+4)/3}) \mid i = 1, 2, \dots, n-2\}$. Here, $\langle D^{(1)} \rangle \cong n ((n-1)/3) P_4$, $\langle D^{(2)} \rangle \cong ((n-1)(n-3)/3) P_4$, $\langle D^{(3)} \rangle \cong (n-2) P_4$. Therefore, $B_2(P_n^+) - (n+1)K_2$ is P_4 -decomposable.

Case 2. $n \equiv 1 \pmod{3}$.

Then the edge set of $B_2(P_n^+)$ can be decomposed into $((4n^2 - 5n - 2)/3) P_4$ and $2nK_2$.

The edge set of $2nK_2$ is given by the set $\bigcup_{i=1}^{n-1} \{(v_i, e_{i+2}), (e_i, e_{i+1})\}, \{(v_1, e_{n-1}), (e_1, f_1)\}$. The

edge sets of $((4n^2 - 5n - 2) / 3) P_4$'s are given by the edge sets $D^{(1)}$ and $D^{(3)}$ as in Case 1, and the set $D^{(4)} = \bigcup_{i=1}^{n-1} (\bigcup_{j=1}^{(n-4)/3} D_{j,i}^{(4)})$, where $D_{j,i}^{(4)} = \{(e_{i+2j}, v_i), (v_i, e_{i+2j-1}), (e_{i+2j-1}, v_{i+3j+1})\}$. Here, $\langle D^{(4)} \rangle \cong (n-1) ((n-4)/3) P_4$. Hence, $B_2(P_n^+) - 2nK_2$ is P_4 -decomposable.

Case 3. $n \equiv 2 \pmod{3}$.

Then the edge set of $B_2(P_n^+)$ can be decomposed into $((4n^2 - 6n - 1) / 3)P_4$ and $(3n-1)K_2$. The edge set of $(3n-1)K_2$ is given by $\bigcup_{i=1}^{n-1} \{(e_i, f_{i+1}), (v_i, e_{i+1}), (v_i, e_{i+2}), (e_1, f_1), (v_1, e_{n-1})\}$. The edge sets of $((4n^2 - 6n - 1) / 3)P_4$ are given by the edge set $D^{(1)}$ and $D^{(3)}$ as in case1, and the set $D^{(5)} = \bigcup_{i=1}^{n-1} (\bigcup_{j=1}^{(n-5)/3} D_{j,i}^{(5)})$, where $D_{j,i}^{(5)} = \{(e_{i+2j+1}, v_i), (v_i, e_{i+2j+2}), (e_{i+2j+2}, v_{i+3j+3})\}$. Here, $\langle D^{(5)} \rangle \cong (n-1) ((n-5)/3)P_4$. Hence, $B_2(P_n^+) - (3n-1)K_2$ is P_4 -decomposable.

Theorem3.5

- (i) For $n \geq 3$ and $n \equiv 0 \pmod{3}$, the graph $B_2(C_n^+) - 2nK_2$ is P_4 -decomposable .
- (ii) For $n \geq 4$ and $n \equiv 1 \pmod{3}$, the graph $B_2(C_n^+) - 3nK_2$ is P_4 -decomposable .
- (iii) For $n \geq 5$ and $n \equiv 2 \pmod{3}$, the graph $B_2(C_n^+) - nK_2$ is P_4 -decomposable .

Proof: Let $V(C_n^+) = \{v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n\}$ where v_1, v_2, \dots, v_n are the vertices of C_n in C_n^+ and u_1, u_2, \dots, u_n are the pendant vertices of C_n^+ and $e_i = (v_i, v_{i+1}), i = 1, 2, \dots, n-1, e_n = (v_n, v_1)$ and $f_i = (v_i, u_i), i=1, 2, \dots, n$ be the edges of C_n^+ . Then $v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n, e_1, e_2, \dots, e_n, f_1, f_2, \dots, f_n \in V(B_2(C_n^+))$. $L(C_n^+)$ is an induced subgraph of $B_2(C_n^+)$. $B_2(C_n^+)$ has $2n$ vertices and $4n^2 - n$ edges.

Let $F_k = \{(v_i, e_j) / \text{for all } i, 1 \leq i \leq n, j \equiv (i+k) \pmod{(n-1)} \text{ and } e_0 = e_n\}$ and $F = \bigcup_{k=1}^{n-3} F_k$.

Let $H_k = \{(v_i, f_j) (u_i, f_j) / \text{for all } i, 1 \leq i \leq n, j \equiv (i+k) \pmod{(n)} \text{ and } f_0 = f_n\}$ and $H = \bigcup_{k=1}^{n-1} H_k$ and $|H| = 2n(n-1)$.

Let $J_k = \{(u_k, e_j) / \text{for all } i, 1 \leq i \leq n-1\}, J = \bigcup_{k=1}^n J_k$ and $|J| = n(n-1)$.

$$E(B_2(C_n^+)) = L(C_n^+) \cup (F \cup H \cup J)$$

In all the sets, the suffices i in v_i, u_i, e_i and f_i are integers modulo n and $v_0 = v_n, u_0 = u_n, e_0 = e_n$ and $f_0 = f_n$.

Case 1. $n \equiv 0 \pmod{3}$.

Then the edge set of $B_2(C_n^+)$ can be decomposed into $((4n^2 - 3n) / 3) P_4$ and $2nK_2$. The edges set $2nK_2$ is given by the set $\{\bigcup_{i=1}^n \{(e_i, f_i), (u_i, e_{n-1+i})\}\}$. The edge sets of

$((4n^2 - 3n) / 3)P_4$ are given by the edge sets $K^{(1)}, K^{(2)}$, and $K^{(3)}$, where

$$K^{(1)} = \bigcup_{i=1}^n (\bigcup_{j=1}^{(n-1)} K_{j,i}^{(1)}), \text{ where } K_{j,i}^{(1)} = \{(v_i, f_{i+j}), (f_{i+j}, u_i), (u_i, e_{i+j-1})\}.$$

$$K^{(2)} = \{(v_i, e_{i+1}), (e_{i+1}, e_{i+2}), (e_{i+2}, f_{i+3}), i = 1, 2, \dots, n\}$$

$$K^{(3)} = \bigcup_{i=1}^n (\bigcup_{j=1}^{(n-3)/3} K_{j,i}^{(3)}) \text{ where } K_{j,i}^{(3)} = \{(e_{i+2j}, v_i), (v_i, e_{i+2j+1}), (e_{i+2j+1}, v_{i+3j+2})\}.$$

Here, $\langle K^{(1)} \rangle \cong n(n-1)P_4, \langle K^{(2)} \rangle \cong nP_4$ and $\langle K^{(3)} \rangle \cong ((n(n-3)/3)P_4$.

Therefore, $B_2(C_n^+) - 2nK_2$ is P_4 -decomposable.

Case 2. $n \equiv 1 \pmod{3}$.

Then the edge set of $B_2(C_n^+)$ can be decomposed into $((4n^2 - 4n) / 3) P_4$'s and $3nK_2$. The edge set of $3nK_2$ is given by the sets $\bigcup_{i=1}^n (u_i, e_{n-1+i}), (e_i, e_{i+1}), (e_i, f_i)\}$. The edge sets of $((4n^2 - 4n) / 3) P_4$'s are given by the edge sets $K^{(1)}$ and $K^{(2)}$ as in Case 1, and the set $K^{(4)} = \bigcup_{i=1}^n (\bigcup_{j=1}^{(n-4)/3} K_{j,i}^{(4)})$, where $K_{j,i}^{(4)} = \{(e_{i+2j+1}, v_i), (v_i, e_{i+2j+2}), (e_{i+2j+2}, v_{i+3j+3})\}$.

Here, $\langle K^{(4)} \rangle \cong n((n-4)/3)P_4$. Hence, $B_2(C_n^+) - 3nK_2$ is P_4 -decomposable.

Case 3. $n \equiv 2(\pmod{3})$.

Then the edge set of $B_2(C_n^+)$ can be decomposed into $((4n^2 - 2n)/3)P_4$ and nK_2 . The edge set of nK_2 is given by $\{\cup_{i=1}^n (u_i, e_{n-1+i})\}$. The edge sets of $((4n^2 - 2n)/3)P_4$ are given by the edge set $K^{(1)}$ are in case 1, $K^{(5)}$ and $K^{(6)}$, where $K^{(5)} = \cup_{i=1}^n \cup_{j=1}^{(n-2)/3} K_{j,i}^{(5)}$ and $K_{j,i}^{(5)} = \{(e_{i+2j-1}, v_i), (v_i, e_{i+2j}), (e_{i+2j}, v_{i+3j+1})\}$, where $K^{(6)} = \{(e_i, f_{i+2}), (f_{i+2}, e_{i+1}), (e_{i+1}, e_{i+2}),$

$i = 1, 2, \dots, n\}$. Here, $\langle K^{(5)} \rangle \cong n((n-2)/3)P_4$ and $\langle K^{(6)} \rangle \cong nP_4$. Hence, $B_2(C_n^+) - nK_2$ is P_4 -decomposable.

Theorem 3.6

For $n \geq 3$, the graph $B_2(K_{1,n}^+) - 2nK_2$ is P_4 -decomposable.

Proof: Let $V(K_{1,n}^+) = \{v, v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n, u_{n+1}\}$, where v is the central vertex and $\langle \{v, v_1, v_2, \dots, v_n\} \rangle \cong K_{1,n}$ and $u_1, u_2, \dots, u_n, u_{n+1}$ are the pendant vertices of $K_{1,n}^+$ and $e_i = (v, v_i)$, $i = 1, 2, \dots, n$ and $f = (v, u)$, $f_i = (v_i, u_i)$, $i = 1, 2, \dots, n$, be the edges of $K_{1,n}^+$. Then $v, v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n, f_1, f_2, \dots, f_n, e_1, e_2, \dots, e_{n-1}, e_n \in V(B_2(K_{1,n}^+))$. $L(K_{1,n}^+)$ is an induced subgraph of $B_2(K_{1,n}^+)$. $B_2(K_{1,n}^+)$ has $4n+3$ vertices and $\frac{1}{2}(9n^2 + 7n)$ edges. In all the sets, the suffices i in v_i, u_i, e_i and f_i are integers modulo n and $v_0 = v_n, u_0 = u_n, e_0 = e_n$ and $f_0 = f_n$.

Let $F_i = \{(v_i, e_j), (v_i, f_j), (u_i, f_j) \mid 1 \leq j \leq n \text{ and } j \neq i\}$. $F = \cup_{i=1}^n F_i$, $|F| = 3n(n-1)$

Let $H_k = \{(u_k, e_j) \mid 1 \leq i \leq n\}$, $H = \cup_{k=1}^n H_k$ and $|H| = n^2$. Let $J = \{(v, f_j) \mid 1 \leq j \leq n\}$.

Let $L = \{(v_i, e), (u_i, e), (u, e_i), (u, f_i) \mid 1 \leq i \leq n\}$

$E(B_2(K_{1,n}^+)) = E(L(K_{1,n}^+)) \cup (F \cup H \cup J \cup L)$.

Let $M^{(1)} = \cup_{i=1}^n M_i^{(1)}$ and $M_i^{(1)} = \{(v, f_{i+1}), (f_{i+1}, u_i), (u_i, e_i)\}$ and $M^{(2)} = \cup_{i=1}^n (\cup_{j=1}^{(n-1)} M_{j,i}^{(2)})$ and $M_{j,i}^{(2)} = \{(v_i, f_{i+j-1}), (f_{i+j-1}, u_i), (u_i, e_{i+j})\}$ and $M^{(3)} = \cup_{i=1}^n M_i^{(3)}$ and $M_i^{(3)} = \{(v_i, f_i), (f_i, u_{n+1}), (u_{n+1}, e_i)\}$. Here, $\langle M^{(1)} \rangle \cong nP_4$, $\langle M^{(2)} \rangle \cong n(n-1)P_4$, $\langle M^{(3)} \rangle \cong nP_4$. The edge set of $2nK_2$ is given by the set $\{(e_i, f_i), (e_i, f_{i+1}), i = 1, 2, \dots, n\}$.

Case 1: n is odd

The edge set of $((3n^2 + n)/2)P_4$ are given by the set $M^{(1)} \cup M^{(2)} \cup M^{(3)} \cup M^{(4)}$, where $M^{(4)} = \cup_{i=1}^n (\cup_{j=1}^{(n-1)/2} M_{j,i}^{(4)})$ and $M_{j,i}^{(4)} = \{(e_{i+2j-1}, v_i), (v_i, e_{i+2j}), (e_{i+2j}, e_i)\}$.

Here, $\langle M^{(4)} \rangle \cong n(n-1)P_4$.

Case 1: n is odd

The edge set of $((3n^2 + n)/2)P_4$ are given by the set $M^{(1)} \cup M^{(2)} \cup M^{(3)} \cup M^{(4)}$, where $M^{(4)} = \cup_{i=1}^n (\cup_{j=1}^{(n-1)/2} M_{j,i}^{(4)})$, where $M_{j,i}^{(4)} = \{(e_{i+2j-1}, v_i), (v_i, e_{i+2j}), (e_{i+2j}, e_i)\}$.

Here, $\langle M^{(4)} \rangle \cong n(n-1)P_4$.

Case 2: n is even

The edge set of $((3n^2 + n)/2)P_4$ are given by the set $M^{(1)} \cup M^{(2)} \cup M^{(3)} \cup M^{(5)} \cup M^{(6)} \cup M^{(7)}$, where $M^{(5)} = \cup_{i=1}^n (\cup_{j=1}^{(n-4)/2} M_{j,i}^{(5)})$, where $M_{j,i}^{(5)} = \{(e_{i+2j-1}, v_i), (v_i, e_{i+2j}), (e_{i+2j}, e_{i+j-1})\}$ and $M^{(6)} = \cup_{i=1}^n M_i^{(6)}$, where $M_i^{(6)} = \{(e_{i+(n-3)}, v_i), (v_i, e_{i+(n-2)}), (e_{i+(n-2)}, e_{i+(n-1)})\}$ and $M^{(7)} = \cup_{i=1}^{n/2} M_i^{(7)}$, where $M_i^{(7)} = \{(v_i, e_{i+5}), (e_{i+5}, e_{i+2}), (e_{i+2}, v_{i+3})\}$. Here, $\langle M^{(5)} \rangle \cong (n(n-4)/2)P_4$, $\langle M^{(6)} \rangle \cong nP_4$, $\langle M^{(7)} \rangle \cong (n/2)P_4$. Therefore, $B_2(K_{1,n}^+) - 2nK_2$ is P_4 -decomposable.

4. Conclusion

In this paper, P_4 -Decomposition of Boolean Function Graph $B(\overline{K_p}, L(G), NINC)$ of path, cycle, stars and corona graphs are obtained.

5. References

- [1] P.Chithra Devi and J. Paulraj Joseph, P_4 - Decomposition of Total Graphs, Journal of Discrete Mathematical Sciences & Cryptography, Vol. 17(2014), No.5 & 6, pp.473-498.
- [2] Harary F, Graph Theory, Addison- Wesley Reading Mass., 1969.
- [3] K. Heinrich, J. Liu and M. Yu, P_4 - Decomposition of regular Graphs, Journal of Graph Theory, Vol.31(2), pp: 135 – 143, 1999.
- [4] T. N. Janakiraman, S. Muthammai, M. Bhanumathi, Domination Numbers on the Boolean Function Graph of a Graph, Mathematica Bohemica, 130(2005), No.2, 135-151.
- [5] T. N. Janakiraman, S. Muthammai, M. Bhanumathi, Domination Numbers on the Complement of the Boolean Function Graph of a Graph, Mathematica Bohemica, 130(2005), No.3, pp. 247-263.
- [6] T. N. Janakiraman, S. Muthammai, M. Bhanumathi, On the Boolean Function Graph of a Graph and on its Complement, Mathematica Bohemica, 130(2005), No.2, pp. 113-134.
- [7] S.Muthammai and S.Dhanalakshmi, Edge Domination in Boolean Function Graph $B(G, L(G), NINC)$ of a Graph, IJIRSET Journal, Vol. 4, Issue 12, December 2015, pp.12346 – 12350.
- [8] S.Muthammai and S.Dhanalakshmi, Edge Domination in Boolean Function Graph $B(G, L(G), NINC)$ of Corona of Some Standard Graphs, Global Journal of Pure and Applied Mathematics, Vol. 13, Issue 1, 2017, pp.152 – 155.
- [9] S.Muthammai and S.Dhanalakshmi, Connected and total edge Domination in Boolean Function Graph $B(G, L(G), NINC)$ of a graph, International Journal of Engineering, Science and Mathematics, Vol. 6, Issue 6, Oct 2017, ISSN: 2320 – 0294.
- [10] S.Muthammai and S.Dhanalakshmi, Domatic edge Domination in Boolean Function Graph $B(G, L(G), NINC)$ of a graph, Journal of Emerging Technologies and Innovative Research(JETIR), Vol .5, Issue 3, March 2018, ISSN: 2349 – 5162.
- [11] C.Sunil Kumar, On P_4 - Decomposition of Graphs, Taiwanese Journal of Mathematics, Vol.7, No.4, pp: 657-664, 2003.