

Studies on antibacterial and antimicrobial Activity of Macrocyclic Cr(III) and Fe(II) Complexes with Schiff base ligands.

Dr. Santosh Kumar Singh.

Govt. SV. Inter College, Bangra Jalalpur Saran, Bihar Pin - 841412

ABSTRACT

All over the world wide intersection which provide to comparative studies on macrocyclic metal Cr(III) and Fe(II) complexes with macrocyclic ligands. Consequently under the current communication, in coordination of our earlier interest. Very recently there has been a phenomenal growth in studies of Cr(III) and Fe(II) complexes. The complexes of the chelating ligands were synthesized using metal halides. The complexes of a new hydrogen bonded macrocyclic ligand derived from metal ion catalysed template synthesis of 1,2-Diamino-4,5-dinitrobenzene and diacetylmonoxime with Cr(III) and Fe(II) have been synthesized. Analyses with characterizations have been done on the basis of their elemental analyses, conductivity, magnetic moment in addition to spectral data and UV- visible. On the basis of electronic spectra and magnetic moment data suggest that the metal complexes have high spin octahedral paramagnetic extra hydrogen bonded more stable macrocyclic ring structure. Spectroscopic studies indicate that the coordination occurs with metal ions through two equatorial oxime-amine chelating ligands by neutral poly dentate donor azomethine nitrogen while other position of metal ions satisfied by two halides ions in facial and axial mode with Cr(III) and Fe (II) and formed three, five member and one six member extra hydrogen bonded more stable cyclic ring structure of complexes. The Cr(III) and Fe (II) complexes and ligands have been screened for antibacterial, antimicrobial activity and in the applied medicinal chemistry with other area of industries.

+2 Inter college, Jalal Pur Bazar, Jalal Pur, Saran Bihar¹.

Keyword- Stable ring structure, Conductivity, Medicinal Chemistry, Azomethine, Facial and axial mode.

INTRODUCTION


Coordination chemistry has received great attention in the field of inorganic chemistry during current time in all over the world. The reasons due to this interest are the fundamental importance in exchange interaction reaction with theory of bonding in new magnetic materials research and in understanding their important roles which play in large number of widely different processes of technological and biological importance⁵⁻⁹. Consequently in the resent communication in continuation of our earlier interest in the field. The metal complexes were prepared by refluxing hot ethyl alcohol solution of metal halide salts with the components of the ligand namely 1,2-Diamino-4,5-dinitrobenzene and diacetylmonoxime. The stoichiometry of the complexes have been found to be of the type $[Cr(H_2L)X_2]$ and $[Fe(H_2L)X_2]$. The chelating of ligand with Cr(III) and Fe(II) are expected to form complexes with different three dimensional cyclic ring geometry. It may enhance their biological activity as antibacterial and antimicrobial activity after chelatio. Which may find their importance in the applied medicinal chemistry with antibacterial and antimicrobial activity and other area of chemical industries.

Materials and method

The metal halide salts are used for synthetic and analytical work in experiment were of reagent grade solvent and purified by standard suitable method and dry before use in laboratory.

Ligand synthesis

The Schiff base ligands (H_2L) was synthesized by the condensation of two reactant molecules of diacetylmonoxime with 1,2-Diamino-4,5-dinitrobenzene under suitable condition in alcoholic medium. The reaction between the reactant molecules is occurs as in Figure - 1



The chemical reaction of synthesis of Ligands (H_2L)

Figure - 1

Preparation of complexes.

1. $[Cr(H_2L)Cl_2]Cl$

0.01 mole of alcoholic solution of chromic chloride with 0.02 mole of diacetylmonoxime and 0.1 mole of dinitroorthophenylenediamine were refluxed for 4 hour in a water bath. The green coloured of solid complexes are separated out. It was filtered, washed dried and analysed. The synthesis chemical reaction is given below.

2. $[Cr(H_2L)Br_2]Br$

0.01 mole of alcoholic solution of chromic bromide with 0.02 mole of diacetylmonoxime and 0.1 mole dinitroorthophenylenediamine were refluxed for 4 hour in water bath. The green coloured of solid complexes are separated out.

3. $[\text{Cr}(\text{H}_2\text{L})(\text{NO}_3)_2]\text{NO}_3$

0.01 mole of alcoholic solution of chromic nitrate with 0.02 mole of diacetylmonoxime and 0.1 mole of dinitroorthophenylenediamine were refluxed for 4 hours in water bath. The green coloured solid complexes are separated out.

4. $[\text{Cr}(\text{H}_2\text{L})\text{ClO}_4]\text{ClO}_4$

0.01 mole of alcoholic solution of perchlorate with 0.02 mole of diacetylmonoxime and 0.1 mole of dinitroorthophenylenediamine were refluxed for 4 hours in a water bath. The green coloured solid complexes are separated out.

5. $[\text{Fe}(\text{H}_2\text{L})\text{Cl}_2]$


The synthesis of Fe (II) complexes with the alcoholic solution of ferric chloride was kept in bright sunlight for six hours. When intense yellow colour of ferric chloride changed to faint green, almost colourless due to reduction of Fe (III) to Fe (II). The components of the ligand were added to it in inert atmosphere. On refluxing dark red colour of solid complexes are separated out. The reaction between them is shown below.

6. $[\text{Fe}(\text{H}_2\text{L})\text{Br}_2]$

The synthesis of Fe (II) complexes with the alcoholic solution of ferric bromide was kept in bright sunlight for six hours. When intense brown colour of ferric bromide changed to faint green, almost colourless due to reduction of Fe (III) to Fe (II). The components of the ligand were added to it in inert atmosphere. On refluxing dark red colour of solid complexes are separated out.

7. $[\text{Fe}(\text{H}_2\text{L})(\text{NO}_3)_2]$

The synthesis of Fe(II) complexes with alcoholic solution of ferric chloride was kept in bright sunlight for six hours. When intense brown colour of ferric nitrate changed to faint green almost coloured due to reduction of Fe (III) to Fe(II).The components of the ligand were added to it in inert atmosphere. On refluxing dark red colour of solid complexes are separated out. The reaction between them is shown below as Figure -2



M = Cr (III), Fe (II)

X = Cl⁻, Br⁻, NO₃⁻, ClO₄⁻

The chemical reaction of preparation of Complexes of [Mn(H₂L)X₂]

Figure- 2

Results and discussion

The complexes with intra-molecular hydrogen bonded macrocyclic ligand derived from metal ion catalysed template synthesis of 1,2-Diamino-4,5-dinitrobenzene and diacetylmonoxime with Cr(III) and Fe(II) have been synthesized. Characterizations have been done on the basis of their elemental analyses, conductivity, Magnetic moment with spectral data of I.R and UV-visible.

Elemental analysis

The physical and analytical data of the complexes are given in table – 1. This is in satisfactory agreement with the calculated values. The suggested molecular formula are supported by the subsequent spectral as well as magnetic moment, molar conductivity in 10⁻³ M solution of DMSO, the value of λ_m (Table –2) show that Cr (III) and Fe (II) complexes are non-electrolytic in nature.

Physical and analytical data for the complexes

Complexes	Colour	Yield %	Elemental analysis Calculation. (Found)			
			M%	N%	C%	H%
[Cr(H ₂ L)Cl ₂]Cl	Green	78	11.75(11.74)	18.27(18.26)	36.56(36.54)	3.48(3.47)
[Cr(H ₂ L)Br ₂]Br	Green	77	9.08(9.07)	14.13(14.12)	28.27(28.26)	2.71(2.68)
[Cr(H ₂ L)(NO ₃) ₂]NO ₃	Green	73	9.839.82)	22.90(22.89)	30.53(30.52)	2.90(2.89)
[Cr(H ₂ L)ClO ₄]ClO ₄	Green	79	8.29(8.28)	12.87(12.86)	25.74(25.74)	2.46(2.45)
[Fe(H ₂ L)cl ₂]	Red	96	13.12(13.10)	19.68(19.66)	39.32(39.32)	3.74(3.73)
[Fe(H ₂ L)Br ₂]	Red	91	10.86(10.83)	16.27(16.26)	32.55(32.54)	3.12(3.11)
[Fe(H ₂ L)(NO ₃) ₂]	Red	92	11.68(11.65)	23.32(23.31)	35.05(34.01)	3.32(3.31)

Table – 1

I.R Spectral data for the complexes

Complexes	V C=N (Oxime)	V C=N	V M=N	V M=O
[Cr(H ₂ L)Cl ₂]Cl	1491	1591	441	396
[Cr(H ₂ L)Br ₂]Br	1494	1596	461	401
[Cr(H ₂ L)(NO ₃) ₂]NO ₃	1501	1601	464	404
[Cr(H ₂ L)ClO ₄]ClO ₄	1504	1606	471	400
[Fe(H ₂ L)cl ₂]	1491	1609	480	406
[Fe(H ₂ L)Br ₂]	1503	1594	491	410
[Fe(H ₂ L)(NO ₃) ₂]	1502	1591	474	406

Table-2

Infrared spectra

Table-2. Lists the most useful infrared assignments for those bands most diagnostic of the mode of coordination of ligand. The most important band in the infrared spectra of 1,2-diamino-4,5-dinitrobenzene is due to ν_s and ν_{as} of -NH₂ groups. The I.R. spectra of diacetylmonoxime exhibits $\nu_{C=O}$ and $\nu_{C=N}$. In the spectra of the complexes ν_{NH_2} of 1,2-diamino-4,5-dinitrobenzene and $\nu_{C=O}$ of diacetylmonoxime is significantly absent indicating Schiff base condensation during macrocyclization. The $\nu_{C=N}$ oxime shows a downward shift indicating its involvement in coordination. In the far I.R. region two bands one in the region 440-610 cm⁻¹ and another in the region 395-415 cm⁻¹ may be assigned to ν_{M-N} and ν_{M-O} respectively. In the halogen complexes, a band in region 295-350 cm⁻¹ may be attributed to ν_{M-X} (X= Cl, Br)

Electronic spectra and magnetic moment studies

The UV-visible spectra of the complexes were recorded for their solutions in ethyl alcohol and DMSO solvents in the range (200-1000) nm. The ligand field spectra of Cr(III) complexes exhibit two bands in the region 624 and 455 nm which are assignable to ${}^4A_{2g} \rightarrow {}^4T_{2g}$ and ${}^4A_{2g} \rightarrow {}^4T_{1g}$ transition the ν_2/ν_1 ratio is 1.35 which is very close of value of 1.42 obtained for pure octahedral Cr(III) complexes. Electronic spectra and magnetic moment data of Cr(III) and Fe(II) complexes support paramagnetic high spin octahedral geometry. These Schiff base ligands and their metal complexes have been investigated as potential antibacterial and antimicrobial agents

Octahedral paramagnetic structure of the metal complexes

Figure- 3

REFERENCE

1. M. S. Refat. Investigation of coordination ability of Mn(II), Fe(III), Co(II), Ni(II), and Cu(II) with metronidazole, the antiprotozoal drug, in alkaline media: Synthesis and spectroscopic studies. *Russian Journal of General Chemistry*. 2017 vol. 87,873–879.
2. Kumar Raman, Synthesis and Spectral studies of macrocyclic Ni(II) complexes with multidentate ligands. *Research Journal of Chemistry and Environment*. 2019, 23 (9), 68-70.
3. S. Malik, S. Ghosh and L. Mitu, Complexes of some 3d-metals with a Schiff base derived from 5-acetamido-1,3,4-thiadiazole-2-sulphonamide and their biological activity. *Journal of the Serbian Chemical Society*. 2011, 76(10),1387-1394
4. Mahasin AliasHuda Kassum CarolinShakir. Synthesis, physical characterization and biological evaluation of Schiff base M(II) complexes. *Journal of the Association of Arab Universities for Basic and Applied Sciences*. 2014, 15, 28-34.
5. MA. Phaniband and SD. Dhumwad. Synthesis, characterization and biological studies of Co^{II}, Ni^{II}, Cu^{II} and Zn^{II} complexes of Schiff bases derived from 4-substituted carbostyrils[quinolin2(1H)-ones]*Trans. Met. Chem*. 2007, 32 (8) 1117-1125.
6. Manishaben jaiswal, "computer viruses: principles of exertion, occurrence and awareness ", *international journal of creative research thoughts (ijcrt)*, issn:2320-2882, volume.5, issue 4, pp.648-651, december 2017, <http://doi.one/10.1729/journal.23273> available at http://www.ijcrt.org/viewfull.php?p_id=ijcrt1133396
7. S. Chandra and SD. Sharma SD. *Trans. Met.* Spectral and antimicrobial studies on tetraaza macrocyclic complexes of Pd^{II}, Pt^{II}, Rh^{III} and Ir^{III} metal ions. 2011,15 (1) 49-54.
8. DHARMPAL SINGH and KRISHAN KUMAR , *Macrocyclic complexes: synthesis and characterization*. *J. Serb. Chem. Soc.* 2010, 75 (4) 475–482.
9. Milena Krstic, Branka Petkovic, Milos Milcic, Dusan Mistic Juan Francisco Santibanez. Synthesis, characterization and biological study of new dinuclear zinc(II) and nickel(II) octaaza macrocyclic complexes. *Macedonian Journal of Chemistry and Chemical Engineering* 2019, vol, 38, No. 1, 346-354.
10. S. Arulmurugan, HP. Kavitha and BR. Venkatraman. BIOLOGICAL ACTIVITIES OF SCHIFF BASE AND ITS COMPLEXES. *Rasayan. J. Chem.* 2010. 3 (3) 385 – 410.
11. A.D. Kulkarni. S.A. Patil. P.S. Badami. Electrochemical Properties of some Transition Metal Complexes: Synthesis, Characterization and In-vitro antimicrobial studies of Co(II), Ni(II), Cu(II), Mn(II) and Fe(III) Complexes . *Int. J. Electrochem. Sci.*, 2009, 4 .717 – 729.
12. AYSEN D. MULAZIMOGLU, IBRAHIM ENDER MULAZIMOGLU and BEDRETTIN MERCIMEK. Synthesis, Characterizations and Investigation of Electrochemical Behaviours of 4-[(2-Hydroxyphenylimino)methyl]benzene-1,3-diol. *Journal of Chemistry* 2009, 6(4), 965-974.
13. Bellam Sreenivasulu, Muthalagu Vetrichelvan. Fei Zhao Song Gao. Jagadese J. Vittal Copper(II) Complexes of Schiff-Base and Reduced Schiff-Base Ligands, Influence of Weakly Coordinating Sulfonate Groups on the Structure and Oxidation of 3,5-DTBC. *European journal of Inorganic Chemistry*. 2005. 22.4635-4645

14. Abd El-Motaleb M. Ramadan , Youssef L. Aly, Saied M. E. Khalil , Magdy Shebl and Ramy A. S. ElNaem. Synthesis and Characterization of Copper(II) Complexes Containing Sulfur/Nitrogen Donor Sets. Mimicking the Function of Phenoxazinone Synthase and Catechol Oxidase. *International Journal of Advanced Research*, 2015, Vol. 3, 7, 10-29.
15. MB. Fugu, NP. Ndahi, BB. Paul and AN. Mustapha. Synthesis, characterization, and antimicrobial studies of some vanillin Schiff base metal (II) complexes. *Journal of Chemical and Pharmaceutical Research*. 2013, 5 (4) 22- 28.
16. H.H. Darji, B. Shah & M.K. Jaiswal. "CONCEPTS OF DISTRIBUTED AND PARALLEL DATABASE", *International Journal of Computer Science and Information Technology & Security (IJCSITS)*, ISSN:2249-9555 vol. 2, no. 6, Dec. 2012, No;224, pg. no: 1150 available at https://www.academia.edu/42308519/CONCEPTS_OF_DISTRIBUTED_AND_PARALLEL_DATABASE
17. Tariq Mahmood Ansari Muhammad Najam-ul-Haq. Essential Trace Metal (Zinc, Manganese, Copper and Iron) Levels in Plants of Medicinal Importance. *Journal of Biological Sciences*, 2004, 4. 95-99.
18. E. Boyer, I. Bergevin, D. Malo, P. Gros, M. F. M. Cellier. Acquisition of Mn(II) in Addition to Fe(II) Is Required for Full Virulence of *Salmonella enterica* Serovar Typhimurium. 2002, 70.11.6032-6042.
19. G. Yan. Y. He, G. Li. Y. Xiong , P. Song and R. M. Wang et. al. Bovine Serum Albumin Metal Complexes for Mimic of SOD. *Journal of chemical sciences*. 2016, Vol. 128. No. 11, 1783-1788.
20. N. Raman, V. Muthuraj, S. Ravichandran, and A. Kulandaisamy, "Synthesis, characterisation and electrochemical behaviour of Cu(II), Co(II), Ni(II) and Zn(II) complexes derived from acetylacetone and *p*-anisidine and their antimicrobial activity, *Journal of Chemical Sciences*, 2003.vol. 115, no. 3, pp. 161–167.
21. Shamim AhmadKhan^aShahab A.A.Nami^bShahnawaz AhmadBhat^aAbdulKareem^a Synthesis, characterization and antimicrobial study of polymeric transition metal complexes of Mn(II), Co(II), Ni(II), Cu(II) and Zn(II). *Microbial Pathogenesis*.2017, Vol.110, 414-425
22. S.Dutt.P.Biswas.Activity of Mn Complexes Incorporating Schiff Bases. *International Journal of Applied Linguistics and Translation*. 2019, 5(2), 27-36.

