

# ELZAKI TRANSFORM AND ADOMIAN POLYNOMIAL FOR SOLVING THIRD ORDER NON-LINEAR KORTEWEG-DE VRIES EQUATIONS

<sup>1</sup>S.Sadiya, <sup>2</sup>A.Poomagal

<sup>1</sup>Assistant Professor, <sup>2</sup>Assistant Professor

<sup>1</sup>Department of Mathematics, Periyar University of Salem,

<sup>1</sup>Adhiyaman Arts and Science College for Women, Uthangarai, TamilNadu, India.

## Abstract:

The combination of Elzaki transform and Adomian was used to obtain the approximate analytical solutions of KdV equations in this research work. In total, three third order KdV equations were considered. Elzaki transform method was applied to obtain the approximate analytical solutions of all the aforementioned equations. Adomian polynomial was introduced as an essential tool to linearize all the associated nonlinear terms in the equations since Elzaki transform cannot handle nonlinear terms. All the problems considered yield exact solutions with few iterations.

**Keywords:** Nonlinear Differential equations, Elzaki transform, Adomian polynomial, Third order KdV equations, Finite difference method

## Introduction

The differential equations have played a central role in every aspect of applied mathematics for every long time and with the advent of the computer, their importance has increased further. Thus investigation and analysis of differential equations arising in applications led to many deep mathematical problems; therefore, there are so many different techniques in order to solve differential equations. In order to solve the differential equations, the integral transforms were extensively used and thus there are several works on the theory and applications of integral transforms such as the Laplace, Fourier, Mellin, Hankel and Sumudu, to name but a few. Recently, Tarig Elzaki introduced a new integral transform, named the Elzaki transform, and further applied it to the solution of ordinary and partial differential equations.

In this paper we derive the formulae for the Elzaki transform of partial derivatives and apply them in solving five types of initial value problems. Our purpose here is to show the applicability of this interesting new transform and its effecting in solving such problems.

The concept of Differential transform was first introduced by Zhou [7, 8], who solved linear and nonlinear initial value problems in electric circuit analysis. Later, DTM was applied to many problems in the literature by several authors [10-13].

Let us consider the KdV equation with given initial condition [6].

$$u_t + puu_x + qu_{xxx} = 0,$$

$$u(x,0) = f(x)$$

where  $p$  and  $q$  are real constants,  $u_t$  and  $u_x$  denote partial derivatives with respect to space  $x$  and time  $t$ , and the nonlinear term  $uu_x$  tends to localize the wave, whereas the wave was spread out by dispersion. The formulation of solitons that have a single humped wave was defined by delicate balance between  $uu_x$  and  $u_{xxx}$ . The displacement which describes how waves evolve under the competing but comparable effects of weak nonlinearity and weak dispersion is denoted by  $u(x,t)$ .

## Objectives

In order to achieve the main aim the following objectives will be carry out:

- 1 Introducing Adomian polynomial into Elzaki Transform method.
- 2 Applying the resulting scheme in 1 to solve nonlinear Kdv Equations which are IVP.
- 3 Finding the numerical solution of KdV equation using finite difference scheme.
- 4 To implement the discretize form or (system of equations) of KdV equation in Matlab.

### 1.1 Elzaki Transformation:

A new transform called Elzaki transform defined for function of exponential order we consider functions in the set A defined by:

$$A = \left( \begin{array}{l} f(t); \exists M, k_1, k_2 > 0, |f(t)| < M e^{\frac{|t|}{k_j}} \\ if t \in (-1)^j X[0, \infty) \end{array} \right)$$

In a set A, M is constant must be finite,  $k_1, k_2$  may be finite or infinite.

The Elzaki transform is defined by

$$E[f(t)] = T(v) = v \int_0^{\infty} f(t) e^{-\frac{t}{v}} dt, t > 0, k_1 \leq v \leq k_2.$$

In this transform the variable v is used to factor the variable t in the argument of function f. This transform has deeper connection with the Laplace transform.

### 1.2 Elzaki Transformations of partial Derivatives

Let T(u) be the Elzaki transform of f(t) i.e,  $E[f(t)] = T(u)$ , then:

- (i)  $E[f'(t)] = \frac{T(v)}{v} - v f(0)$
- (ii)  $E[f''(t)] = \frac{T(v)}{v^2} - f(0) - v f'(0)$
- (iii)  $E[f^n(t)] = \frac{T(v)}{v^n} - \sum_{k=0}^{n-1} v^{2-n+k} f^k(0)$ .

Note that  $E[f(t)] = T(u)$  means that T(u) is the Elzaki transform of f(t), and f(t) is the inverse Elzaki transform of T(u). i.e,

$$f(t) = E^{-1}[T(u)]$$

**Proof:** (i) From the equation  $E[f'(t)]$  is given by:

$$E[f(t)] = T(v) = v \int_0^{\infty} f(t) e^{-\frac{t}{v}} dt, t > 0, k_1 \leq v \leq k_2.$$

By integrating by part we have

$$\begin{aligned} \int_0^{\infty} f'(t) |t| e^{-\frac{t}{v}} dt &= e^{-\frac{t}{v}} \Big|_0^{\infty} + \frac{1}{v} \int_0^{\infty} f(t) e^{-\frac{t}{v}} dt. \\ &= -f(0) + \frac{1}{v} \int_0^{\infty} f(t) e^{-\frac{t}{v}} dt. \end{aligned}$$

Which yields to

$$\begin{aligned} E[f'(t)] &= u \left[ -f(0) + \frac{1}{v} \int_0^{\infty} f(t) e^{-\frac{t}{v}} dt \right] \\ &= -f(0) + \frac{1}{v} \int_0^{\infty} f(t) e^{-\frac{t}{v}} dt. \end{aligned}$$

Recall that

$$T(v) = v \int_0^{\infty} f(t) e^{-\frac{t}{v}} dt$$

Therefore

$$E[f'(t)] = \frac{T(v)}{v} - v f(0)$$

(ii) Using equations again, we have

By integrating by part we have

$$E[f^{(1)}(t)] = v \left[ -f'(0) + \frac{1}{v} \int_0^\infty f'(t) e^{-\frac{t}{v}} dt \right]$$

$$E[f^{(1)}(t)] = -vf'(0) + \frac{E(f'(t))}{v}$$

$$= \frac{E(f'(t))}{v} - vf'(0)$$

By considering the result from (i)

$$E[f^{(1)}(t)] = \frac{T(v)}{u^2} - f(0) - vf'(0)$$

(iii) For general form we are going to prove by induction

Given

$$E[f^{(n)}(t)] = \frac{T(v)}{v^n} - \sum_{k=0}^{n-1} v^{2-n+k} f^{(k)}(0), \text{ for all } n \geq 1$$

STEP 1: From equation when  $n=1$ , we have:

$$E[f^{(1)}(t)] = \frac{T(v)}{v} - vf(0)$$

$\frac{T(v)}{v} - vf(0)$  is clearly the same as  $\frac{T(v)}{v^1} - v^{2-1+0} f^{(0)}(0)$ , and the latter corresponds to equation when  $n=1$ .

This shows that equation holds for  $n=1$ .

STEP 2: Assume that equation holds for  $n=N$  i.e.,

$$E[f^{(N)}(t)] = \frac{T(v)}{v^N} - \sum_{k=0}^{N-1} v^{2-N+k} f^{(k)}(0)$$

We prove it equally hold for  $N+1$ , i.e.,

$$E[f^{(N+1)}(t)] = \frac{T(v)}{v^{N+1}} - \sum_{k=0}^N v^{2-(N+1)+k} f^{(k)}(0)$$

To begin with, note that from STEP 1 above

$$E[f^{(N+1)}(t)] = E[f^{(N)}(t)']$$

$$= \frac{E[f^{(N)}(t)]}{v} - v f^{(N)}(0)$$

From equation

$$E[f^{(N+1)}(t)] = \frac{T(v)}{v^{N+1}} - \sum_{k=0}^{N-1} v^{2-N+k-1} f^{(k)}(0) - v f^{(N)}(0)$$

$$= \frac{T(v)}{v^{N+1}} - \sum_{k=0}^N v^{2-(N+1)+k} f^{(k)}(0)$$

The last equally corresponds to equation when  $n=N+1$

## 2.1 Application of Elzaki transform and Adomian Polynomial for solving third order KdV equations

In this section we want to incorporate the Adomian polynomial into the Elzaki transform to solve both third-order KdV Equations, and to generalized the method of solving any kind of these equations by using both Elzaki transform and Adomian polynomial. Let's take a look on our previous research work before solving KdV equations by Elzaki transform.

### Application to third-order KdV Equations

Use the Elzaki transform method and Adomian polynomial to solve the following problems

### 2.2 Numerical Applications

#### Problem 1:

Consider the homogeneous KdV equation

$$u_t - 6uu_x + u_{xxx} = 0 \quad \rightarrow(1)$$

with initial conditions

$$u(x,0) = 10x$$

Let us rewrite Equation as:

$$u_t = [6uu_x - u_{xxx}]$$

Applying Elzaki transform to both side

$$E[u_t] = E[6uu_x - u_{xxx}] \quad \rightarrow(2)$$

By Elzaki transform property

$$E[u_t] = \frac{U(x,v)}{v} - v u(x,0)$$

$$\frac{U(x,v)}{v} - v u(x,0) = E[6uu_x - u_{xxx}] \quad \rightarrow(3)$$

Applying the given initial conditions on Equation and simplifying, we obtain;

$$U(x,v) = 10v^2x + vE[6uu_x - u_{xxx}] \quad \rightarrow(4)$$

Applying the inverse Elzaki transform,

$$u(x,t) = E^{-1}[10v^2x] - E^{-1}\{vE[6uu_x - u_{xxx}]\}$$

The resulting expression is

$$u(x,t) = 10x - E^{-1}\{vE[6uu_x - u_{xxx}]\} \rightarrow(5)$$

From  $u_0 = 10x$

The recursive relation is given as:

$$u_{n+1} = E^{-1}\left\{vE\left[6A_n - \frac{\partial^3 u_n}{\partial x^3}\right]\right\} \quad \rightarrow(6)$$

Note that  $A_n$  is the Adomian polynomial to decompose the nonlinear terms by using the relation:

$$A_n = \frac{1}{n!} \frac{d^n}{d\lambda^n} f\left[\sum_{i=0}^{\infty} \lambda^i u_i\right]_{\lambda=0} \quad \rightarrow(7)$$

Let the nonlinear term be represented by

$$f(u) = u \frac{\partial u}{\partial x} \quad \rightarrow(8)$$

By using Equation, we obtain;

$$A_0 = u_0 \left[ \frac{\partial u_0}{\partial x} \right]$$

$$A_1 = u_1 \left[ \frac{\partial u_0}{\partial x} \right] + u_0 \left[ \frac{\partial u_1}{\partial x} \right]$$

$$A_2 = u_2 \left[ \frac{\partial u_0}{\partial x} \right] + u_1 \left[ \frac{\partial u_1}{\partial x} \right] + u_0 \left[ \frac{\partial u_2}{\partial x} \right]$$

From Equation (6)

When  $n=0$ , we have,

$$u_1 = E^{-1} \left\{ vE \left[ 6A_0 + \frac{\partial^3 u_1}{\partial x^3} \right] \right\}$$

$$u_1 = E^{-1} \left\{ vE \left[ 6u_0 \left[ \frac{\partial u_0}{\partial x} \right] - \frac{\partial^3 u_n}{\partial x^3} \right] \right\}$$

since  $u_0 = 10x$  we have,

$$u_1 = E^{-1} \{ vE [600x] \} \quad \rightarrow(9)$$

$$u_1 = 600xt \quad \rightarrow(10)$$

When  $n = 1$ , we have;

$$u_2 = E^{-1} \left\{ vE \left[ 6A_1 - \frac{\partial^3 u_1}{\partial x^3} \right] \right\}$$

since  $u_1 = 600xt$  we have,

$$u_2 = E^{-1} \left\{ vE \left[ 6u_1 \left[ \frac{\partial u_0}{\partial x} \right] + 6u_0 \left[ \frac{\partial u_1}{\partial x} \right] - \frac{\partial^3 u_1}{\partial x^3} \right] \right\} \quad \rightarrow(11)$$

$$u_2 = 36000xt^2$$

When  $n = 2$ , we have;

$$u_3 = E^{-1} \left\{ vE \left[ 6A_2 - \frac{\partial^3 u_2}{\partial x^3} \right] \right\}$$

$$u_3 = E^{-1} \left\{ vE \left[ 6u_2 \left[ \frac{\partial u_0}{\partial x} - \frac{\partial^3 u_0}{\partial x^3} \right] + 6u_1 \left[ \frac{\partial u_1}{\partial x} - \frac{\partial^3 u_1}{\partial x^3} \right] + 6u_0 \left[ \frac{\partial u_2}{\partial x} - \frac{\partial^3 u_2}{\partial x^3} \right] + \frac{\partial^5 u_2}{\partial x^5} - \left( \frac{\partial u_2}{\partial x} \right)^2 \right] \right\} \quad \rightarrow(12)$$

$$u_3 = 2160000xt^3$$

The approximate series solution is,

$$u(x,t) = u_0 + u_1 + u_2 + \dots$$

$$u(x,t) = 10x + 600xt + 36000xt^2 + 2160000xt^3 + \dots$$

This can be written as

$$u(x,t) = 10x [1 + 60t + 3600t^2 + 216000t^3 + \dots]$$

By using Taylor's series, the closed form solution will be as follows

$$u(x,t) = 10x \left[ \frac{1}{1-60t} \right]$$

$$u(x,t) = \frac{10x}{1-60t}$$

## Conclusion

Generalizations of all existing Elzaki differentiation, integration and convolution theorems in the existing literature are demonstrated and so also generalizing all existing Elzaki shifting theorems. The Laplace – Elzaki duality (L E D ) will be used to invoke a complex inverse Elzaki transform.

Moreover, the problems considered show that the Elzaki transform method and Adomian polynomial is a very powerful integral transform method in solving both third order KdV equations. We have obtained the approximate analytical solutions of several third order and fifth order KdV equations using the combination of Elzaki transform method and Adomian polynomial which was meant for linearizing the nonlinear terms. Using this method make us to realise how potent this method is because all the problems considered yield exact solutions with few iterations. Solving any other nonlinear differential equations or ordinary is very easy by using this powerful method.

## REFERENCES

- [1] Yıldırım Keskin, Sema Servi, and Galip Oturanc, . Reduced differential transform method for solving klein gordon equations. In Proceedings of the World Congress on Engineering, volume 1, page 2011, 2011.
- [2] Yıldırım Keskin, İbrahim C, ađlar, and Ays,e Bet`ul Koc, . Numerical solution of sinegordon equation by reduced differential transform method. In Proceedings of the

World Congress on Engineering, volume 1, pages 6–8, 2011.

[3] Tuncay Aktosun and Cornelis van der Mee. Explicit solutions to the Korteweg–de Vries equation on the half line. *Inverse Problems*, 22(6):2165, 2006.

[4] G Adomian. Solving frontier problems of physics: The decomposition method. Kluwer, Boston, MA, 1994.

[5] Mariam BH Abushammala. Iterative Methods for the Numerical Solutions of Boundary Value Problems. PhD thesis, American University of Sharjah, 2014.

[6] S.Sadiya and T.Shanmuga Priya. Lie Symmetries and Reductions of (2+1) Dimensional Modified Equal Width Wave Equation With Damping Term, *JETIR* Volume 5, Issue 10, 2018.

[7] Norman J Zabusky and Martin D Kruskal. Interaction of "solitons" in a collisionless plasma and the recurrence of initial states. *Physical review letters*, 15(6):240, 1965.

[8] Peter S Riseborough. The Korteweg–de Vries equation: its place in the development of nonlinear physics. *Philosophical Magazine*, 91(6):997–1006, 2011.

[9] Chengri Jin and Mingzhu Liu. A new modification of Adomian decomposition method for solving a kind of evolution equation. *Applied Mathematics and Computation*, 169(2):953–962, 2005.

[10] Abdul-Majid Wazwaz and Alice Gorguis. Exact solutions for heat-like and wavelike equations with variable coefficients. *Applied Mathematics and Computation*, 149(1):15–29, 2004.

