

# EFFECT OF INDUSTRIAL WASTEWATER IRRIGATION ON THE SOIL CHARACTERISTICS OF NEARBY AREA IN BULDANA DISTRICT OF MAHARASHTRA.

G.D.Tambatkar

Department of Chemistry , Shri.D.M.Burungale Science & Arts College,Shegaon – 444 203

Dist:- Buldana ( M.S.) [ India ]

## ABSTRACT:-

Safe disposal of industrial wastewater has become a challenge for industrial managers and for scientists also. Rapidly increasing industrialization and urbanization have resulted into the generation of huge quantities of wastewater. Use of untreated industrial wastewater for crop irrigation is a common practice by the farmers. The immediate solution that seems to the problem of water shortage is the reuse of wastewater generated from industrial operations and domestic usage after the suitable treatment in agriculture and related activities.

The present research work deals with the study of effects of industrial wastewater irrigation on the soil characteristics of nearby area in Buldana District of Maharashtra. The data obtained in present study indicates that EC, Phosphate, TS, TDS, TSS, Ca, Mg etc., values in fertilizer wastewater whereas pH, TA, TH, Cl<sub>2</sub> values are maximum in sewage water of Khamgaon MIDC. At Sugar Industry sites, EC, Temp., phosphate, TDS, Ca, Mg, Cl<sub>2</sub> values in wastewater are maximum. Organic Carbon , Available N, K, P, Na, SO<sub>4</sub>, Cl<sub>2</sub>, Mg, Fe, Cu, Zn and Mn were comparatively higher in wastewater irrigated soils than the tube well irrigated soil.

**KEY WORDS:** - Wastewater, Soil Characteristics, Physico-Chemical Study, Buldana District.

**INTRODUCTION :** The requirement of water is increasing day by day due to tremendous increase in our demands on account of increasing domestic and industrial activities. Rapidly increasing industrialization and urbanization have resulted into the generation of higher quantities of wastewater. Use of untreated industrial wastewater for crop irrigation is a common practice by the farmers.

Irrigation with sewage and pulp paper mill effluent had enriched the soil mainly with respect to N, P, K and enhances the crop yields considerably<sup>1</sup>. Wastewater application for crop irrigation increases the zinc concentration in soil and sugarcane<sup>2</sup>. Increase in the organic carbon is beneficial for crops as it increases the update of potassium. Tannery and textile industrial wastewater contain appreciable amounts of plant nutrients such as N, P, K, Ca, Mg and S. These nutrients can be used for plant growth after proper treatment. Enrichment of soil organic carbon in soil due to addition of industrial wastewater is reported is reported by Chhonkar et.al<sup>3</sup>. The use of paper mill wastewater has favourably influenced the crop production, its continuous application for a number of years may results in enrichment of heavy metals in the top layers of soil<sup>4</sup>. Untreated industrial effluents contains higher amounts of Cd, Pb, Zn, Cu, Mn and Fe enhances the concentration of heavy metals in surface at irrigated soil<sup>5</sup>. Higher amounts of heavy metals like Cu, Zn, Fe and Mn is recorded in irrigated soil near in industrial complex<sup>6</sup>. Long term and indiscriminate application of raw sewage effluent result in the accumulation of heavy metals in surface and sub-surface soils<sup>7</sup>. Excessive accumulation of heavy metals such as Cd, Pb, Cr and Ni in the soil and the resultant phytotoxicity reported by Tsakou and Peralta<sup>8</sup>.

**MATERIALS AND METHODS :-** Wastewater and soil samples were collected from Khamgaon Industrial area of Buldana District, Maharashtra. At Khamgaon industrial site, soap unit is discharging wastewater into common drainage , which is being used for crop irrigation by local farmers. Wastewater samples were collected from the identified sites during the different seasons, i.e., premonsoon, monsoon and postmonsoon. Tube well water samples (control) were also collected for comparative studies. Samples were processed for the analysis of various physico-chemical parameters viz., pH, EC, DO, BOD, COD, TS, TSS, TDS, TH, Ca cations, Mg anions,

Cl<sub>2</sub> as per the standard methods of APHA-AWWA-WPCF ( 1985 ) . The heavy metal analysis samples were processed by acid digestion with 1:4 mixture of HClO<sub>4</sub> and HNO<sub>3</sub> and measured by atomic absorption Spectrophotometer.

Composite soil samples , irrigated with industrial wastewater were collected in three different seasons at depth of 0-15 and 15-30 cm. Samples were air dried and processed for the measurement of physico-chemical parameters viz., pH, EC by pH meter and conductivitymeter<sup>9</sup>, organic carbon<sup>10</sup>, available Phosphorus<sup>11</sup>, available K and Na<sup>12</sup>.

## RESULTS AND DISCUSSION

### Wastewater characteristics :

Wastewater and control samples analysis was characterised for pH, EC, organic carbon, total solid, hardness, alkalinity and heavy metals. Result of analysis of waste water and soil presented in table 1 and 2. of both sides and heavy metals of wastewater and soil presented in fig.2 and 3. pH value of khamgaon industry site was slightly alkaline in nature (7.1-8.2). EC values (1.3-4.0) were high in wastewater at Khamgaon site than control( 1.0-2.0). EC values low in monsoon season in all sites than premonsoon and postmonsoon season due to dilution by rainwater.

Dissolved oxygen(DO) values were low in industrial and sewage water than control, it could be due to high organic load in wastewater. BOD values were high in industrial and sewage wastewater (42-89) in comparison to control(0.98-1.8). COD values similarly high in wastewater at Khamgaon and Sugar mill sites than control. Average values of BOD and COD of wastewater were high in comparison to control the permissible limit (30 and 250 mg/l resp.) prescribed for the discharge of the effluent into inland surface water(EPR,1993)<sup>13</sup>. Temperature values marginally vary from site to site. Average values (176-374) of total alkalinity (TA) were high in wastewater in premonsoon and postmonsoon season than control(183-348). Phosphate values were comparatively high in industrial wastewater(1.8-5.6) than control. Total solid (TS) values including total dissolve and total suspended solid average values were high in waste water (1432-1903) than control (643-1730). In comparison to prescribed values of total suspended solid (TSS) for discharging waste water was (100mg /l). High concentration of suspended solid were found in monsoon season, it could be due to addition of loose soil particles from the adjacent land along with run off water. Total hardness values were high in waste water. Ca and Mg cation values were high in wastewater than control. Chloride values (251-596) low in control than wastewater (348-889). Average values of chloride were below maximum tolerance limit (600mg /l) for irrigation quality standard except Khamgaon industries and common drainage wastewater at sugar industries site. Heavy metals such as Cu, Zn, Fe

And Mn concentration were high in wastewater than control, whereas Fe values were high in control at Khamgaon Industries. Concentration of heavy metals were below the maximum recommended limit (100µg/l) prescribed for irrigation water<sup>14</sup>.

**Soil Characteristics :-** Results of the different parameters of soil analysis at both the sites are presented in Table No.3 and 4. Data indicates that pH values were high (7.4-9.0) in wastewater irrigated soil than control(7.1-8.5). The negative impact on soil pH can be attributed to wastewater irrigation, higher Na concentration increases pH of the soil. EC values were high in wastewater irrigated soil(0.4-3.0) than control values(0.25-0.6). High values of EC indicate that it could be due to availability of salt components K, Na, Mg, Cl and SO<sub>4</sub> in higher concentration in wastewater. EC values increases during the period of wastewater irrigation<sup>15</sup>. Values of EC were comparatively low in monsoon season than premonsoon and postmonsoon seasons due to dilution by rainwater. Moisture content values (0.32-6.3) were high in control soil in monsoon season than wastewater irrigated soil(0.37-6.0). Porosity values were low in wastewater irrigated soil (43.4-58.4) than control (49.8-59.3) it could be due to accumulation of salts in surface layer soil. Higher concentrations of Na, Ca, Mg salts and heavy metals affect the soil properties due to decrease in porosity after irrigation with wastewater. Organic Carbon values (0.37-0.94) were comparatively high in wastewater irrigated soil than control(0.11-0.79), due to high organic load in wastewater. Nutrient content N, P, K values high in wastewater irrigated soil in premonsson and postmonsson season than control. Increase in N, P, K concentration after wastewater irrigation without any significant negative impact has been reported by Vindhya et.al.2001<sup>16</sup>. Na and Mg values were high in wastewater irrigated soil in monsoon season at Sugar Industry

site. Observations indicate that the soil infested with moderate sodicity problem because of long term application of wastewater. Wastewater contains Na, Ca and Mg salts in higher concentration affects soil by direct accumulation of large quantities, as well as the interaction Na with exchange complex, which causes the deflocculation of soil particles<sup>17</sup>. Cl<sub>2</sub> and SO<sub>4</sub> values were low in control than wastewater irrigated soil. Carbonate content in soil layer was not detectable because of its possible conversion in CaCO<sub>3</sub>. However HCO<sub>3</sub> values (0.23-0.57) were high in wastewater irrigated soil in comparison to control (0.24-0.50). Analyzed values were high in premonsoon and postmonsoon than monsoon season. Concentration of Cu, Fe, Zn and Mn were comparatively high in wastewater irrigated soil than control. Heavy metal concentration increases in soil after wastewater irrigation<sup>18</sup>. Long term application sewage sludge containing higher Zn and Cr content exerts adverse effect on plant growth<sup>19</sup>. Analysis of heavy metal indicates that the long term and indiscriminate application of wastewater may cause accumulation of heavy metals in surface soil and it will be harmful for plants and animals. Therefore, extensive research is required for the long term assessment of heavy metals accumulation in soil, crops and consumers due to wastewater irrigation.

#### ACKNOWLEDGEMENT

The authors are thankful to HOD, Department of Zoology, Sant Gadge Baba Amravati University, Amravati for providing necessary laboratory facilities for research work.

#### REFERENCES :

- APHA, AWWA AND WPCF, 1985 Standard Methods for the Examination of Water and Wastewater. 16<sup>th</sup> Ed. APHA Inc., New York pp. 1268.
1. Nan J. and Chung G., 2001. Accumulation of Cd and Pb in spring wheat (*Triticum aestivum* L.) grown in calcareous soil irrigated with wastewater. *Bull. Environ. Contam. Toxicol.* 66: 748-754.
  2. Terras C., Navarro M., Martin-lagos, F. Gimenez R., Olalla M., Lopez H. and Lopez M., 2002. Influence of pH and industrial activity on total zinc concentrations in agricultural soils, sewage sludge and beach sands: relation with plant availability. *Bull. Environ. Contam. Toxicol.* 68: 224-229.
  3. Chhonkar P.K., Datta S.P., Joshi H.C. and Pathak H., 2000. Impact of industrial effluents on soil health and agriculture, part II – Tannery and Textile industrial effluents. *J. of Scientific and Industrial Research*, 59: 446-454.
  4. Gupta S.K., Mitra A. and Adhikari S., 1990. Proc. Natn. Symp. on Protec. of Environ. of city Wat. Fronts. Central Water Commission, New Delhi, IV-7.
  5. Xiog X., Stagnitri F., Peterson J., Allinson G. and Turoczy N., 2001. Heavy metal contamination of pasture soils by irrigated municipal sewage. *Bull. Environ. Contam. Toxicol.* 67: 535-540.
  6. Barman S.C., Kisku G.C., Salve P.R., Mishra D., Sahu R.K., Ramteke P.W. and Bhargawa S.K., 2001. Assessment of industrial effluent and its impact on soil and plants. *J. Environ Biol.* 22, 4 : 251-256.
  7. Datta S.P., Biswas D.R., Sharon N., Ghose S.K. and Rattan R.K., 2000. Effect of long term application of sewage effluents on organic carbon, available phosphorus, potassium and heavy metal status and content of heavy metals in crops grown thereon. *J. Indian Soc. Sci.* 48, 4: 836-839.
  8. Tsakou A., Roulia M. and Christodoulakis N.S., 2001. Growth of cotton plant (*Gossypium hirsutum*) as affected by water and sludge from sewage treatment plant, plant pathology and plant development. *Bull. Environ. Contam. Toxicol.* 66: 735-742.
  9. Piper C.S., 1950. Soil and Plant Analysis, Interscience publishers, Univ. Adelaide, Australia.
  10. Walkely A.C. and Black T.A., 1934. Estimation of soil organic carbon by the chromic acid filtration method. *Soil Sci.*, 47: 27-38.
  11. Olsen S.R., Cole C.L., Watnabe F.C. and Dean D.D., 1964. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *USDA. Circs.* 939.
  12. Jackson C.S. 1973. Soil Chemical Analysis. Prentice Hall of India, New Delhi.
  13. EPR 1993. In the Gazette of India: Extraordinary (Part – II Sec. 3), Min. Environ. Forest Govt. of India.
  14. Pratt P.F., 1972. Quality criteria for trace elements in irrigation waters. Publications of Univ. California Expt. Stn. Riverside, California, USA.

15. Singh G., Bala N., Rathod T.R. and Singh B., 2001. Effect of Textile effluent on tree plantation and soil chemistry. *J. Environ.Biol.*, 22,1:59-66.
16. Vindhya V.,Ramesh Babu,Kale C.K., Kothanderman V.,Manivell U., Jayabaiou R. and Subramanian Y.V., 2001. Irrigation water soil quality with reference to operation of a small scale chemical industry, *I.J.E.P.*,21,5:409-415.
17. Mitra A. and Gupta S.K. , 1999. Effect of sewage water irrigation on essential plant nutrient and pollutant element status in a vegetable growing area around Calcutta. *J. of Indian Soc.Soil.*, 47,1:99-105.
18. Tyagi P. and Joshi N., 2001. Acomparative study of some physico-chemical parameters of soil irrigated with recycled and tube well water.*I.J.E.P.*, 21,6:5125-528.
19. Wong J.W.C., Lai K.M.,Su.D.S. and Fang M.,2001. Availability of heavy metals for *Brasica Chinensis* Grown in an acidic loamy soil amended with a domestic and an industrial sewage water sludge water.*Water,Air and Soil Pollution*,128,3-4:339-353.

