

PHARMACOLOGICAL POTENTIAL OF AVERRHOA CARAMBOLA: A REVIEW

¹Kumari Sonu *, ²Dr. Dev Prakash, ³Balwan Singh
Student of M. Pharm Pharmacology, Principal cum Director, HOD
¹Pharmacy
HIMT College of Pharmacy, Greater Noida, U.P., India

Abstract

Averrhoa carambola L. (Oxalidaceae), normally known as star fruit bears a great consequence in traditional medicine. Traditionally, *A. carambola* was used in ailments such as arthralgia, chronic headache, boils and pyodermas, colds, cough, epistaxis, spermatorrhea, fever, food poisoning, gastroenteritis, malaria, malarial splenomegaly, oliguria, postpartum edema, sore throat, subcalorism and traumatic injury. Pharmacological investigations on *A. carambola* have established anti-inflammatory, antimicrobial, antifungal, antitumor and anti-ulcer activities. In addition, the plant possesses hypocholesterolemic, hypoglycemic, hypotensive, nephrotoxic, neurotoxic, negative inotropic and chronotropic effects. Phytochemical investigations have exposed the occurrence of saponins, tannins, alkaloids and flavonoids. This review is an attempt to modernize the pharmacological activities and clinical studies on *A. carambola*.

Key words: Averrhoa carambola, Oxalidaceae, pharmacological activities, chemical constituents

1. INTRODUCTION

Star fruit, *Averrhoa carambola* (Oxalidaceae) is originate from America, Brazil, and Australia, South-East Asia including Malaysia, Southern China, Taiwan and India. *A. carambola* tree is frequently 3 to 5 m in height and can reach a greatest height of 10 m, with a finely fissured light brown bark and 15 to 20 cm long leaves. It bears big indehiscent yellowish-green berry fruit of 5 to 8 cm long with a characteristic form similar to a five pointed star, and every cell of the fruit contains five arillate seeds [1-5]. In traditional medicine, the fruit was used for treating ailments such as cough, food poisoning, sore throat and malarial splenomegaly; the root for treating arthralgia, chronic headache, epistaxis and spermatorrhea; the leaves for treating boils, colds, gastroenteritis, oliguria, postpartum edema, pyodermas and traumatic injury, whilst the flowers for treating fever, malaria and subcalorism [6-9]. There are preceding reviews on *A. carambola* describing mainly the botanical and cultivation aspects, traditional uses and some pharmacological activities [10-12]. In this review, we recapitulate and bring up to date the phytochemical, pharmacological and clinical investigations on *A. carambola*.

Classification [13]

Scientific Name: *Averrhoa carambola*

Kingdom: Plantae – Plants

Subkingdom: Tracheobionta -Vascular plants

Superdivision: Spermatophyta

Division: Magnoliophyta – Flowering plants

Class: Magnoliopsida – Dicotyledons

Subclass: Rosidae

Order: Geraniales

Family: Oxalidaceae – Wood-Sorrel family

Genus: Averrhoa Adans. – Averrhoa

Species: *Averrhoa carambola* L. – carambola

Vernacular names [14, 15]

- Sanskrit : Karmaranga
- English : Starfruit, Chinese gooseberry
- Hindi : Kamrakh, Karmal
- Bengali : Kamranga
- Assamese : Kordoi/ rohdoi
- Gujarati : Kamrakh
- Marathi : Karambal
- Telugu : Ambanamkaya
- Tamil : Thambaratham/Tamarattai
- Malayalam : Caturappuli
- Sinhala : Kamaranga
- Filipino : Balimbing, saranate
- Indonesian : Belimbing
- Malay : Belimbing

Origin and distribution

Star fruit is supposed to have originated in Ceylon and the Moluccas, but it has been cultivated in Southeast Asia and Malaysia for hundreds of years. [16] The perennial herb is usually grown in Malaysia, Taiwan, Thailand, Israel, Florida, Brazil, Philippines, China, Australia, Indonesia, in the warmer parts of India, Bangladesh and other areas of the world with the same climate [17].

Description

Botanical Description/ macroscopy- *Averrhoa carambola* is a small, attractive, multi stemmed, sluggish growing evergreen tree with a short trunk or a shrub, 5-7 m of height or rarely, 10 m high, spreading 20-25 ft in diameter. It has a bushy shape with many branches producing a broad, rounded crown. At the base, the trunk reaches a diameter of 15 cm [18, 19].

Chemical constituents

Preliminary phytochemical analysis of carambola fruit showed the attendance of saponins, alkaloids, flavonoids and tannins [21]. The fruit was too established to hold proanthocyanidins, epicatechin, gallic acid in gallotannin form and L-ascorbic acid [22]. It was reported that the chief sterols present in the fruits of carambola are β -sitosterol, campesterol, lupeol and isofucosterol; it also contained the four major plant fatty acids – palmitic, oleic, linoleic and linolenic acid [23]. Carambola have identified o-glycosyl flavonoid constituent such as quercetin-3-o- β -d glycoside and rutin. Edible segment of the fruit is a excellent source of dropping and non-reducing sugars, minerals, volatile favours, tannins, dietary fibers, pectin, cellulose, hemicelluloses, iron, calcium, phosphorous and carotenoid compositions [24]. Other components identified are cyaniding-3-o- β -dglucoside, cyaniding-3-5-o- β -d-diglycoside, β -amirin [25] and C glycoside flavones, such as apigenin-6-C- β -L-fucopyranoside and apigenin-6-C- β -1-fucopyranoside. this latter compound are also known a carambola flavone [26]. Occurrence of p-Anisaldehyde, 5-hydroxymethyl-2-furfur-al, Gallic acid and dihydroabscissic alcohol in the stem bark of *A. carambola* [27].

Furthermore, fifteen compounds, which included six chiral lignans and nine phenolic glycosides, were separated from the butanol fraction of *Averrhoa carambola* L. root and identified. All of the compounds, namely 3,4,5-trimethoxy-phenol-1-O- β -D-glucopyranoside (**1**), benzyl-1-O- β -D-glucopyranoside (**2**), (+)-5'-methoxyisolariciresinol 3 α -O- β -D-glucopyranoside(**3**), (+)-isolariciresinol 3 α -O- β -D-glucopyranoside (**4**), koaburaside (**5**), (+)-lyoniresinol 3 α -O- β -D-glucopyranoside (**6**), (-)-lyoniresinol 3 α -O- β -D-glucopyranoside (**7**),(-)-5'-methoxyisolariciresinol 3 α -O- β -D-glucopyranoside (**8**), (-)-isolariciresinol 3 α -O- β -D-glucopyranoside (**9**), 3,5-dimethoxy-4-hydroxyphenyl 1-O- β -apiofuranosyl (1"→6')-O- β -D-glucopyranoside (**10**), 3,4,5-trimethoxyphenyl 1-O- β -apiofuranosyl (1"→6')- β -glucopyranoside (**11**), methoxy-hydroquinone-4- β -D-glucopyranoside (**12**), (2S)-2-O- β -D-glucopyranosyl-2-hydroxy-phenylacetic acid (**13**), 3-hydroxy-4-methoxyphenol-1-O- β -D-apiofuranosyl-(1"→6')-O- β -D-glucopyranoside (**14**) and 4-hydroxy-3-methoxyphenol 1-O- β -Dapiofuranosyl-(1"→6')-O- β -D-glucopyranoside (**15**) were isolated from this plant for the first time [28].

Nutritional value of star fruit

Averrhoa carambola is rich with all vital nutrients. It is a good source of natural antioxidants like L-ascorbic acid, (-) epicatechin and gallic acid in gallotannin forms [22]. Consuming 100 g of this fruit can provide, 35.7 g calories, 0.38 g proteins, 9.38 g carbohydrates, 0.80 g-0.90 g dietary fibre, 0.8 g fat, 4.4-6.0 mg calcium, 0.32-1.65 mg iron, 15.5-21.0 mg phosphorus, 2.35 mg potassium, 0.003-0.552 mg of carotene, 4.37 mg tartaric acid, 9.6 mg oxalic acid, 2.2 mg α -ketoglutaric acid, 1.32 mg citric acid. Moreover, various amino acids like 0.03-0.038 mg of thiamine, 0.019-0.03 mg of riboflavin, 0.294-0.38 mg of niacin, 3 mg of tryptophan, 2 mg of methionine and 26 mg of lysine are also present in 100 g of the fruit [29].

Pharmacological properties

Antioxidant capacity

Luximon-Ramma *et al.*, (2003) examined the antioxidant capacity, total phenolics, proanthocyanidins, flavonoids and vitamin C contents of 17 commonly consumed exotic fruits from Mauritius including *A. carambola*. The antioxidant activities of *A. carambola* were found to be ranged from 11 to 17 μ mol Trolox equivalent antioxidant capacity (TEAC)/g fresh weight, and 9 to 22 μ mol ferric reducing antioxidant power (FRAP)/g fresh weight, for the acid and sweet fruits correspondingly. The total phenolics content was 1429 and 2099 μ g/g fresh weight; proanthocyanidins 896 and 1321 μ g/g fresh weight; total flavonoids 103 and 148 μ g/g fresh weight and vitamin C content was 190 and 144 μ g/g fresh weight, for the acid and sweet fruits correspondingly. There were strong correlations among antioxidant activity (assessed by both TEAC and FRAP) and total phenolics and proanthocyanidins contents. Flavonoids seemed to add less to the antioxidant prospective of the fruit, whilst very poor association was observed between ascorbate content and antioxidant activity. The antioxidant power of *A. carambola* was establish to be amongst the highest in the study group and the study accomplished *A. carambola* to be a considerable source of phenolic antioxidants, thus it may demonstrate potent health benefits. Shui and Leong, (2006) showed that the residue of *A. carambola* fruit, which is normally redundant during juice drink processing, contains much higher antioxidant activity than the extracted juice using different methods for assessing antioxidant activity. Under optimized extraction conditions, the remains accounted for around 70 % of total antioxidant activity and total polyphenolic contents, nevertheless it only contributed 15 % of the weight of the entire fruit. Freeze-dried residue-powder, which accounted for approximately 5 % of total weight, had total polyphenolic content of 33.2 ± 3.6 mg gallic acid equivalent (GAE)/g sample, total antioxidant activity of 3490 ± 310 and 3412 ± 290 mg L-ascorbic acid corresponding antioxidant capacity or 5270 ± 468 and 5152 ± 706 mg trolox corresponding antioxidant capacity per 100 g sample obtained by 2,2'-azino-bis-(3-ethylbenzthiazoline-6-sulfonic acid) free radical and 1,1-diphenyl-2-picryl-hydrazyl scavenging assays, respectively. It was also found to have 510.3 ± 68.1 mol ferric reducing/antioxidant power per gram sample. The residue-extract also shows strong antioxidant activity in delaying oxidative rancidity of soya bean oil at 110 °C. Antioxidant activity and polyphenolic contour of residue extracts were compared with extracts of standardized pyconogenol. High performance liquid chromatography joined with mass spectrometry revealed that major proanthocyanidins in *A. carambola* fruit were different from their isomers in the standardised extracts of pyconogenol. The high content of phenolics and strong antioxidant activity of residue extract indicate that the residue powder may yield health benefits when used in functional food products and that residue extract should also be regarded as potential source of nutrients in future. Ali *et al.*, (2011) screened 20 different kinds of fruits including *A. carambola* from Aizawal Market of Mizoram India for their antioxidant potential using DPPH, FRAP and total phenolics content assay. *A. carambola* showed moderate antioxidant activity among the tested fruits, the activity was found to be 81.03 ± 1.97 g of Trolox equivalent/100 g of fruit (DPPH assay), 78.770 ± 0.33 g of Trolox equivalent/100 g of fruit (FRAP assay) and total phenolics contents were found to be 54.45 ± 0.43 g of gallic acid equivalent/100 g of fruit. Moresco *et al.*, (2012) reported isolation of β - sitosterol from the *n*-hexane fraction;

apigenin-6-C- β -L-fucopyranoside and apigenin-6-C-(2"-O- α -L-rhamnopyranosyl)- β -L-fucopyranoside from the ethyl acetate fraction, and a new compound apigenin-6-C-(2"-O- α -L-rhamnopyranosyl)- β -D-glucopyranoside from the *n*-butanol fraction. The antioxidant potential of the fractions was examined using the DPPH radical scavenging assay and reducing power of iron (III) to iron (II) ions. The ethyl acetate and *n*-butanol fractions

showed the highest antioxidant potential to scavenge DPPH, with the IC₅₀ values of 90.92 and 124.48 μ g/mL, correspondingly. In the assay of reducing power, these fractions presented 135.64 and 125.12 mg/g of ascorbic acid equivalents, correspondingly. The antioxidant activity exhibited a momentous correlation with the phenolic content ($r^2 = 0.997$), but a poor correlation with the flavonoids content ($r^2 = 0.424$). The *n*-hexane fraction was the only fraction to show toxicity on *A. salina* with LC₅₀ of 800.2 μ g/mL.

Anti-inflammatory activity

Cabrini *et al.*, (2011) examined the ethanol extract of *A. carambola* leaves, its hexane, ethyl acetate and butanol fractions and two isolated flavonoids, apigenin-6-C- β -L-fucopyranoside and apigenin-6-C-(2"-O- α -L-rhamnopyranosyl)- β -L-fucopyranoside for anti-inflammatory activity. The ethanol extract reduced edema in a dose-dependent manner, resulting in a maximum inhibition of $73 \pm 3\%$ and an ID₅₀ value of 0.05 (range: 0.02–0.13) mg/ear. Myeloperoxidase activity was also inhibited by the ethanol extract, resulting in a maximum inhibition of $60 \pm 6\%$ (0.6 mg/ear). The tested fractions introverted edema formation and myeloperoxidase activity, the ethyl acetate fraction was the majority successful fraction, resulting in inhibition levels of 75 ± 5 and $54 \pm 8\%$ for edema formation and myeloperoxidase activity, correspondingly. On the other hand, the isolated compounds did not show any considerable anti-inflammatory activity.

Acetylcholinesterase inhibitory activity

Teh *et al.*, (2010) investigated the effects of *A. carambola* juice kept at dissimilar storage conditions on the activity of acetylcholinesterase in a variety of organs of Sprague Dawley (SD) rats. Three groups (5 animals per group) of rats were administered with distilled water, and dissimilar star fruit preparations, freshly prepared star fruit juice and after 3 hours storage, correspondingly. The results showed a considerable reduce in the hepatic acetylcholinesterase activity in rats treated with star fruit juice maximum activity was observed for the freshly prepared juice.

Antimicrobial and antifungal activity

Mia *et al.*, (2007) isolated two compounds *p* anisaldehyde and β -sitosterol from the bark of *A. carambola*. The petroleum ether, carbon tetrachloride and chloroform fractions of the methanol extract of *A. carambola* when subjected to antimicrobial showing at 400 μ g/disc established mild inhibition on microbial growth. The average zone of inhibition produced by the petroleum ether, carbon tetrachloride and chloroform soluble fractions were 8-12 mm, 8-12 mm, and 8-15 mm, respectively. The petroleum ether extract moderately inhibited the growth of *E. coli* and *S. dysenteriae* having the zone of inhibition of 12 mm each. On the other hand, the chloroform soluble fraction strongly inhibited the growth of *E. coli* with zone of inhibition 15 mm. In the case of fungi, mild inhibitory activity was exhibited by all extractives. The LC₅₀ values obtained from cytotoxicity evaluation were 0.32, 0.70, 0.06 and 3.14 μ g/mL for standard vincristine sulfate, petroleum ether, carbon tetrachloride and chloroform soluble fractions, respectively.

Antitumor activity

Li *et al.*, (2012) studied the biotransformation of dihydro-epi-deoxyarteannuin B by using suspension cultured cells of *A. carambola*. One novel sesquiterpene, 7 α -hydroxy-dihydro-epideoxyarteannuin B, and one known sesquiterpene, 3- α -hydroxydihydro-epideoxyarteannuin B, were obtained upon the addition of dihydro-epi-deoxyarteannuin B. The study concluded that, cultured cells of *A. carambola* have the capacity to hydroxylate sesquiterpene compounds in a regio- and stereo selective manner. The inhibitory effects of 7 α -hydroxy-dihydroepideoxy arteannuin B and 3- α -hydroxy-dihydroepideoxyarteannuin B on proliferation of K562 and HeLa cell lines were (59.29 ± 0.99 , 84.04 ± 0.27 μ mol/mL) and (40.63 ± 1.45 , 41.54 ± 0.82 μ mol/mL), respectively.

Anti-ulcer activity

Goncalves *et al.*, (2006) examined the water-alcohol extract of leaves of *A. carambola* for its antiulcerogenic power. The ethanolic extract at doses of 800 and 1200 mg/kg *p.o.*, only showed important anti-ulcer activity in the acidified-ethanol-induced ulcer model in rats. However, the extract did not show any activity in the indomethacin and acute stress ulcerogenic models. Thus, the study concluded ethanolic extract of *A. carambola* as having little antiulcer activity.

Negative inotropic and chronotropic effect

Vasconcelos *et al.*, (2005) examined the atrial isometric force in inspired left atria and determined the chronotropic changes in impulsively thrashing right atria. *A. carambola* leaves extracts (1.5 mg/mL) abolished the contractile force in a concentration dependent manner in cardiac contractility in the guinea-pig atria. Amongst the methanolic, ethanolic, aqueous, and acetic acid extracts, the aqueous extract was establish to be the most potent (EC₅₀ = 520 ± 94 μ g/mL). The aqueous extract abolished the positive. Bowditch staircase phenomenon and reduced the inotropic response to CaCl₂, events that are dependent on the cellular Ca₂₊ inward current. In impulsively beating atria, the aqueous extract promoted a negative chronotropic effect that was antagonized by 0.1 μ M isoproterenol bitartrate. The EC₅₀ of the aqueous extract increased from 133 ± 58 to 650 ± 100 μ g/mL in the presence of this agonist. Subsequently, Vasconcelos *et al.*, (2008) investigated the effects of the aqueous extract of *A. carambola* leaves on the cellular calcium influx by examining the left atrium of guinea pig and the pituitary GH3 cells. In the atrium, the aqueous extract (1500 μ g/mL) shifted to the right the concentration effect

Curve of the positive inotropic effect produced by (\pm) BAY K 8644, an L-type calcium channel agonist. The aqueous extract increased EC₅₀ (concentration required to promote 50% of the maximum effect) of the inotropic effect of BAY K 8644 from 7.8 ± 0.38 to 115.1 ± 0.44 nM. In GH3 cells treated with 500 μ g/mL of aqueous extract, the L-type calcium inward current declined by

30%. The extract did not change the voltage correspondent to the peak current. It was suggested that, at least in part, the negative inotropic effect of aqueous extract on the guinea pig atrium is due to a reduction of the L-type calcium current.

Electrophysiological effects

Vasconcelos *et al.*, (2006) examined the effects of the aqueous extract of *A. carambola* leaves on guinea pig heart. In this study, the aqueous extract induced many kinds of atrioventricular blocks (1st, 2nd, and 3rd degrees); depressed the cardiac rate from 136 ± 17 to 89 ± 14 bpm, increased the QRS complex duration from 27 ± 3.1 to 59 ± 11 ms and increased the QT interval from 229 ± 23 to 264 ± 19 ms. Additionally, it increased the conduction time between the right atrium and the His bundle ($27 \pm 6.5\%$), reduced the intra ventricular pressure ($86 \pm 6\%$) and decreased the conduction velocity of atrial impulse ($17 \pm 3\%$). However, the transmission time from the His bundle to the right ventricle was not altered. Atropine sulfate did not change moreover the electrocardiographic parameters or the intra ventricular pressure effects promoted by the *A. carambola* aqueous extract. Therefore, the learn suggested avoiding the use of such extracts since it can induce electrical and mechanical changes in the normal heart.

Hypotensive activity

Soncini *et al.*, (2011) studied the hypotensive effect of the aqueous extract of *A. carambola* and its underlying mechanisms using isolated rat aorta. *In vitro*, the aqueous extract caused a reduction in the E_{max} response to phenylephrine devoid of a change in sensibility. Additionally, in a depolarized Ca^{2+} -free medium, the aqueous extract inhibited $CaCl_2$ -induced contractions and caused a concentration-dependent rightward shift of the response curves, suggesting that the aqueous extract inhibited the contractile mechanisms involving extracellular Ca^{2+} influx. The study concluded the aqueous extract to be hypotensive in nature and suggested the effects to be in part, due to the inhibition of Ca^{2+} , which is in conformity with its uses in conventional medicine.

Hypocholesterolemic activity

Wu *et al.*, (2009) examined the potential hypocholesterolemic activity of dissimilar insoluble fibers prepared from *A. carambola* with or devoid of micronization processing. After micronization, the cation-exchange and water-holding capacities of the *A. carambola* pectic polysaccharide-rich unsolvable fibres were effectively increased from 8.5 to 22.4 mL/g. The *A. carambola* micronized insoluble fibers abridged the concentrations of serum triglyceride by and total cholesterol by 15.6 % and 15.7 %, correspondingly by means of enhancing the excretion of cholesterol and bile acids in the feces. The study suggests a new move near of micronization of the fruit which may help to progress physiological functions of food fibers in fiber-rich functional food applications. The study also indicates that the particle size is an important factor in affecting the characteristics and physiological functions of insoluble fibers.

Hypoglycaemic activity

Ferreira *et al.*, (2008) examined the effects of the hydroalcoholic extract of leaves of *A. carambola* L. on fasting blood glucose. The hydroalcoholic extract treated animals showed considerably inferior fasting blood glucose ($p < 0.05$). In contrast, livers from hydroalcoholic extract of these animals showed considerably superior ($p < 0.05$) glucose production from L-alanine. This consequence was mediated, at least part of it, by an activation of the catabolism of L-alanine inferred by the augmented hepatic urea and L-lactate production. Unlike L-alanine, the glucose production from L-glutamine, L-lactate and glycerol was not affected by the treatment. Likewise, the hydroalcoholic extract conduct did not affect the glucose uptake in soleus muscles, inferred by the incorporation of $[14C]$ -glucose to glycogen (glycogen synthesis) and $[14C]$ -lactate production. Thus, the study suggests that lessening of fasting blood glucose promoted by the treatment with hydroalcoholic extract of *A. carambola* was not mediated by an inhibition of hepatic gluconeogenesis and/or an augmented glucose uptake by muscles. Chau *et al.*, (2004b) investigated the hypoglycaemic effects of numerous insoluble fiber-rich fractions including insoluble dietary fiber, alcohol insoluble solid and water-insoluble solid obtained from the pomace of *A. carambola*, by some *in vitro* methods. It was establish that the three insoluble fiber rich fractions could efficiently adsorb glucose, retard glucose diffusion, reschedule the release of glucose from starch, and inhibit the activity of α -amylase to different extents. The hypoglycemic effects of these unsolvable fiber-rich fractions were considerably ($P < 0.05$) stronger than that of cellulose. Thus, it was accomplished that these fiber-rich fractions could be incorporated as low-calorie bulk ingredients in high fiber foods to reduce calorie level and assist the control of blood glucose level.

Nephrotoxic effect

Neto *et al.*, (2009) studied the effects of *A. carambola* fruit on patients with renal failure, who are not yet on dialysis and accomplished that the fruit and its juice induce nephrotoxicity and neurotoxicity, which may occasionally be fatal. Patients with chronic kidney disease (stages 3–5 but not on dialysis) should be forbidden from eating the fruit. People with normal renal function must also be warned to avoid eating of large amounts of the fruit or juice particularly on a vacant stomach.

Neurotoxic effect

Carolino *et al.*, (2005) examined the effects of a neurotoxic fraction isolated from *A. carambola* on GABAergic and glutamatergic transmission systems. The fraction was obtained by chronological chromatography of the aqueous extract on anion swap column, then on cation exchange column and, finally on reversed-phase high performance liquid chromatography (HPLC). The fraction did not affect GABA/glutamate uptake or release, or on glutamate binding, but effected GABA binding in a concentration-dependent manner (IC_{50} 0.89 mM). Video-electroencephalogram recordings showed that cortical management of neurotoxic fraction, induce behavioral changes in animals, including tonic-clonic seizures, evolving into status epilepticus, accompanied by cortical epileptiform activity. The neurotoxic fraction has not yet been chemically characterized but the preliminary data suggested that it contains a non peptide molecule conflicting from oxalic acid and bearing a molecular weight less than 500.

Phytochemistry

Chemical constituents isolated from *A. carambola*

Averrhoa carambola has been the subject of phytochemical studies since late 1970s. The following are the chemical constituents isolated for *A. carambola*.

C₁₃-aroma compounds

Herderich *et al.*, (1992) reported identifications of glycosidically bound constituents from star fruit (*A. carambola* L.) using HRGC and HRGC-MS techniques. The constituents were obtained from the extracts by Amberlite XAD-2 adsorption followed by methanol elution. The compounds identified were the ionone derivatives, namely 4-hydroxy- β -ionol, 3-hydroxy- β -ionol, 4-oxo- β -ionol, 3-hydroxy- β -ionone, 3-oxo- α -ionol, 3-oxo-*retro*- α -ionol (2 isomers), 3-oxo-4,5-dihydro- α -ionol, 3-oxo-7,8-dihydro- α -ionol (blumenol C), 3,5-dihydroxy-megastigma-6,7-diene-9-one (grasshopper ketone), 3-hydroxy- β -damascone, 3-hydroxy-5,6-epoxy- β -ionone, 3-hydroxy-5,6-epoxy- β -ionol, 3,4-dihydro-3-hydroxyactinidol, vomifoliol (blumenol A), 4,5-dihydrovomifoliol and 7,8-dihydrovomifoliol (Blumenol B). Several of these new constituents are easily degraded upon heat-treatment at natural pH condition of the fruit pulp, thus rationalizing the formation of a number of C₁₃-aroma compounds, which have recently been reported as star fruit volatiles.

B-Galactosidase

Balasubramaniam *et al.*, (2005) isolated β -Galactosidase (EC. 3.2.1.23) from *A. carambola* fruit and fractionated it using a combination of ion exchange and gel filtration chromatography into four isoforms, β -Galactosidase I, II, III and IV. The β -galactosidases exhibited the molecular weights 84, 77, 58 and 130 kDa, respectively. B-Galactosidase I, was the most prominent isoform. The purified β -galactosidase I was highly active in hydrolyzing (1 \rightarrow 4) β -linked spruce and a mixture of (1 \rightarrow 3) β - and (1 \rightarrow 6) β -linked gum arabic galactans. The β -galactosidase I also exhibited the capacity to solubilize and depolymerize structurally intact pectins as well as to modify alkaline-soluble hemicelluloses, indicating in part changes that occur during ripening.

Pharmacognostic and phytochemical analysis

Thomas *et al.*, (2008) conducted preliminary pharmacognostic and phytochemical analysis and showed the presence of saponins, tannins, alkaloids and flavonoids. Furthermore, the study revealed that the water-soluble and alcohol soluble extractive values decreased gradually with ripening of fruit.

Analysis of oxalic acid

Wilson *et al.*, (1982) quantified oxalic acid in 15 Florida-grown *A. carambola* cultivars using simple and rapid high-performance liquid chromatographic (HPLC) method. Oxalic acid levels were found to vary among the cultivars examined from 0.08 to 0.73 g per 100/g of fruit.

Fruit and fibre-rich fractions

Compositions

Fruit composition

Narain *et al.*, (2001) studied the composition of the fruit during maturation. The pH of the fruit increased with the advance in maturity, whereby the pH values were 3.44, 2.40 and 2.71 for ripe, green mature and half-ripe fruits, respectively. Increased calcium contents were observed at ripe stage (4.83 ± 0.27 mg/100 g of edible fruit) and it was significantly different than the fruits at green mature (3.55 ± 0.85 mg/100 g of edible fruit) or half-ripe stages (4.83 ± 0.27 mg/100 g of edible fruit). The titratable acidity, reducing sugars and tannin contents of the fruits vary significantly among the fruits at different stages of maturity.

Fiber-rich fractions

Chau *et al.*, (2004a) reported that the pomace of *A. carambola* possess a high level of insoluble fiber-rich fractions including insoluble dietary fiber, alcohol insoluble solid, and water-insoluble solid (46.0–58.2 g/100 g of pomace). These fiber-rich fractions were mostly composed of pectic substances and hemicellulose. The physicochemical properties of these fiber-rich fractions (water-holding capacities, swelling properties, and cation-exchange capacities) were significantly ($P < 0.05$) higher than those of cellulose. The capacity of these fiber-rich fractions to adsorb glucose and reduce amylase activity implied that they might help control postprandial serum glucose. Thus the study suggested that the consumption and application of the insoluble fiber-rich fractions as low-calorie bulk ingredients in fiber enrichment.

Clinical studies

Clinical symptoms and outcomes in uraemic patients given *A. carambola*

Investigations of the clinical symptoms and outcomes of 32 uraemic patients ingesting star fruit showed that most common symptoms were persistent and intractable hiccups (in 30 patients; 93.75%), vomiting (in 22 patients; 68.7%), variable degrees of disturbed consciousness (mental confusion, psychomotor agitation; in 21 patients; 65.6%), decreased muscle power, limb numbness, paresis, insomnia and paresthesias (in 13 patients; 40.6%) and seizures (in seven patients; 21.8%). Patients who were promptly treated with hemodialysis, including those with severe intoxication, recovered without sequelae. The study concluded that

hemodialysis, particularly on a daily basis, is the best treatment for star fruit intoxication. In adverse cases, continuous methods of replacement therapy may provide a superior initial procedure, since rebound effects are a common event. Peritoneal dialysis is of no use as a treatment, especially when consciousness disorders ensue (Neto *et al.*, 2003).

Conclusion

A. carambola has long been used in traditional medicine and has demonstrated pharmacological activities which support its use in traditional uses as an important therapeutic agent. However, the scientific evidences are limited in respect to its bioactive constituents, bioavailability, pharmacokinetics, and therapeutic importance including clinical trials. Thus, future work on bioactivity-guided isolation of the active secondary metabolites responsible for the pharmacological activities of *A. carambola*, and studies on their structure-activity relationships, Mechanisms of actions, pharmacokinetics and toxicity are required for the development of *A. carambola* as a therapeutic agent. Furthermore, the extracts or fractions with potential bioactivities may be standardized and employed in clinical studies.

References

1. Ali MA, Nayan V, Chanu KV, Ralte L, Devi LI. 2011. Antioxidant activity of fruits available in Aizawal Market of Mizoram India. *World J Agric Sci* 7: 327 - 332.
2. Araho D, Masazumi M, Wen-hua C, Toshimitsu K, Kenji M, Takao I. 2005. A new flavone Cglycoside from the leaves of *Averrhoa carambola*. *National Medical (Tokyo)* 59: 113 - 116.
3. Artani N, Yeli M, Mohammed H. 2006. Isolation and identification of active antioxidant compound from star fruit (ethanol extract). *J Applied Sci* 6: 1659 - 1663.
4. Avinash P, Swapneel K, Darshana P, Anita P. 2012. A comprehensive review of an important medicinal plant – *Averrhoa carambola* L. *Pharmacog Comm* 2: 13 - 17.
5. Balasubramaniam S, Lee HC, Lazan H, Othman R, Ali ZM. 2005. Purification and properties of a β - galactosidase from carambola fruit with significant activity towards cell wall polysaccharides. *Phytochemistry* 66: 153 - 163.
6. Cabrini DA, Moresco HH, Imazu P, Silva CD, Pietrovski EF, Mendes DAGB, Prudente AdS, Pizzolatti MG, Brighente I, Costa sM, Otuki MF. 2011. Analysis of the Potential Topical Anti-Inflammatory Activity of *Averrhoa carambola* L. in Mice. *Evid-Based Compl Alt Med* Doi: 10.1093/ecam/nej026 Article ID908059
7. Carolino ROG, Belebony RO, Pizzo AB, Vecchio FD, Garcia-Cairasco N, Moyses-Neto M, Santos WFd, Coutinho-Netto J. 2005. Convulsant activity and neurochemical alterations induced by a fraction obtained from fruit *Averrhoa carambola* (Oxalidaceae: Geraniales). *Neurochemistry International* 46: 523 - 531.
8. Chau CF, Chen CH, Lee MH. 2004a. Characterization and physicochemical properties of some potential fibres derived from *Averrhoa carambola*. *Food Nahrung* 48: 43 - 46.
9. Chau CF, Chen CH, Lin CY. 2004b. Insoluble fiberrich fractions derived from *Averrhoa carambola*: hypoglycemic effects determined by in vitro methods. *LWT - Food Sci Technol* 37: 331 - 335.
10. Ferreira EB, Fernandes LC, Galende SB, Cortez DAG, Bazotte RB. 2008. Hypoglycemic effect of the hydroalcoholic extract of leaves of *Averrhoa carambola* L. (Oxalidaceae). *Rev Bras Farmacogn* 18: 339 - 343.
11. Gheewala P, Kalaria P, Chakraborty M, Kamath JV. 2012. Phytochemical and Pharmacological profile of *Averrhoa carambola* Linn: An overview. *Int Res J Pharm* 3: 88 - 92.
12. Goncalves ST, Baroni S, Bersani-Amado FA, Melo GAN, Cortez DAG, Bersani-Amado CA, Cuman RKN. 2006. Preliminary studies on gastric anti-ulcerogenic effects of *Averrhoa carambola* in Rats. *Acta Farmac Bonaerense (Latin Am J Pharm)* 25: 245 - 247.
13. Gunasegaran R. 1992. Flavonoids and anthocyanins of three Oxalidaceae. *Fitoterapia* 63:89 - 90.
14. Guanghou S, Lai PL. 2004. Analysis of polyphenolic antioxidants in star fruit using liquid chromatography and mass spectrometry. *J Chromatography A* 1022: 67 - 75.
15. Henrique H, Moresco, Gustavo S, Queiroz, Moacir G, Pizzolatti, Inês MC, Brighente. 2011. Chemical constituents and evaluation of the toxic and antioxidant activities of *Averrhoa carambola* leaves. *Braz J Pharmacogn* 22: 319 - 324.
16. Herderich M, Neubert C, Winterhalter P, Schreier P, Skouroumounis GK. 1992. Identification of C13-norisoprenoid flavour precursors in starfruit (*Averrhoa carambola* L). *Flav Frag J* 7: 179 - 185.
17. Li X, Yang L, Yu R, Zhu J, Tian T, Song G, Song L. 2012. Biotransformation of dihydro-epideoxyarteannuin B by suspension-cultured cells of *Averrhoa carambola*. *Afr J Biotechnol* 11: 1724 - 1728.
18. Luximon-Ramma A, Bahorun T, Crozier A. 2003. Antioxidant actions and phenolic and vitamin C contents of common Mauritian exotic fruits. *J Sci Food Agric* 83: 496 - 502.
19. Manda H, Vyas K, Pandya A, Singhal G. 2012. A complete review on: *Averrhoa carambola*. *World J Pharm Pharmac Sci* 1: 17 - 33.
20. Margen S. 1992. *The Wellness Encyclopedia of Food and Nutrition*, New York: Health Lett Assoc 271 - 272.
21. Mia MM, Rahman MS, Begum K, Begum B, Rashid MA. 2007. Phytochemical and biological studies of *Averrhoa carambola*. *Dhaka Univ J Pharmac Sci* 6: 125-128.
22. Moresco HH, Queiroz GS, Pizzolatti MG, Brighente IMC. 2012. Chemical constituents and evaluation of the toxic and antioxidant activities of *Averrhoa carambola* leaves. *Rev Bras Farmacogn* 22: 319 - 324.
23. Morton JF. 1987. *Fruits of warm climates*, Miami: Flair Books 125 - 128.
24. Narain N, Bora PS, Holschuh HJ, Vasconcelos Da S. MA. 2001. Physical and chemical composition of carambola fruit (*Averrhoa carambola* L.) at three stages of maturity. *Food Science and Technology. Ciencia y Tecnologia Alimentaria* 3: 144 - 148.
25. Neto MM, Silva GEB, Costa RS, Vieira Neto OM, Garcia-Cairasco N, Lopes NP, Haendchen PFC, Silveira C, Mendes AR, Filho RR, Dantas M. 2009. Star fruit: simultaneous neurotoxic and nephrotoxic effects in people with previously normal renal function. *Nephrology Dialysis Transplantation Plus* 2: 485 - 488.
26. Neto MM, Costa JAC, Garcia- Cairasco N, Netto JC, Nakagawa B, Dantas M. 2003. Intoxication by star fruit (*Averrhoa carambola*) in 32 uraemic patients: treatment and outcome. *Nephrology Dialysis Transplantation* 18: 120 - 125.

27. Ranganayaki S, Singh R, Singh AK. 1980. The chemical examination of the bark of *Avehrroa carambola*. Proc Nat Acad Sci India - Section A 50: 61 - 63.
28. Shui G, Leong LP. 2006. Residue from star fruit as valuable source for functional food ingredients and antioxidant nutraceuticals. Food Chem 97: 277 - 284.
29. Soncini R, Santiago MB, Orlandi L, Moraes GOI, Peloso ALM, dos Santos MH, Alves-da-Silva G, Paffaro Jr VA, Bento AC, Giusti-Paiva A. 2011. Hypotensive effect of aqueous extract of *Averrhoa carambola* L. (Oxalidaceae) in rats: An in vivo and in vitro approach. J Ethnopharmacol 133: 353 - 357.
30. Sung CK, Kimura T, But PPH, Guo J-X. 1998. International Collation of Traditional and Folk Medicine: Northeast Asia, Singapore: World Scientific Publishing Company: 75-76.
31. Tadros SH, Sleem AA. 2004. Pharmacognostical and biological study of the stem and leaf of *Avehrroa carambola* L. Bull Fac Pharm 42: 225 - 246.
32. Teh CC, Khoo ZY, Khursiah F, Rao NK, Chin JH. 2010. Acetylcholinesterase inhibitory activity of star fruit and its effect on serum lipid profiles in rat. Int Food Res J 17: 987 - 994.
33. Thomas S, Patil DA, Patil AG, Chandra N. 2008. Pharmacognostic evaluations and physicochemical analysis of *Averrhoa carambola* L. fruit. J Herb Med Toxicol 2: 51 - 54.
34. Tiwari KP, Masood M, Minocha PK. 1979. Chemical constituents of *Gemlina philippinensis*, *Adenocalymna nitida*, *Allamanda cathartica*, *Averrhoa carambola* and *Maba buxifolia*. J Ind Chem Soc: 56: 944.
35. Vasconcelos CML, Araújo MS, Conde-Garcia EA. 2006. Electrophysiological effects of the aqueous extract of *Averrhoa carambola* L. leaves on the guinea pig heart. Phytomedicine 13: 501 - 508.
36. Vasconcelos CML, Araújo MS, Silva BA, Conde- Garcia EA. 2005. Negative inotropic and chronotropic effects on the guinea pig atrium of extracts obtained from *Averrhoa carambola* L. leaves. Braz J Med Biol Res 38: 1113 - 1122.
37. Vasconcelos CML, Gondim ANS, Cruz JS, Mafra RA, Silva BA, Conde-Garcia EA. 2008. Aqueous leaf extract of *Averrhoa carambola* L. (Oxalidaceae) reduces both the inotropic effect of BAY K 8644 on the guinea pig atrium and the calcium current on GH3cells. Rev Bras Farmacog 18: 539 - 543.
38. Wilson CW, Shaw PE, Knight RJ. 1982. Analysis of oxalic acid in carambola (*Averrhoa carambola* L.) and spinach by high-performance liquid chromatography. J Agric Food Chem 30: 1106 - 1108.
39. Wu SC, Wu SH, Chau CF. 2009. Improvement of the hypocholesterolemic activities of two common fruit fibers by micronization processing. J Agric Food Chem 57: 5610 - 5614.

