

# Alleviation of salinity stress mediated by Salicylic acid in wheat: An overview

Rajan Saini

School of Agriculture, Department of Agronomy, Lovely Professional University,  
Phagwara, (Punjab), 144411, India.

**Abstract:** Salicylic acid (SA) shows major effect on plant resistance to various environmental stress factors. Wheat plants treated with salicylic acid shows an increment in the level of cell division inside the apical meristem of seedling roots which results in increased plant growth. Adverse environmental conditions or stress like heat stress, salinity, and water stress hampers the growth of wheat. To mitigate this adverse effect of various types of stress we use SA of different concentrations in wheat either by foliar application or by seed priming. SA accelerates the growth and development of plant during stress conditions so it is known as the plant growth promoter.

## Introduction:

Wheat (*Triticum aestivum* L.) is a most popular cereal crop around the world and it come under family Poaceae. Wheat gives about 55% of the starches and 20% of the food calories burned through worldwide (Breiman and Graur, 1995). Wheat is getting cultivated under wide range of climatic conditions. India secured first position in area but ranks second in production after China with the average productivity of 3117 kg/ha. Wheat cultivation is supposed to begun in India 5000 years ago (Feldman, 2001). Today, India positions second in wheat production worldwide. It's area under cultivation is 28MHa. To tackle increasing demand of wheat without increasing area, we need to incorporate new physiological tools. These tools will add up for the advancement of breeding programme under abiotic stress environment.

**Salicylic acid:** Salicylic acid ( $C_7H_6O_3$ ) is a beta hydroxyl acid also known as an endogenous growth regulator of phenolic type, which affects a range of different processes in plants, including seed germination, stomatal closure, ion up-take and transport, membrane permeability photosynthetic and growth rate. SA is additionally significant signal molecule for adjusting plant responses to environmental stress. SA is well known to provide protection against a number of abiotic stresses (Barkosky and Einhellig, 1993; Harper and Balke, 1981; Cutt and Klessig, 1992; Senaratna *et al.*, 2000).

**Salinity:** Wheat production is significantly influenced by salinity stress. The cellulose content, lignin content of both shoots and roots show downfall with rise in Sodium Chloride concentration. Expansion in NaCl fixation results in increased  $Na^{2+}$  and  $Ca^{2+}$  accumulation and lowers  $K^{+}$  and  $Mg^{2+}$  concentration in various organs of wheat plant. Soaking of grains in SA could result in handling the antagonistic impacts of NaCl

salinity by inducing the accumulation of proline in wheat seedlings. Therefore, salinity plays a major role in the growth and development of wheat. Therefore, salinity should be neutralised in order to get higher wheat yields.

**Salinity stress in wheat:** High salinity results in lower germination rate, photosynthesis, transpiration, and higher accumulation of  $\text{Na}^+$  and  $\text{Cl}^-$  ions which disturb the normal metabolic processes i.e. germination, growth, photosynthesis, water relation, cellular damage, ion uptake, yield of wheat plants. Higher salinity results in stomatal closure in wheat hence reduce  $\text{CO}_2$  availability in the leaves and inhibits carbon fixation. Hasanuzzaman *et al.*, (2017). Emergence percentage of wheat under salinity stress is greatly reduced as compared to non-saline conditions.

Root dry weight and shoot dry weight also gets hampered under salinity. Chlorophyll content and osmotic potential also shows negative results in wheat due to the influence of salicylic acid.

#### Response of salinity stress:

- 1- **OS induction:** Under severe saline conditions, rate of ROS generation exceeds the scavenging potential of cellular defense system resulting in oxidative stress. Oxidative stress damages cellular components of wheat plant which results in dis-function and finally the death of plant.
- 2- **Reduced water uptake:** High salt accumulation leads to osmotic stress and causes low water potential which results in reduced water uptake which ultimately results in the drought stress and in severe cases death of the plant takes place.
- 3- **Ion toxicity:** Under saline conditions, there is higher accumulation of  $\text{Na}^+$  and  $\text{Cl}^-$  ions which results in ion toxicity.
- 4- **Nutrient imbalance:** Under saline conditions the uptake of major primary nutrients N, P, K is greatly affected. Due to this nutrient deficiency the normal metabolic processes of wheat are hampered.
- 5- **Hampered growth:** Under moderate saline conditions wheat plant shows hampered growth as growth processes are hampered. There will be low germination percentage, reduction in plant height and other morphological characteristics (Nadeem *et al.*, 2013).
- 6- **Reduction in yield:** Salinity directly affects the yield as it negatively affects the growth processes in wheat plants. Due to this detrimental effect in growth factors, normal metabolic processes are disturbed and there is reduction in yield (Zeng, 2001 and Carillo, 2001).
- 7- **Cellular damage:** Uncertain growth and irregular uptake of water and other essential nutrients ultimately result in destruction of cell membrane properties of wheat plants. Wheat seedlings experienced lipid peroxidation, accumulation of hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), and increased membrane permeability under salt stress. Cellular damage mainly occurs under highly saline conditions (Nxele, *et al.*, 2017 and Shabala and Munns, 2012).

## Role of Salicylic under Salt Stress:

Antioxidants help in maintaining the reactive oxygen species levels in plants. Phenolic compounds and tocopherol are one of the examples. The higher concentrations of antioxidants in plants help to overcome salt stress. Therefore, Salicylic acid plays a major role in controlling salinity stress by inducing antioxidants. Osmolytes are small and minute molecules which are mostly soluble (AbdElgawad et al., 2016; Gupta and Huang, 2014). To mitigate the salt stress, plants accumulate various osmolyte like proline, glycine and betaine (Qureshi *et al.*, 2013 and Siddique *et al.*, 2018). The phenolic group compounds are secondary metabolites in plants. Salicylic acid is also one of them. It acts a signaling molecule that respond against environmental stress in which salinity stress is one of them. Salicylic acid helps the plants in the absorption of mineral nutrients from soil during salinity stress which helps the plant to stay strong. It regulates the osmotic potential in favour of the plant. Plant hormones are the chemicals that are produced within plants in minute quantity that helps the plants in regulating normal metabolic processes in adverse environmental conditions. Some plant hormones like Salicylic acid produced within plants in minute quantity which help to overcome against stress. So that overall these compound mediate in ROS detoxification and in the protection of photosynthetic system which ultimately leads in alleviating the cellular damage against salt stress (Das and Roychoudhury, 2014 and Sharma et al., 2012)

## Conclusion:

Soil salinity is a major issue in agriculture that leads to great loss in economics. Salinity at lower concentrations causes hampered growth while at higher concentrations leads towards the death of the plant. Plant treated with salicylic acid, leads to produce Osmolytes, Phenolic compounds and stress growth hormone while also lead to enhance the level of nutrient and water uptake. So that over all ROS detoxification get activate hence the cell and its membrane become protected against salinity stress.

## References

- Breiman, A., & Graur, D. (1995). Wheat evolution. *Israel Journal of Plant Sciences*, 43(2), 85-98.
- Shaked, H., Kashkush, K., Ozkan, H., Feldman, M., & Levy, A. A. (2001). Sequence elimination and cytosine methylation are rapid and reproducible responses of the genome to wide hybridization and allopolyploidy in wheat. *The Plant Cell*, 13(8), 1749-1759.
- Barkosky, R. R., & Einhellig, F. A. (1993). Effects of salicylic acid on plant-water relationships. *Journal of Chemical Ecology*, 19(2), 237-247.
- Cutt, J. R., & Klessig, D. F. (1992). Pathogenesis-related proteins. In *Genes involved in plant defense* (pp. 209-243). Springer, Vienna.

- Harper, J. R., & Balke, N. E. (1981). Characterization of the inhibition of K<sup>+</sup> absorption in oat roots by salicylic acid. *Plant Physiology*, 68(6), 1349-1353.
- Senaratna, T., Touchell, D., Bunn, E., & Dixon, K. (2000). Acetyl salicylic acid (Aspirin) and salicylic acid induce multiple stress tolerance in bean and tomato plants. *Plant Growth Regulation*, 30(2), 157-161.
- Hasanuzzaman, M., Nahar, K., Rahman, A., Anee, T. I., Alam, M. U., Bhuiyan, T. F., ... & Fujita, M. (2017). Approaches to enhance salt stress tolerance in wheat. *Wheat Improvement, Management and Utilization*, 151-187.
- AbdElgawad, H., Zinta, G., Hegab, M. M., Pandey, R., Asard, H. and Abuelsoud, W. (2016). High salinity induces different oxidative stress and antioxidant responses in maize seedlings organs. *Frontiers in plant science*, 7, 276.
- Carillo, P., Annunziata, M. G., Pontecorvo, G., Fuggi, A., and Woodrow, P. (2011). Salinity stress and salt tolerance. *Abiotic stress in plants—Mechanisms and adaptations*, 1, 21-38.
- Das, K., and Roychoudhury, A. (2014). Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Frontiers in environmental science*, 2, 53.
- Gupta, B., and Huang, B. (2014). Mechanism of salinity tolerance in plants: physiological, biochemical, and molecular characterization. *Int. J Genomics* 2014:18. doi: 10.1155/2014/701596.
- Nadeem, S. M., Zahir, Z. A., Naveed, M., and Nawaz, S. (2013). Mitigation of salinity-induced negative impact on the growth and yield of wheat by plant growth-promoting rhizobacteria in naturally saline conditions. *Annals of Microbiology*, 63(1), 225-232.
- Nxele, X., Klein, A. and Ndimba, B. K. (2017). Drought and salinity stress alters ROS accumulation, water retention, and osmolyte content in sorghum plants. *South African Journal of Botany*, 108, 261-266.
- Qureshi, M., Abdin, M., Ahmad, J., and Iqbal, M. (2013). Effect of long-term salinity on cellular antioxidants, compatible solute and fatty acid profile of Sweet Annie (*Artemisia annua* L.). *Phytochemistry* 95, 215–223. doi: 10.1016/j.phytochem.2013.06.026.
- Shabala, S., and Munns, R. (2012). Salinity stress: physiological constraints and adaptive mechanisms. *Plant stress physiology*, 1(1), 59-93.
- Sharma, P., Jha, A. B., Dubey, R. S., and Pessarakli, M. (2012). Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of botany*, 2012.
- Siddique, A., Kandpal, G., and Kumar, P. (2018). Proline accumulation and its defensive role under Diverse Stress condition in Plants: An Overview. *Journal of Pure and Applied Microbiology*, 12(3), 1655-1659.
- Zeng, L., Shannon, M. C., and Lesch, S. M. (2001). Timing of salinity stress affects rice growth and yield components. *Agricultural Water Management*, 48(3), 191-206.