



ASSESSMENT OF STORAGE AND HANDLING PRACTICES ON QUALITY OF HOUSEHOLD DRINKING WATER

Sandhya Tambekar Wanjari

Department of Microbiology

Dhote Bandhu Science College,

Gondia – 441614, Maharashtra (India)

E-mail: drsandhyatambekarwanjari@gmail.com

<https://orcid.org/0000-0002-6226-826X>

Abstract:

Waterborne diseases constitute a major public health problem in India and consumption of contaminated water causes health risks to the public. Safe drinking water is essential to the health and well-being of human beings. Contamination in stored household drinking water is associated with the methods of water storage, handling practices, and domestic and personal hygiene practices. To find out the probable cause of the contamination of drinking water; a survey was conducted in 100 families and household drinking water quality (before and after use) was analyzed for its potability. The objective of this study was to assess the contamination potentials of different house water handle portability rage practices The study indicated that the storage container and their hygienic practices, number of family members, children in the family, and their socioeconomic status, education of housewives significantly affect the quality of drinking water in the houses.

(Keywords: Household water, potability of water, Personal and domestic hygiene)

Introduction:

Household water storage remains a necessity in many communities worldwide, especially in developing countries. Water storage often using tanks/vessels is envisaged to be a source of water contamination, along with related user practices (Musa, 2021). Access to clean and safe water, good sanitation, and hygiene practices are necessary for a healthy population. The importance of education, sociocultural acceptance, and changing people's beliefs and behaviors achieved sustainability and affordability in the provision of safe water (Agensi, 2019). Provision of safe household water included the conditions and practices of water collection, storage, handling, and the choice of storage containers or vessels (Chalchisa *et al*, 2017; Tambekar and Mahore, 2005). Microbial contamination of collected and stored household water is caused not only by the collection and use but unsanitary methods to dispense water from household storage vessels, including contaminated hands and dippers, and inadequate cleaning of vessels, which lead to the accumulation of sediments and pathogens. Even though water is treated before supplies but may become contaminated after collection, either during transport or storage in the home ((Alexander *et al*, 2019, Duru *et al*, 2013).

The reasons for the contamination of drinking water in the home may be due to personal and domestic inadequate and unsanitary storage conditions that allowed, for the introduction and proliferation of disease-causing microbes (Jagals, *et al*, 2003). Personal and domestic hygienic practices indirectly depend on the education of the family members, as educated people are more conscious about their health and always try to maintain hygienic conditions in their homes which leads to contamination in drinking water ((Tambekar, *et al* 2016). Introduction of hands or fingers or dippers into stored water contaminants the water which

suggested the importance of hand washing, and domestic and personal hygiene practices might have prevented the chance of infection in a community (Nala *et al.*, 2003; Yvan Hutin *et al.*, 2003).

Hence a study was conducted on the personal and domestic hygienic practices in houses that affect the quality of household drinking water. The factors such as the education of housewives, socioeconomic status of the family, personal and domestic hygiene, number of family members, water handling and withdrawal practices, etc were studied for keeping the water free from diseases-producing microorganisms.

Materials and Methods:


The study was conducted to observe the effect of handling and personal hygienic practices along with the education and socio-economic status, in houses on the quality of drinking water after storage in the container. For the study, 100 families were identified from the various colonies of Gondia city, and data was collected as per prepared questionnaires from respective houses. Two samples of drinking water, one before and the other after the use of water were collected from storage containers for the potability test of each house. A total of 200 drinking water samples were collected and Manja's H₂S test was performed for potability (Manja *et al.*, 2001) by adding approximately 20 ml water sample into the medium contain bottle. These bottles were then incubated at room temperature for 24, 48, and 72 h to determine the extent of blacking in the bottle by producing hydrogen sulfide gas. The MPN was calculated by Multiple Tube Fermentation Technique (MTFT); nine multiple tube dilution techniques using double and single-strength Bromo-Cresol Purple MacConkey medium, by adding 10ml, 1ml, 0.1ml water samples in appropriate test medium and incubated for 24-48h at 37°C. The MPN Index was calculated from the MPN table and the index of water more than 10coliforms/dl was designated as polluted or unhealthy for drinking purposes or non-potable (APHA Standard Methods for the Examination of Water and Wastewater, 23rd Edition, 2017).

The data were collected using interview and observation methods. In the interview; information on the education of spouse, number of children, the income of a family, method of collection and storage of water, water handling practices, etc. were collected from the hundred households from where water samples were collected for analysis, by filling a prepared questionnaire. Collected data were analyzed by using SPSS software about the potability of drinking water.

Results and Discussion:

Waterborne diseases constitute a major public health burden in India and consumption of contaminated water causes health risks to the public, and the situation is alarming in rural areas. The most common and widespread health risk associated with drinking water is contamination, either directly or indirectly by the method of water storage, handling practices, domestic and personal hygiene practices (Packiyam *et al.*, 2016). This study aimed to assess the contamination potentials of different house water handling and storage practices in Gondia. To find out the probable cause of the contamination in drinking water this survey was conducted and household drinking water samples (before and after use) were analyzed for coliform contamination.

The water sample from 100 families was collected and analyzed for potability of the drinking water and found that, out of these, 82% were found to be potable and 18% non-potable. The water quality after use in those families using earthenware without a tap showed 90% potable while earthenware with a tap showed 100% portable. The water in stainless steel and plastic container showed 89 and 3% potable after use. The data showed water in the water filter was 100% potable (Fig. 2).






The withdrawing utensil may be one of the causes which made the potable drinking water non-potable because while withdrawing water there was the chance of water getting contaminated with the fingers of a person or with contaminated dipper. Families, who were using a dipper without a handle for water withdrawal from the container, the water was 77% potable as compare to families either using the short handle (86% potability) or long handle dipper or tap for withdrawing water (100% potable). Long or short handle dipper or tap reduces the chances of contact of hands or fingers with the water and prevents secondary contamination in the stored water. Moreover washing the hands before collecting or withdrawing water reduced the chances of contamination up to zero percent and kept the water potability 100% (Fig. 2).

The socioeconomic condition of families is one factor that also plays an important role in contaminating the household drinking water (Gorter *et al.*, 1998). The families from the lower income group kept the water up to 77% potable after use while middle-class families kept it 89% and the of the higher middle class kept it 94% potable. This indicates that the socio-economical status of the family also affects the hygienic practices of the family (Fig. 2).

The education of spouses, the number of family members, and their hygienic practice also have a prominent role in making potable water into non-potable (Kern and Harada, 2002; WHO, 2003). The illiterate spouses kept the water potable up to 78% while the primary educated kept it 89%, and the SSC, HSC, and graduate kept it 91%, 93%, and 95% respectively. It indicates that the education of the housewife or family members plays a prominent role in personal and domestic hygiene which leads to good hygienic practices and the quality of household drinking water ((Fig. 2).

The quality of drinking water is inversely proportional to the number of family members. A family with more numbers contaminates the water highly as compared to a family with few members, as they used it often (Tambekar, 2017). This indicated that the more the family member more the contamination because the more the members in the family, the more times of use dipper or handling which results in more contamination. The families with 3 members, 5 members, and 7 members kept the water 87%, 82%, and 79% potable after use. This also indicates that the number of children may affect the quality of drinking water because children may withdraw water unhygienically, such as without washing their hands or directly dipping their hands in the stored water, which makes water contaminated. Washing the water pot or dipper also affects the quality of drinking water indicating that a higher frequency of washing water containers or dippers is good quality or potability of the water (Table 2).



The study recommended that long or short-handle dippers should be used for the withdrawal of water from the storage container. The education of the housewife also affects the storage, handling, and hygienic practices of drinking water. The r educated housewives stored water more hygienically as they were more conscious about health as compared to illiterate housewives. Socioeconomic status also affects the hygienic conditions of the family and invariably affects the quality of water storage. The higher the monthly income higher the hygienic quality of drinking water in the houses. At the same time the family member or children, have poor quality water as everybody may not use proper hygienic procedures or wash their hands or dipper before withdrawing the water. Thus, the study indicated that the storage and handling practices, number of family members or children in the family, socio-economic status of the family, and their hygienic practices significantly affect the quality of drinking water in the houses

The study recommended that long or short-handle dippers should be used for the withdrawal of water from the storage container. The education of the housewife also affects the storage, handling, and hygienic practices of drinking water. The r educated housewives stored water more hygienically as they were more conscious about health as compared to illiterate housewives. Socioeconomic status also affects the hygienic conditions of the family and invariably affects the quality of water storage. The higher the monthly income higher the hygienic quality of drinking water in the houses. At the same time the family member or children, have poor quality water as everybody may not use proper hygienic procedures or wash their hands or dipper before withdrawing the water. Thus, the study indicated that the storage and handling practices, number of family members or children in the family, socio-economic status of the family, and their hygienic practices significantly affect the quality of drinking water in the houses

References:

1. Agensi A, Tibyangye J, Tamale A, Agwu E, & C. Amongi., 2019. Contamination Potentials of Household Water Handling and Storage Practices in Kirundo Subcounty, Kisoro District, Uganda. *Journal of Environmental and Public Health*, pp. 1–8.
2. Alexander A, J.Tibyangye, A.Tamale, E., and C. Amongi (2019). Contamination Potentials of Household Water Handling and Storage Practices in Kirundo Subcounty, Kisoro District, Uganda. *Journal of Environmental and Public Health* Volume 2019, Article ID 7932193, 8 pages <https://doi.org/10.1155/2019/7932193>
3. APHA Standard Methods for the Examination of Water and Wastewater 23rd Edition, 2017
4. Chalchisa D, Megersa M, Beyene A (2017) Assessment of the quality of drinking water in storage tanks and its implication on the safety of urban water supply in developing countries. *Environ Syst Res* 6:1.
5. Duru M, Amadi C, Amadi B, Nsofor C, Humphrey N (2013) Effect of different storage vessel on water quality. *Global Res J Sci* 2:9–13.

6. Gorter, A. C., Peter Sandtford and Rosa Marfa, (1998). Hygiene in rural Nicaragua in relation to diarrhea. *International J. Epidemiology*, 27: 1090-1100.
7. Jagals P, Jagals C, Bokako T (2003). The effect of container-biofilm on the microbiological quality of water used from plastic household containers. *J Water Health* 1(3):101–108.
8. Karn, K. and H. Harada, (2002). Field survey on water supply sanitation and associated health impact in urban poor communities- a case from Mumbai city India. *Water Sci. Technol.* 46(11): 265-276.
9. Manja K S, Sambasiva R, Chandra S K V, Nath K J, Dutta S, Gopal K, Lyengar L, Dhindsa SS and Parija SC (2001), “Report of study on H2S test for drinking water”, UNICEF, New Delhi.
10. Musa M., T.G. Ngobi, L. Okeny, P. Acheng, H. Namakula, E. Kyaterekera, I. Nansubuga and N. Kibwami (2021) The effect of household storage tanks/vessels and user practices on the quality of water: a systematic review of literature. *Environmental Systems Research* vol 36. Open Access. 22 Feb. 2021.
11. Nala, N.P., P Jagals and G. Joubert (2003). The effect of a water hygiene educational program on the microbiological quality of container stored water in household. *Water SA*. 29(2), 171-176. <http://www.int.water/san/>
12. Packiyam R. , S. Kananan, S. Pachaiyappan and U. Narayanan (2016) Effect of Storage Containers on Coliforms in Household Drinking Water. *Int.J.Curr. Microbiol.App.Sci* (2016) 5(1): 461-477.
13. Tambekar D.H. and A.J. Mahore. Studies on personal and domestic hygienic practices on storage and handling of drinking water in houses. *Pollution Research*, 2005: 24 (2): 377-381
14. Tambekar DH , SD Tambekar, Dhote SV and Solanke RS . Drinking water quality deterioration in hostels in Amravati city. *Indian Journal of Applied Research*, 2017: 7(4):188-190.
15. Tambekar DH, Dhote SV and Shinde GM, 2016. Household drinking water: Assessment of microbiological contamination between source and point of use. *Bioscience Discovery*, 7(2):147-157.
16. World Health Organisation (WHO), 2005. Revised drinking water guidelines to help prevent water related outbreaks and disease, *Public Health News*.
17. Yvan Hutin and Stephen Luby, (2003). A large outbreak in Kona city, Nigeria: the importance of hand washing with soap and the danger of street-vended water. *J. Water and Health*. 1 (1), 45-52.

