

Preliminary Screening of AminoAcids from a Medicinal Plant - *Embelia basal*

Dr. Gayatri S. Kamble¹

Assistant Professor

¹Department of First Year Engineering,
Anantrao Pawar College of Engineering and Research, Parvati, Pune – 411009, Maharashtra, India

¹gayatri.kamble@abmspcorpune.org

Abstract: *Embelia basal*, a shrub from family *Myrsinaceae* have various medicinal values. Detection of amino acids from various extracts of leaves of *Embelia basal* was carried out by using different mobile phases with paper chromatographic technique. The results obtained were compared with standard amino acids in respective mobile phases. Proline and serine were found to be present in all different phases examined for leaves of *Embelia basal*. Total 19 amino acids were detected from different extracts of the plant. The maximum separation and detection of amino acids was observed in two specific mobile phases

Index Terms - *Embelia basal*, Paper chromatography, Amino acids.

I. INTRODUCTION

Natural products are used as traditional medicines from ancient times. They are having a great importance in Ayurveda. The present study deals with the leaves of *E. basal* that is Vidanga, a well known Ayurvedic medicine. Various species of *Embelia* are used in Ayurveda for therapeutic purposes (1). In Ayurveda Vidanga is used to cure pyorrhea, diarrhea and flatulence, for the treatment of cough and as an antifertility drug. Fresh juice is cooling, diuretic and laxative. Fruits are acrid, light, astringent, carminative, anthelmintic, stimulant, alternative. Its reputation is due to the action of expelling tapeworms. Fruits cure dental, oral and throat troubles. A paste of seeds is applied locally against ringworm and seed powder is used as an errhine in cold and headache. A decoction of seeds is beneficial in fever, skin diseases and chest complaints (2).

Literature survey revealed wide biological activity of family *Myrsinaceae*. One of the species, *Embelia ribes* is used in dental caries (3). It is also used in the process of formulating Anti-AIDS Ayurvedic pharmaceutical compositions (4). The antibacterial activity of embelin isolated from berries of *Embelia ribes* has been reported (5). The most of the work has been carried out on *Embelia ribes*. This species shows an antispermatogenic effect (6). It also acts as a contraceptive (7). This family is widely spread and has applications from antediluvian days. Taking into consideration such wide biological activity, the qualitative analysis of amino acids is performed.

E. basal having similar botanical characters are always mixed with or replaced for *Embelia ribes*, even though the therapeutic property of the formulation does not change. This suggests equal importance of *Embelia basal*. Fruits of *Embelia basal* were also composed of the essential amino acids (8). Therefore, *Embelia ribes* a medicinally important plant with wide biological activities attracted us to study the similar species, *Embelia basal*.

II. EXPERIMENTAL

The leaves of *Embelia basal* (R & S) A. Dc. family *Myrsinaceae* were authenticated by Agharkar Research Institute, Pune, India. Whatmann filter paper no. 1 was used for paper chromatography. Dried and pulverized material of leaves was used to prepared extracts. These extracts of definite concentrations were prepared by using chloroform, acetone, ethanol and water. These extracts were repeatedly treated with chloroform for the removal of chlorophyll. The chloroform layer was separated and the remaining part was used for amino acid analysis.

The following three mobile phases were used in paper chromatographic technique for amino acid estimation.

Phase 1: Pyridine:Isoprophyllalcohol: Aceticacid:Water (8:8:1:3)

Phase 2: *n*-Butanol: Aceticacid:Water (4:1:5)

Phase 3: *n*-Butanol:Pyridine:Aceticacid:Water (6:10:1:3)

The chromatographic paper was dried and the spots were developed using ninhydrin as a spraying reagent. The different extracts showed the presence of various amino acids in different mobile phases as reported in Tables 1-3.

III. RESULTS AND DISCUSSION

The amino acids are basic units of proteins and therefore their presence was detected. Similar to Fruits, leaves of *Embelia basal* were also found to be a rich source of various amino acids. Qualitative estimation of amino acids by paper chromatography showed the presence of proline and serine in all the phases. Amino acids like butyric acid, valine, glutamic acid and methionine are found to be present commonly in phases 1 and 2; while lysine, ornithine, iso-leucine and phenyl alanine were detected in phases 3. Alanine was detected in phase 1, Threonine was detected in phase 2 and remaining amino acids like aspartic acid, histidine, hydroxyproline and L-dopa were detected in phase 3.

**AMINO ACIDS DETECTED IN THE PHASE 1:
PYRIDINE: ISOPROPHYL ALCOHOL: ACETIC ACID: WATER (8:8:1:3)**

Name of amino acids	R _f for standard amino acids	R _f for sample
Alanine	0.330	0.320
Ornithine	0.100	0.100
Proline	0.410	0.400
Butyric acid	0.460	0.460
Valine	0.530	0.540
Glutamic acid	0.220	0.210
Iso-Leucine	0.640	0.640
Methionine	0.550	0.550
Phenyl alanine	0.600	0.610
Serine	0.210	0.200

**AMINO ACIDS DETECTED IN THE PHASE 2:
N-BUTANOL: ACETIC ACID: WATER (4:1:5)**

Name of amino acids	R _f for standard amino acids	R _f for sample
Proline	0.360	0.360
Threonine	0.300	0.290
Butyric acid	0.430	0.430
Valine	0.530	0.520
Glutamic acid	0.310	0.300
Methionine	0.550	0.550
Serine	0.250	0.260
Tryptophan	0.590	0.580

**AMINO ACIDS DETECTED IN THE PHASE 3:
N-BUTANOL: PYRIDINE: ACETIC ACID: WATER (6:10:1:3)**

Name of amino acids	R _f for standard amino acids	R _f for sample
Cysteine	0.440	0.420
Lysine	0.100	0.100
Ornithine	0.080	0.070
Proline	0.330	0.330
Threonine	0.280	0.290
Aspartic acid	0.500	0.500
Histidine	0.130	0.130
Iso-Leucine	0.580	0.590
Phenyl alanine	0.580	0.580
Serine	0.190	0.210
Tryptophan	0.530	0.530
Hydroxyproline	0.230	0.230
L-Dopa	0.430	0.420

IV. ACKNOWLEDGEMENTS

Author is thankful to the BCUD-Savitribai Phule Pune University for the funding. Author is also thankful to the Principal and the Head, Department of First Year Engineering, ABMSP's Anantrao Pawar College of Engineering and Research Pune, India for providing the necessary laboratory facilities for the work.

REFERENCES

- [1] V.S. Agarwal, Drug Plant of India, Kalyani Publishers, Ludhiana, Vol. 1, p. 353 (1997).
- [2] S.G. Joshi, Medicinal Plants, Oxford and IBH Publishing Co. Pvt. Ltd. New Delhi, p. 289 (2000).
- [3] T. Namba, M. Tsunozuka, K. Saito and U. Pilapitiya, Studies on Dental Caries Prevention by Traditional Medicines (Part-VII), Vol. 39(2), pp. 146-153 (1985).
- [4] S. Khanna, A Process of Preparing Anti-AIDS, GB Patent, GB2314270 (1998).
- [5] M. Chitra, C.S.S. Devi and E. Sukumar, Fitoterapia, 74, 401 (2003).
- [6] S.D. Seth, N. Johri and K.R. Sundaram, Indian J. Pharmacol., 14, 207 (1982).
- [7] T.K. Chatterjee, Herbal Options, Books and Allied (P) Ltd., Kolkata, pp. 98-99 (2000).
- [8] G.S. Kamble, S.M.Devare, S.M.Shaha, N.R.Deshpande, J.P.Salvekar, Asian J. of Chemistry, Vol. 21, No. 6(2008), 4429-4432, 2008.

