

# PSEUDO $\tilde{M}$ -PROJECTIVE CURVATURE TENSOR ON A RIEMANNIAN MANIFOLD

<sup>1</sup>Suraj Prakash Mishra, <sup>2</sup>Bhagwat Prasad, and <sup>3</sup>Virendra Upadhyaya

<sup>1</sup>Department of Physical Sciences MGCGV, Chitrakoot, Satna, M.P.

<sup>2</sup>Department of Mathematics S.M.M. TOWN PG College, Ballia-277001

<sup>3</sup>Department of Physical Sciences MGCGV, Chitrakoot, Satna, M.P.

**Abstract:** Yano and Sawaki(1972) introduced quasi conformal curvature tensor in a Riemannian manifold. Recently one of the author Prasad(2002) investigated pseudo projective curvature tensor in a Riemannian manifold. In this paper we introduced a new curvature tensor named as Pseudo  $\tilde{M}$ -projective curvature tensor on a Riemannian manifold. Some properties for Pseudo  $\tilde{M}$ -projective curvature tensor are investigated. Finally a particular case has been shown.

**Key words and phrases:** Conformal curvature tensor C, projective curvature tensor P, conharmonic curvature tensor H, concircular curvature tensor V, quasi conformal curvature tensor  $\tilde{C}$ , pseudo projective curvature tensor  $\tilde{P}$ , quasi concircular curvature tensor  $\tilde{V}$ , M-projective curvature tensor M, pseudo M-projective curvature  $\tilde{M}$  and  $W_i$  – curvature tensor ( $i=1, 2, \dots, 9$ ).

## 1. Introduction

The Weyl conformal curvature tensor (Eisenhart 1949) is the well known tensor, which is invariant of every conformal transformation. In particular, if a conformal transformation transforms a harmonic function into a harmonic function, then it is called a conharmonic transformation, and the conharmonic curvature tensor (Ishii, 1957) is an invariant under conharmonic transformations. Next it is well known that a semi-Riemannian manifold is locally projectively flat if and only if the projective curvature vanishes. Apart from this the concircular curvature tensor (Yano, 1940) is the next most important curvature tensor from the semi – Riemannian point of view, which is an invariant of concircular transformations.

Let conformal curvature tensor C, projective curvature tensor P, conharmonic curvature tensor H and concircular curvature tensor V are given by (Mishra, 1984) .

$$C(X,Y)Z = R(X,Y)Z - \frac{1}{n-1} [S(Y,Z)X - S(X,Z)Y + g(Y,Z)QX - g(X,Z)QY] + \frac{r}{(n-1)(n-2)} [g(Y,Z)X - g(X,Z)Y],$$

where R is the curvature tensor, S is the Ricci tensor and r is the scalar curvature,

$$C(X,Y,Z,W) = g(C(X,Y)Z,W)$$

$$R(X,Y,Z,W) = g(R(X,Y)Z,W)$$

$$\text{and } S(X,Y) = g(QX, Y).$$

$$P(X,Y)Z = R(X,Y)Z - \frac{1}{n-1} [S(Y,Z)X - S(X,Z)Y]$$

$$H(X,Y)Z = R(X,Y)Z - \frac{1}{n-2} [S(Y,Z)X - S(X,Z)Y + g(Y,Z)QX - g(X,Z)QY] \text{ and}$$

$$V(X,Y)Z = R(X,Y)Z - \frac{r}{n(n-1)} [g(Y,Z)X - g(X,Z)Y]$$

These curvatures satisfies the symmetric and skew symmetric property as well as cyclic property possessed by the curvature tensor  $R(X,Y)Z$ .

In 1970-72 and 1982 Mishra and Pokhariyal defined the following curvature tensors on Riemannian manifolds

$$W_1(X,Y)Z = R(X,Y)Z + \frac{1}{n-1} [S(Y,Z)X - S(X,Z)Y]$$

$$W_2(X,Y)Z = R(X,Y)Z + \frac{1}{n-1} [g(X,Z)QY - g(Y,Z)QX]$$

$$W_3(X,Y)Z = R(X,Y)Z + \frac{1}{n-1} [g(Y,Z)QX - g(X,Z)QY]$$

$$W_4(X,Y)Z = R(X,Y)Z + \frac{1}{n-1} [g(X,Z)QY - g(X,Y)QZ]$$

$$W_5(X,Y)Z=R(X,Y)Z+ \frac{1}{n-1} [g(X,Z)QY - S(X,Z)Y]$$

$$W_6(X,Y)Z=R(X,Y,Z) + \frac{1}{n-1} [g(X,Y)QZ - S(Y,Z)X]$$

$$W_7(X,Y)Z=R(X,Y)Z + \frac{1}{n-1} [g(Y,Z)QX - S(Y,Z)X]$$

$$W_8(X,Y)Z=R(X,Y)Z+ \frac{1}{n-1} [S(X,Y)Z - S(Y,Z)X]$$

$$W_9(X,Y)Z=R(X,Y)Z + \frac{1}{n-1} [S(X,Y)Z - g(Y,Z)QY]$$

Some properties of these curvature tensor have been studied by many authors such as (De and Ghosh,1994,1995and 1996), (De and Yieldiz, 2010), (Pokhariyal, 1982), (Prasad, 1997, 2002 and 2003), (Prakash, 2010), (Pokharyal etal,2017) and many others.

Yano and Sawaki in 1972 introduced quasi conformal curvature tensor in Riemannian manifold as follows.

$$C(X,Y)Z= aR(X,Y)Z+b[S(Y,Z)X-S(X,Z)Y+g(Y,Z)QX-g(X,Z)QY]- \frac{r}{n} \left( \frac{a}{n-1} + 2b \right) . [g(Y,Z)X - g(X,Z)Y],$$

where a and b are constants such that  $a \neq 0, b \neq 0$

Some properties quasi conformal curvature tensor have been studied by (Amun and Maralabhavi, 1977), (Chaki and Ghosh, 1997), (De and Shaikh, 1997), (Kumar Prasad and Verma, 2009), (Prakash and Singh, 2009), (Mallick, zhau al De, 2016),(Shukla and Singh, 2011), (Venkatesh and Kumar, 2016) and many other workers.

Recently one of the author (Prasad, 2002) introduced pseudo projective curvature tensor in a Riemannian manifold ( $n > 2$ ) by

$$\tilde{P}(X,Y)Z = a R(X,Y)Z+b[S(Y,Z)X-S(X,Z)Y] - \frac{r}{n} \left( \frac{a}{n-1} + b \right) . [g(Y,Z)X - g(X,Z)Y]$$

where a and b are constants such that  $a, b \neq 0$ .

In recent papers (Bagewadi, Prakash and Venkatesh, 2007), (Narain, Prakash and Prasad, 2009), (Jaisaval and Ojha, 2010), (Guler, Demirabag, 2016), (Kumar, 2019) etc explored various geometrical properties by using this curvature tensor on LP – Sasakian manifold, K-contact manifold and Trans-Sasakian manifold, weakly symmetric manifold and Riemannian manifold.

Further in 2004, 2007 and 2008 (Prasad & Maurya) defined pseudo  $W_2$  – curvature tensor, quasi – concircular curvature tensor and pseudo  $W_4$  – curvature tensor on Riemannian manifold as follows respectively

$$\tilde{W}_2(X,Y)Z=a R(X,Y,Z) + b [g(Y,Z)QX-g(X,Z)QY]- \frac{r}{n} \left( \frac{a}{n-1} + b \right) . [g(Y,Z)X - g(X,Z)Y],$$

$$\tilde{V}(X,Y)Z=aR(X,Y)Z+\frac{r}{n} \left( \frac{a}{n-1} + 2b \right) . [g(Y,Z)X - g(X,Z)Y],$$

and

$$\tilde{W}_4(X,Y)Z=a R(X,Y)Z + b [g(X,Y)QZ-g(X,Z)QY]- \frac{r}{n} \left( \frac{a}{n-1} + b \right) . [g(X,Y)Z - g(X,Z)Y],$$

(Narain, Prakash and Prasad, 2009), (Kumar, Prasad and Verma 2009), (Prakash, 2011), (Kumar , 2012) and (Prasad etal,2010) extended this notation to LP- Sasakian manifold, P-Sasakian manifold and LP- Sasakian manifold with coefficient  $\alpha$ .

In continuation of above study, the object of this paper is to define another a new curvature tensor on a Riemannian manifold which we call it Pseudo  $\tilde{M}$ -projective curvature tensor. The paper is organized as follows. In section 2, the definition of Pseudo  $\tilde{M}$ -projective curvature tensor is given. We give properties and some identities of this curvature tensor.

In section 3, it is proved that pseudo  $\tilde{M}$ -projectively flat manifold is of zero scalar curvature provided  $a(n+1)+bn(3n+2)+ c(n-1) \neq 0$

In the last section, a necessary and sufficient condition for  $(M^n, g)$  to be Pseudo  $\tilde{M}$ -projectively conservative is obtained.

**2. Pseudo  $\tilde{M}$ -projective curvature tensor:**

Let  $(M^n, g)$  be an n-dimensional Riemannian manifold and  $\chi(M^n)$  the Lie algebra of vector fields in  $M^n$ . Throughout the paper we assume that  $X, Y, Z, W \in \chi(M^n)$ , unless specifically stated otherwise.

In an n-dimensional Riemannian manifold  $(M^n, g)$ , pseudo  $\tilde{M}$ -projective curvature tensor of type (1,3) is defined by

$$\tilde{M}(X, Y)Z = aR(X, Y)Z + b[S(Y, Z)X - S(X, Z)Y] + c[g(Y, Z)QX - g(X, Z)QY] + \frac{r}{n} \left( \frac{a}{n-1} + 3b - c \right) \cdot [g(Y, Z)X - g(X, Z)Y], \tag{2.1}$$

where a, b and c are constants such that a, b, c ≠ 0 and R, S, Q and r are the curvature tensor, the Ricci tensor, the Ricci operator of the type(1,1) and the scalar curvature respectively.

If a=1, b=c=  $-\frac{1}{2(n-1)}$ , then (2.1) takes the form

$$\tilde{M}(X, Y)Z = R(X, Y)Z - \frac{1}{2(n-1)} [S(Y, Z)X - S(X, Z)Y + g(Y, Z)QX - g(X, Z)QY] = M(X, Y)Z,$$

where  $M(X, Y)Z$  is the M-Projective curvature tensor (Pokhariyal, 1982). Thus the M-projective curvature tensor is a particular case of the tensor  $\tilde{M}(X, Y)Z$ . For this reason  $\tilde{M}$  is called the  $\tilde{M}$ -Projective curvature tensor.

Note that Pseudo  $\tilde{M}$ -projective curvature tensor satisfies the following algebraic properties:

- (i)  $\tilde{M}(X, Y, Z, W) + \tilde{M}(Y, X, Z, W) = 0$
- (ii)  $\tilde{M}(X, Y, Z, W) + \tilde{M}(X, Y, W, Z) = (b-c)[S(Y, Z)g(X, W) - S(X, Z)g(Y, W) + S(Y, W)g(X, Z) - S(X, W)g(Y, Z)]$
- (iii)  $\tilde{M}(X, Y, Z, W) + \tilde{M}(Z, W, X, Y) = (b+c)[S(Y, Z)g(X, W) + g(Y, Z)S(X, W)] - 2[bS(X, Z)g(Y, W) + cS(W, Y)g(Z, X)]$ .
- (iv)  $\tilde{M}(X, Y, Z, W) + \tilde{M}(Y, Z, X, W) + \tilde{M}(Z, X, Y, W) = 0$ ,  
 where  $\tilde{M}(X, Y, Z, W) = g(M(X, Y)Z, W)$  and  $R(X, Y, Z, W) = g(R(X, Y)Z, W)$ .

Let  $\{e_i\}$  be an orthonormal basis of the tangent space at each point of the manifold where  $1 \leq i \leq n$ . Now from (2.1), we have

$$\sum_{i=1}^n \tilde{M}(X, Y, e_i, e_i) = \sum_{i=1}^n \tilde{M}(e_i, e_i, Z, W) = 0$$

Particular case:

If the manifold becomes an Einstein manifold then from (2.1), we get

$$M(X, Y)Z = aR(X, Y)Z + \frac{r}{n} \left[ \frac{a}{n-1} + 4b \right] [g(Y, Z)X - g(X, Z)Y] = \overset{\star}{V}(X, Y)Z \tag{2.2}$$

(a) new curvature tensor

If a=1, b=  $-\frac{1}{2(n-1)}$  then (2.2) taken the form  $\overset{\star}{V}(X, Y)Z = R(X, Y)Z - \frac{r}{n(n-1)} [g(Y, Z)X - g(X, Z)Y] = V(X, Y)Z$ ,

where  $V(X, Y)Z$  is the concircular curvature tensor. Thus we observe that the

concircular curvature  $V$  is a special case of the tensor  $\overset{\star}{V}(X, Y)Z$ , which seems to be the generalization of concircular curvature tensor.

Hence we call it as generalized concircular curvature tensor  $\overset{\star}{V}$ .

**3. Pseudo  $\tilde{M}$ -projectively flat manifold.**

For pseudo  $\tilde{M}$ -projectively flat manifold, we have

$$\tilde{M}(X, Y)Z = 0 \tag{3.1}$$

Then from (2.1) and (3.1) we obtain

$$aR(X, Y)Z + b[S(Y, Z)X - S(X, Z)Y] + c[g(Y, Z)QX - g(X, Z)QY] + \frac{r}{n} \left[ \frac{a}{n-1} + 3b - c \right] \cdot [g(Y, Z)X - g(X, Z)Y] = 0 \tag{3.2}$$

From (3.2), we get

$$aR(X, Y, Z, W) + b[S(Y, Z)g(X, W) - S(X, Z)g(Y, W)] + c[g(Y, Z)S(X, W) -$$

$$g(X,Z)S(Y,W)] + \frac{r}{n} \left[ \frac{a}{n-1} + 3b - c \right] [g(Y,Z)g(X,W) - g(X,Z)g(Y,W)] = 0 \tag{3.3}$$

Putting  $X=W=e_i$  in (3.3),  $\{e_i\}$  is an orthonormal basis of the tangent space at a point of the manifold and taking summation over  $e_i, 1 \leq i \leq n$ , we get

$$S(Y,Z)[c - a + b(n - 1)] = \frac{r}{n} [a + 3b(n - 1) + c]g(Y,Z),$$

which gives on further contraction

$$r[a(n+1) + b(3n+2)n + c(n-1)] = 0 \tag{3.4}$$

If  $a(n+1) + b(3n+2)n + c(n-1) \neq 0$ , then from (3.4) we get  $r=0$ .

Hence, we have the following theorem:

**Theorem (3.1):** Pseudo  $\tilde{M}$ -projectively flat manifold is of zero scalar curvature, provided  $a(n+1) + b(3n+2)n + c(n-1) \neq 0$ .

**4. Pseudo  $\tilde{M}$ -projectively conservative ( $M^n, g$ ).**

A manifold  $(M^n, g)$  shall be called pseudo  $\tilde{M}$ -projectively conservative if

$$\text{div } \tilde{M} = 0 \text{ (Hicks, 1969)} \tag{4.1}$$

Now differentiating (2.1) covariantly, we get

$$\begin{aligned} (D_U R)(X, Y)Z + b[(D_U \tilde{M})(X, Y)Z] &= a(D_U R)(X, Y)Z + b[(D_U S)(Y, Z)X - (D_U S)(X, Z)Y] + \\ & c[g(Y, Z)(D_U Q)(X) - g(X, Z)(D_U Q)(Y)] - \\ & \frac{D_U r}{n} \left[ \frac{a}{n-1} + 3b - c \right] \cdot [g(Y, Z)X - g(X, Z)Y], \end{aligned}$$

$$\begin{aligned} (\text{div } \tilde{M})(X, Y)Z &= a(\text{div } R)(X, Y)Z + b[(D_X S)(Y, Z) - (D_Y S)(X, Z)] + \\ & c[g(Y, Z)dr(X) - g(X, Z)dr(Y)] - \frac{1}{n} \left[ \frac{a}{n-1} + 3b - c \right] [g(Y, Z)dr(X) - g(X, Z)dr(Y)] \end{aligned} \tag{4.2}$$

$$\text{From (Eisenhart, 1926), we have } (\text{div } R)(X, Y)Z = (D_X S)(Y, Z) - (D_Y S)(X, Z) \tag{4.3}$$

$$\begin{aligned} \text{In view of (4.2) and (4.3), we get } (\text{div } \tilde{M})(X, Y)Z &= (a+b)[(D_X S)(Y, Z) - \\ & (D_Y S)(X, Z)] + \left[ \frac{c(n^2-1) - (a+3b(n-1))}{n(n-1)} \right] [g(Y, Z)dr(X) - g(X, Z)dr(Y)] \end{aligned} \tag{4.4}$$

Suppose the Ricci tensor  $S(X, Y)$  is of Codazzi type i.e

$$(D_X S)(Y, Z) = (D_Y S)(X, Z) \tag{4.5}$$

Then from (4.4) and (4.5), we get

$$(\text{div } \tilde{M})(X, Y)Z = \left[ \frac{c(n^2-1) - (a+3b(n-1))}{n(n-1)} \right] [g(Y, Z)dr(X) - g(X, Z)dr(Y)] \tag{4.6}$$

$$\text{Hence if } (\text{div } \tilde{M})(X, Y)Z = 0, \text{ then from (4.6), } c(n^2-1) - (a+3b(n-1)) [g(Y, Z)dr(X) - g(X, Z)dr(Y)] = 0$$

Since  $c(n^2-1) - (a+3b(n-1)) \neq 0$ , here

$$g(Y, Z)dr(X) - g(X, Z)dr(Y) = 0, \text{ which shows that } r \text{ is constant. Again if } r \text{ is constant then from (4.6), we get } (\text{div } \tilde{M})(X, Y)Z = 0.$$

Hence we can state the following theorem:

**Theorem (4.1).** If in a Riemannian manifold  $(M^n, g)$ , the Ricci tensor is Codazzi type then the manifold Pseudo  $\tilde{M}$ -projectively conservative if and only if the scalar curvature is constant, provided  $c(n^2-1) - (a+3b(n-1)) \neq 0$ ,

According to our assumption  $a+b \neq 0$ , then from (4.4), we have

$$\frac{(\text{div } \tilde{M})(X, Y)Z}{a+b} = [(D_X S)(Y, Z) - (D_Y S)(X, Z)] + \left[ \frac{c(n^2-1) - (a+3b(n-1))}{n(n-1)} \right] [g(Y, Z)dr(X) - g(X, Z)dr(Y)] \tag{4.7}$$

Hence from (4.7), we can state theorem as follows

**Theorem (4.2).** If in a Riemannian manifold  $(M^n, g)$ , Pseudo  $\tilde{M}$ -projective curvature tensor is such that  $a+b \neq 0$ , the manifold is pseudo  $\tilde{M}$ -projectively conservative if and only if

$$[(D_X S)(Y, Z) - (D_Y S)(X, Z)] = \left[ \frac{(a+3b(n-1)) - ((n^2-1))}{n(n-1)} \right] \cdot [g(Y, Z)dr(X) - g(X, Z)dr(Y)].$$

## References

- [1] Yano , K and Sawaki, S,1972. Riemannian manifolds admitting a conformal transformation group, J.of Diff. Geo, 2, 161-165.
- [2] Yano, K,1940. Conircular Geometry I.Conircular transformations, Math. Institute, Tokyo Imp. Univ. Proc, 16, 195-200.
- [3] Eisenhart, L.P. 1949. Riemannian Geometry Prinuton University Press.
- [4] Ishii, Y.1957. On conharmonic transformations, Tensor .N.S, 7,73-80.
- [5] Mishra, R.S.1984. Structures on a differentiable manifold and their applications. Chandrama Prakashan, 50 A Balrampur house, Allahabad, India.
- [6] Mishra, R.S and Pokhariyal, G.P.1970, curvature tensors and their relativistic significance I. The Yokohoma Mathematical Journal, XVIII, (2),105-108.
- [7] Mishra, R.S and Pokhariyal, G.P.1971. Curvature tensors and their relativistic significance II. The Yokohama Math Jour. XIX(2), 97-103.
- [8] Pokhariyal G.P.1972. Curvature tensors and their relativistic significance III,The Yokohama Math Jour. XX(2), 115-119.
- [9] Pokhariyal, G.P.1982, Relativistic significance of curvature tensors, Internat J.Math Sci, 5(1), 133-139.
- [10] Pokhariyal, G.P.1982, study of new curvature tensor in a Saskian manifold, Tensor N.S. 36, 222-226.
- [11] De, UC, and Ghosh, J.C.1994. On Pseudo  $W_2$ - symmetric manifolds, Bull. Cal. Math. Soc 86, 521- 526.
- [12] De, UC and Ghosh, J.C.1995. On generalized pseudo  $W_2$ -symmetric manifolds, Bull. Cal. Math. Soc 87, 339- 334.
- [13] De, UC and W Ghosh, J.C.1996 On a type of Sasakian manifold, Ganita, 47(1), 51 - 55.
- [14] De UC and Yildiz, A.2010. On a type of kenmotsu manifolds, DGDS, 12, 289-398.
- [15] Prasad, B.1997.  $W_2$ -curvature tensor on Kenmotsu manifold, Indian J. Maths 39(3), 287-291.
- [16] Prasad, B.2002. On semi-pseudo  $W_4$ -symmetric manifold, Ganita, 53(2), 185-191.
- [17] Prasad, B.2002. On pseudo projective curvature tensor on a Riemannian manifold, Bull. Cal. Math. Soc, 94, 163-166.
- [18] Prasad, B.2003. On generalized  $W_4$ -recurrent manifolds, Tensor N.S. 64.
- [19] Prasad, B. and Maurya 2007. Quasi con circular curvature tensor on a Riemannian manifold, News Bull. Col. Math. Soc. 30(1-3), 5-6.
- [20] Prakasha, D.G. 2010. On generalized  $W_2$ -recurrent  $(LCS)_n$ -manifold, JTS, 4, 33-40.
- [21] Njori, P.W., Moindi, S.k and Pokhariyal, G.P.2017. A study on  $W_8$ -curvature tensor in a Sasakian manifolds, Pioneer J. of Math and Mathematical Sc., 20(1), 1-11.
- [22] Amur, K and Maralabhavi, Y.B.1977. On quasi- conformally space, Tensor N.S.31, 194-197.
- [23] Chaki, M.C. and Ghosh, M.L.1997. On quasi- conformally and quasi-conform ally conservative Riemannian manifold, AL.I.CUZA. Tomnl XLIII, 375-38.
- [24] De, U.C. and Shaikh A.A.1997. K-contact and Sasakian manifolds with conservative quasi-conformal curvature tensor, Bull. Cal. Math. Soc.89, 349-354.
- [25] Kumar, R, Prasad, B and Verma, S.K.2009. On P-Sasakian manifolds satisfying certain conditions on the quasi-conformal and quasi-concircular curvature tensor, Rev. Bull. Call. Math. Soc., 17(1&2), 85-90.
- [26] Prakash, A. and Singh, A.2009.Quasi-conformal curvature tensor on LP-Sasakian manifold. JTS, 3, 59-70.
- [27] Shukla, S.S. and Singh, D.D.2011. On conformal curvature tensor of (E)- para-Sasakian manifolds, SUT Journal of Maths, 47(2), 119-128.
- [28] Mallick, S.Zhao, P. and De, U.C.2016. Spacetimes admitting quasi-conformal curvature tensor, Iranian Mathematical Society, 42(6).1535-1546.
- [29] Venkatesha and Kumar, R.T.N.2016. Quasi-conformal curvature tensor  $N(K)$ -contact metric manifolds, Acta. Math. Univ. Comeniance,LXXXV (1), 125-135.

- [30] Bagewadi, C.S., Prakash, D.G. and Venkatesha 2007. On pseudo protectively flat LP-Sasakian manifold with coefficient  $\alpha$ , Anna. Univ. Curic-Sklodowska Lubin-Polonia, IXI, 1-8.
- [31] Narain, D, Prakash, A and Prasad, B. 2009. A pseudo projective curvature tensor on LP-Sasakian manifold, AL.I CUZA, "Tomnl LV-f-2, 275-284.
- [32] Jaisawal,J.P. and Ojha, R.H.2010. On weakly pseudo-protectively symmetric manifold, DGDS, 12, 83-94.
- [33] Guler , S. and Demirbag, S.A.2016, Riemannian manifold satisfying certain conditions on pseudo projective curvature tensor, Filomat 30(3), 721-31.
- [34] Kumar, Rajesh 2019. On a type of generalized symmetric manifold, common Korean Math, Soc. 37(3), 921-934.
- [35] Prasad, B and Maurya. A.M.2004. A pseudo  $W_2$  –curvature tensor on a Riemannian manifold. J. of Pure Math, 21, 81-85.
- [36] Prasad, B., Narain, S. and Maurya, A. 2008. A pseudo  $W_4$ - curvature tensor on a Riemannian manifold, Act a ciencia Indica Math., XXXIV 357-360.
- [37] Narain, D., Prakash, A.and Prasad, B. 2009.Quasi-concircular curvature tensor on LP-Sasakian manifold, Bull. Call. Math. Soc.,101, 387-394.
- [38] Prakash, A. 2011, $\emptyset$ -pseudo  $\tilde{W}_4$ -flat LP-Sasakian manifolds. The Journal of Mathematics and Comp. Sci, 3(3), 301-305.
- [39] Kumar, Rajesh 2012. en pseudo  $\tilde{W}_2$ -flat LP-Sasakian manifold with coefficient  $\alpha$ , Science Vision, 12(1), 17-21.
- [40] Prasad, B., Gupta, N.and Pandey, S.N. 2010. Pseudo  $\tilde{W}_2$ -curvature tensor on a L.P.-Sasakian manifold, Journal of progressive science 1(2).
- [41] Jaeman Kim 2016.On pseudo  $W_4$ -symmetric manifolds, Honam Math. J.38 (1), 39-45.
- [42] Hicks, N.J. 1969.Notes on differential geometry, D. Van. Nost. Camp.Inc. Princetor, New York.

