



# A REVIEW ON MOUTH DISSOLVING TABLET

Sachin Munde<sup>1</sup>, Avinash Dhoble<sup>2</sup>, Santosh Waghmare<sup>3</sup>

Department of Pharmaceutics, Loknete Shri Dadapatil Pharate College of Pharmacy Mandavgan Pharata. Tal- Shirur, Dist- Pune.

## ABSTRACT

The interest for MDT (Mouth Disintegrating Tablet) has been extending from the most recent ten years particularly in geriatric, pediatric and patient with inadequacies or something to that affect in swallowing... MDTs are those tablets which when set in mouth get broken up quickly in spit without the need of fluid and can be gulped. European pharmacopeia embraced the term or dispersible tablet for MDTs. Mouth breaking down tablets are otherwise called Fast softening tablets, Or dispersible tablets, quick dissolving/scattering tablets or liquefy in mouth tablets. This article surveys the potential advantages presented by MDTs as an oral medication conveyance framework for different sorts of patients experiencing various illnesses and incapacities. Wanted attributes and difficulties for growing quick deteriorating drug conveyance frameworks, quality control tests, and different strategies utilized in the readiness of quick crumbling drug conveyance frameworks like Lyophilization advancements, tablet forming strategy, sublimation procedures, splash drying strategies, mass expulsion innovation, direct pressure technique and employments of super-breaks down. It likewise surveys the licensed advancements for quick dissolving tablets, benefits and weaknesses of various innovations for planning quick deteriorating dose structure, future imminent for MDTs. The developing significance for MDTs is because of the potential benefits presented by this innovation. MDT is a New Drug Delivery framework with least breaking down time and simplicity of self-organization.

**Keywords:** MDT, Orodispersible tablets, fast dissolving/dispersing tablets, Melt in mouth tablets, Mass extrusion, Superdisintegrants

## INTRODUCTION

Numerous patients, particularly older think that it is troublesome in gulping tablets, containers, liquids and in this manner don't conform to solution, which brings about high frequency of rebelliousness situated examination has brought about bringing out numerous more secure and new medication conveyance framework. Quickly crumbling/dissolving tablet is one of such model, for the explanation of fast crumbling or even with spit. Thinking about personal satisfaction, the majority of these endeavors have been centered around simplicity of drug. Among the different measurement structures created to work on the simplicity of organization, the mouth dissolving tablet (MDT) is the most generally favored business items. The oral hole is an appealing site for the organization of medications due to simplicity of organization. Different dose structures like Tablets, Cases, and Liquid arrangements are managed by oral course. During the last decade, mouth dissolving tablet (MDT) advancements that cause tablets to deteriorate in the mouth without biting and extra water admission have drawn a incredible arrangement of consideration. The MDT is otherwise called quick dissolving, quick scattering, fast break up, quick soften, as well as speedy breaking down tablet. All MDTs supported by the Food and Drug Administration (FDA) are named orally deteriorating tablets. As of late, the European Pharmacopeia embraced the term or dispersible tablet for a tablet that scatters or on the other hand breaks down in under 3minutes in the mouth previously gulping. Such a tablet deteriorates into more modest granules or dissolves in the mouth from a hard strong to a gel-like construction, permitting simple gulping by patients. The deterioration time for great MDTs fluctuates from a few seconds to about a moment. Orally crumbling tablets give a benefit especially for pediatric and geriatric populace who experiences issues in gulping regular tablets and containers. Moreover pediatrics patients might experience the ill effects of ingestion issues as consequences of immature strong and anxious control. Also, patients going with almost no overabundance to water, limit utility of orally manage convectional tablet case. MDT brings about fast

disintegration and quick ingestion which give quick beginning of activity. Additionally, drug applicants that go through pre-gastrics ingestion when planned as MDTs may show expanded oral bioavailability. It gives great soundness exact dosing, and simple assembling [1 - 4].

### **Ideal Properties [2]**

An ideal MDT should:

- 1) Require no water for oral organization.
- 2) Have a satisfying mouth feel.
- 3) Have a satisfactory taste concealing property.
- 4) Be harder and less friable.
- 5) Leave negligible or no buildup in mouth after organization.
- 6) Exhibit low aversion to ecological conditions (Temperature and moistness).
- 7) Allow tablet producing by ordinary handling and bundling supplies.

### **Advantages [3]**

- 1) Administration to the patients who can't swallow, for example, the old, disabled patients, patients impacted by renal disappointment & patients who won't swallow, for example, pediatric, geriatric and mental patients.
- 2) Rapid medication treatment mediation.
- 3) Achieve expanded bioavailability/quick ingestion through pre-gastric ingestion of medications from mouth, pharynx and throat as salivation passes down.
- 4) Convenient for organization and patient agreeable for debilitated, out of commission patients and for explorers and occupied individuals, who don't dependably approach water.
- 5) The danger of choking or suffocation during oral organization of ordinary details due to physical impediment is kept away from, accordingly giving further developed wellbeing.
- 6) New business opportunity will get produced due to the item separation.

### **Salient Features [4]**

- 1) Ease of organization to patients who won't swallow a tablet, like pediatric and geriatric patients and, mental patients.
- 2) Convenience of organization and precise dosing as contrasted with fluids.
- 3) Rapid disintegration of medication and assimilation which may produce fast, beginning of activity.
- 4) Some medications are ingested from the pharynx and throat as the spit passes down into the stomach, in such cases bioavailability of medications is expanded.
- 5) Ability to give benefits of fluid medicine in the type of strong arrangement.
- 6) Pre-gastric ingestion can bring about further developed bioavailability furthermore because of decreased dose, worked on clinical execution through a decrease of undesirable impacts.

### **Limitations of Mouth Dissolving Tablets [4]:**

- 1) The tablets normally have lacking mechanical strength. Consequently, cautious dealing with is required.
- 2) The tablets might leave upsetting taste and additionally abrasiveness in mouth if not figured out as expected.
- 3) Drugs with generally bigger portions are hard to figure out into MDT for example anti-microbials like ciprofloxacin with grown-up portion tablet containing around 500 mg of the medication.

4) Patients who simultaneously take anticholinergic meds may not be the best possibility for MDT. Also patients with dryness of the mouth because of diminished spit creation may not be great contender for these tablet plans.

### Desired characteristics and challenges for developing fast disintegrating drug delivery systems:

- 1) Time needed for deterioration
- 2) MDTs ought to deteriorate/break down/scatter or liquefy in mouth without the need of water in extremely brief span of time, conceivably inside 60 seconds.
- 3) Taste of the dynamic fixing
- 4) As most medications are unpalatable, quick deteriorating drug conveyance frameworks as a rule contain the medicament in taste masked structure. Conveyance frameworks break up or crumble in patient's mouth, accordingly delivering the dynamic fixings which interact with the taste buds and consequently, taste concealing of the medications becomes basic to patient consistence.
- 5) Tablet strength, Friability and porosity: In request to permit quick crumbling tablets to break down in the mouth, they are made of either exceptionally permeable or delicate formed networks or compacted into tablets with extremely low pressure power, which makes the tablets friable just as powerless, which are difficult to manage, consistently requiring explicit strip off bother packaging.
- 6) Hygroscopic nature: Several quick deteriorating drug conveyance dose structures are hygroscopic and can't keep up with physical honesty under typical condition from stickiness which calls for particular item bundling.
- 7) Mouth believe: Mouth feel is basic, and patients ought to get an item that feels wonderful. Any enormous particles from the deteriorating tablet that are insoluble or gradually solvent in spit would prompt an undesirable dirty inclination. This can be defeat by keeping most of the particles underneath the recognizable size limit. Sometimes, certain flavors can soak up further developed mouth feel discernment, bringing about an item that is seen as being less abrasive, regardless of whether the main change is the flavor. Foam can be added to help crumbling and further develop mouth feel by decreasing the "dryness" of an item [5]

### Various techniques used in the preparation of mouth disintegrating drug delivery systems

- 1) Freeze–drying (Lyophilization technologies)
- 2) Tablet molding method
- 3) Sublimation techniques
- 4) Spray drying techniques
- 5) Mass extrusion technology
- 6) Direct compression method
- 7) Use of disintegrates

#### 1) Freeze drying or Lyophilization technology

A cycle by which, water get sublimated from item later freezing. Lyophilization is a drug innovation which permits drying of hotness delicate medications and biologicals at low temperature under conditions that permits expulsion of water by sublimation. Lyophilization brings about arrangements, which are profoundly permeable, with an extremely high explicit surface region, which break up quickly and show further developed retention and bioavailability [6, 7].

R. P. Scherer protected Zydis innovation by utilizing freeze drying process for the planning of mouth dissolving tablet. Based on patent gave to Gregory et al. [8] Seager talked about arrangement, process innovation and bioavailability of quick dissolving tablets arranged by utilizing Zydis innovation [8].

#### 2) Molding technique

Shaped tablets are ready by utilizing water-solvent fixings with the goal that the tablet break down or crumble quickly what's more totally. Powder is saturated with the assistance of hydro

Alcoholic dissolvable and afterward shaped into tablets under tension not exactly the ordinary measurements structure. The solvents are taken out via air-drying. The tablet Possesses permeable construction, which works with simple disintegration. Adding sucrose, acacia or PVP k30 may build the mechanical strength of tablet [6, 7].




Figure 1: Steps involved in freeze drying technology


### 3) Sublimation technique

The fundamental guideline engaged with planning quick dissolving tablets by sublimation strategy is expansion of an unpredictable salt to the tableting parts, blending the parts to acquire a Considerably homogeneous blend and volatizing an unpredictable salt. The evacuation of unpredictable salts makes pores in the tablet, which help in accomplishing fast breaking down when the tablet comes in contact with salivation. Camphor, Naphthalene, Urea, ammonium

Bicarbonate, and so on, can be utilized to get ready permeable tablets of good mechanical strength [6-7]. Koizumi et al. involved mannitol as diluent and camphor as an unstable material to get ready permeable packed tablets [9]. The tablets were exposed to vacuum at 80°C for 30 min to dispose of the camphor and in this manner structure the pores in the tablet. Makino et. All used water as a pore framing material to plan permeable tablets with incredible mechanical strength and disintegration character [6-7, 9]

Drug + volatilizing agents + other ingredients


COMPRESSION



Volatilizing agent

Compressed tablet

SUBLIMATION



Pores appeared on sublimation of volatilizing solvent

#### 4) Spray drying technique

Shower drying is an interaction by which profoundly permeable, fine powders can be delivered. Splash dryers are perpetually utilized in the drug industry to deliver exceptionally permeable powders. Allen et al. have revealed applying this interaction to the creation of quick dissolving tablets [10]. Splash Drying can be used to get ready quickly dissolving tablet. This method is

In light of a particulate help grid that is ready by splash drying and fluid structure containing support lattice and different parts to frame an exceptionally permeable and fine powder. This is then blended in with dynamic fixing and packed into tablet. Quick dissolving tablet arranged by shower drying procedure deteriorated inside 20 seconds [6-7, 10]. Licensed advancements for mouth dissolving tablets [11]

##### 1) Zydis Technology:

Zydis definition is a novel freeze dried tablet in which medication is truly entangled or disintegrated inside the grid of quick dissolving transporter material. When zydis units are placed into the mouth, the freeze-dried design crumbles quickly and doesn't expect water to help gulping. The zydis framework is made out of numerous materials to accomplish various targets. To confer strength and strength during taking care of, polymers like gelatin, dextran or alginates are consolidated. These structure a lustrous undefined structure, which bestows strength. To acquire crystallinity, class and hardness saccharides like mannitol or sorbitol are consolidated. Water is utilized in the assembling system to guarantee creation of permeable units to accomplish quick deterioration while different gums are utilized to forestall sedimentation of scattered medication particles in the assembling process. Breakdown protectants, for example, glycine forestalls the shrinkage of zydis units during freeze drying process or long-term stockpiling. Zydis items are stuffed in rankle packs to shield the detailing from dampness in the climate [11].




Figure No. 3 Steps Involved spray drying method

**2) Durasolv Technology:**

Durasolv is the licensed innovation of CIMA labs. The tablets made by this advancement contain of medication, filler and an ointment. Tablets are ready by utilizing ordinary tableting hardware and have good rigidity. These can be bundled into ordinary bundling framework like rankles. Durasolv is a fitting innovation for item requiring low measures of dynamic fixings [11]

**3) Orasolv Technology:**

CIMA labs have made Orasolv. In this framework dynamic medicament is taste veiled. It likewise contains bubbly crumbling specialist. Tablets are made by direct pressure procedure at low pressure power to limit oral disintegration time. Ordinary blenders and tablet machine is utilized to deliver the tablets. The tablets delivered are delicate and friable [11].

**4) Flash Dose Technology:**

Flash portion innovation has been protected by fuisz. Nurofen meltlet, one more kind of ibuprofen as dissolve in mouth tablets arranged utilizing streak portion innovation is the primary business item sent off by Biovail Corporation. Streak portion tablets comprise of self-binding shear structure grid named as "floss". Shear structure networks are arranged by WOW signifies "Without Water". In this interaction, blend of low mouldability saccharides and high mouldability saccharides is used to acquire a quickly liquefying solid tablet. The dynamic Fixing is blended in with a low mouldability saccharide (for example lactose, glucose, and mannitol) and granulated with a high mouldability saccharide (for example Maltose, oligosaccharides) [11]

**5) Flash tab Technology:**

Prographarm labs have protected the Flash tab innovation. Tablet arranged by this Framework comprises of a functioning fixing as miniature precious stones. Drug miniature granules might be ready by utilizing the regular strategies like coacervation, miniature exemplification furthermore expulsion spheronisation. All the handling used ordinary tablet innovation [11]

## Super Disintegrants Used in MDTs

As day's passes, interest for quicker breaking down definition is expanded. Along these lines, drug specialist needs to form disintegrants for example Super disintegrants which are powerful at low focus also have more noteworthy deteriorating effectiveness and they are more viable intragranularly. This super disintegrants act by expanding and because of enlarging pressure applied in the external course or spiral bearing, it makes tablet burst or the sped up ingestion of water prompting a huge expansion in the volume of granules to advance crumbling[12].

Different kinds of Super disintegrants utilized are as per the following -

- 1) Crosspovidone
- 2) Microcrystalline cellulose
- 3) Sodium starch glycollate
- 4) Sodium carboxy methyl cellulose/Cross carmelose sodium
- 5) Crosscarmellose sodium
- 6) Calcium carboxy methyl cellulose
- 7) Modified corn starch
- 8) Kyron

## Factors to be considered for selection of super disintegrants [12]

1. It should deliver mouth dissolving when tablet meets salivation in the mouth
2. It ought to be adequately compactable to create less friable tablets. It can ready to create great mouth feel to the patient. Along these lines, little molecule size is liked to accomplish patient Consistence.
3. It ought to have great stream since it work on the flowability of the absolute mix.

## Evaluation of mouth dissolving tablets [13, 14]

MDTs formulations have to be evaluated for the following evaluation test:

- 1) General Appearance:** The outward presentation of tablets incorporates size, shape, shading, taste, scent, and surface.
- 2) Size, Shape, Thickness and width:** The size and shape of the tablet can be correspondingly depicted, observed furthermore controlled. Thickness of tablets is a significant trademark for appearance and furthermore in counting by utilizing filling gear. Some filling gear uses the uniform thickness of the tablets as an including component. Ten tablets ought to be taken and their thickness is required to gauge by vernier caliper.
- 3) Uniformity of weight:** In Indian pharmacopeia method for consistency of weight was followed, ten or twenty tablets were taken and their still up in the air separately and by and large on an advanced gauging balance. Then, at that point, the normal load of one tablet is needed to be determining from aggregate weight. The weight variety test would be an agreeable technique for deciding the drug content consistency.
- 4) Hardness of tablets:** Hardness of tablet is characterized as the power applied across the width of the tablet in the request to break the tablet. Opposition of the tablet to chipping, scraped area or breakage under state of capacity change and taking care of before utilization relies upon its hardness. Hardness of the tablet of every plan was decided utilizing Monsanto hardness analyzer.
- 5) Friability of tablets:** Friabilator comprise of plastic chamber spins at 25 rpm, dropping those tablets at distance of 6 crawls with every insurgency. The tablets were turned in the friabilator for somewhere around 4 min. Toward the finish of these test tablets are needed to be dedusted and rechecked, the misfortune in the heaviness of tablet is the deliberate of friability and is communicated in rate as

$$\% \text{friability} = \frac{\text{initial weight} - \text{final weight}}{\text{Initial weight}} \times 100$$

**6) Disintegration time:** As depicted in pharmacopeia, tablets are set in the crumbling cylinder and time is noted. As indicated by the European pharmacopeia the quick deterioration or or dispersible tablets ought to break down inside 3 min without leaving any buildup on the screen.

**7) In-vitro dispersion time test:** To decide scattering time take a 10ml of estimating chamber and pour a 6ml of distil water in it, then, at that point, drop a tablet in the equivalent. At last the time needed for complete not set in stone as a scattering time.

**8) Wetting time:** Take five roundabout tissue papers of 10 cm measurement and put them in a petridish with a 10 cm distance across. Ten millimeters of water containing Eosin, a water-solvent color, is needed to add in petridish. Then,

at that point, place a tablet cautiously on the outer layer of the tissue paper. The time needed for water to arrive at upper surface of the tablet is noted as a wetting time.

$$R = 10 \left( \frac{W_a}{W_b} \right)$$

Where-  $W_b$  is weight of tablet before water absorption &  $W_a$  is weight of tablet after water absorption

**10) In vitro disintegration test:** In vitro crumbling audit should be performed by using USP type II Apparatus (paddle type) [Electro lab (ETC - 11L) Tablet disintegration tester] at 50 rpm. Phosphate cushion pH 6.8, 900 ml is fundamentally utilized as disintegration medium which is needed to keep up with at  $37 \pm 0.5^\circ\text{C}$ . Aliquot of (10ml) disintegration medium is needed to pull out at explicit time span (2min) and afterward it is needed to subject for cycle of filtration. How much medication not entirely settled by UV

Spectrophotometer (Shimadzu, japan) by estimating the absorbance of the example. Three preliminaries of each group were performed and normal % drug discharge with standard deviation was determined and recorded.

**11) Accelerated Stability study:** The Orally breaking down tablets are stuffed in reasonable bundling and put away under the accompanying conditions for a period as recommended by ICH rules for sped up investigations.

(I)  $40 \pm 1^\circ\text{C}$

(ii)  $50 \pm 1^\circ\text{C}$

(iii)  $37 \pm 1^\circ\text{C}$  and Relative Humidity=  $75\% \pm 5\%$

Pull out the tablets following a time of 15 days and examined for actual portrayal (Visual imperfections, Hardness, Friability, Disintegrations, and Dissolution and so forth) and medication content. The information acquired is fitted into first request conditions to decide the energy of corruption. Sped up strength information are plotting agreeing Arrhenius condition to decide the timeframe of realistic usability at  $25^\circ\text{C}$  [15] Future Prospective for MDTs Now there are different items accessible industrially in market which is delivered by quick dissolving tablet innovations. Still there is wide region for research on this innovation. A portion of the difficulties like planning a medication of harsh taste and dampness retaining nature make issues for definition researcher. At the point when the portion of medication is huge it causes issue of expanded breaking down time. The two focuses to be considered if there should be an occurrence of MDTs are shortening. Future forthcoming with proceeded with advancements in drug excipients, one can expect the development of more clever advances for MDTs in the days to come. These developments might include changing detailing structure and handling to accomplish new execution end-focuses or the consolidation of new mechanical advances with customary drug handling methods for the creation of novel mouth dissolving measurements structures. It is sensible to expect that future patterns in developments of medication conveyance frameworks will keep on uniting diverse mechanical disciplines to make novel technologies. He deterioration time simultaneously keeping different boundaries like friability, taste, and mouth feel and tablet strength inside the acknowledged reach. Utilizing taste covering specialists and super-deteriorating specialists without huge expansion in the weight and volume of definite measurements structures. Likewise, there is a degree to develop better packaging system to make FDTs more stable during handling.

## CONCLUSION

The MDTs enjoy expected upper hands over customary dose structures, with their worked on persistent consistence, accommodation, bioavailability and quick beginning of activity had drawn the consideration of many produces more than 10 years. MDTs definitions acquired by a portion of these innovations have adequate mechanical strength, fast crumbling/disintegration in the mouth without water. There is an unmistakable chance for new improved oral items emerging inside this market fragment. Around 33% of the populace, essentially the geriatric and pediatric populaces, has gulping challenges, bringing about helpless consistence with oral tablet drug treatment which prompts diminished by and large treatment viability... These tablets are intended to break down or deteriorate quickly in the spit for the most part inside <60 seconds (scope of 5-50seconds). The improvement of a quick dissolving tablet additionally gives an open door to a line augmentation in the commercial center; a wide scope of medications (e.g., neuroleptics, cardiovascular medications, analgesics, allergy medicines, and medications for erectile brokenness) can be viewed as possibility for this dose structure. As a medication substance approaches the finish of its patent life, it is normal for drug producers to foster a given medication element in a better than ever measurement structure

**REFERENCES**

- [1] Modasiya M, Lala I, Prajapati B, Patel V, Shah D. Design and characterization of Fast Disintegrating Tablets of Piroxicam. *International Journal of PharmaTech Research*, **2(1)**, 353-357 (2009).
- [2] Sahoo S, Mishra B, Biswal P, Panda O, Mahapatra S, Jana G. Fast Dissolving Tablet: As A Potential Drug Delivery System. *Drug Invention*, **2(2)**, 130-135 (2010).
- [3] Kaur T, Gill B, Kumar S, Gupta G. Mouth Dissolving Tablets: A Novel Approach to Drug Delivery. *International Journal of Current Pharmaceutical Research*, **3(1)**, 1-7 (2011).
- [4] Nayak U, Patra S, Rout P, Patro B, Nayak B. Development and optimization of prometazine theoclate Mouth dissolving tabltes. *The Indian Pharmacist*, **4**, 65-68 (2008).
- [5] Reddy L, Ghosh B, Rajneesh R. Fast Dissolving Drug Delivery system: A review of the literature. *Indian Journal of Pharmaceutical Sciences*, **64(4)**, 1-3 (2002).
- [6] Kuchekar B, Arumungam V. Fast Dissolving Drug Delivery System. *Indian Journal of pharmaceuticals and research education*, **35**, 150-155 (2001).
- [7] Bhushan Y, Sambhaji S, Anant R, Kakasaheb R. New drug delivery system for elderly. *Indian Drugs*, **37**, 312-318 (2000).
- [8] Gregory G, Ho D, U.S.Patent 4, 305,502 (1981).
- [9] Koizumi K. New Method of Preparing high porosity rapid saliva soluble compressed tablet using mannitol with camphour, a subliming material. *International Journal of Pharmaceutics*, **152(1)**, 127-131 (1997).
- [10] Allen V, Wang B, U.S. Patent U55587180 (1996).
- [11] Gupta A, Mittal A, Jha K. Fast Dissolving Tablet- A Review. *The Pharma Innovation*, **1**, 1-8 (2012).
- [12] Panigrahi R, Behera S. A Review on Fast Dissolving Tablets. *Webmed Central Quality and patient safety*, **9**, 1-15 (2010).
- [13] Mishra U, Prajapatil S, Bharadvaj P. A Review Article on Mouth Dissolving Tablet. *International Journal of Pharma Research & Review*, **3(6)**, 24-34 (2014).
- [14] Indian Pharmacopoeia. Controller of Publications. 4th edition, Volume 2, New Delhi, pp735-736; 1996.
- [15] Abdul S, Mohammad H. Mouth dissolving tablets: An overview. *International Journal of Research in Pharmaceutical and Biomedical Sciences*, **2(3)**, 959-970 (2011).

