



Antimicrobial Activity of Root Extract of *Aristolochia indica* L.

¹N. Yasothkumar, ²K. Rajendran, ²S. Shanmugam, ³S. Venkatesh

¹Department of Botany, Yadava College, Tiruppalai, Madurai – 625014, Tamil Nadu, India

²Post Graduate and Research Department of Botany, Thiagarajar College, Teppakulam, Madurai – 625009, Tamil Nadu, India

³Post Graduate and Research Department of Botany, Saraswathi Narayanan College, Perungudi, Madurai – 625022, Tamil Nadu, India

ABSTRACT

The medicinal properties of plants could be based on their pharmacological effects of the phytochemicals present in them. Hence plant derived antimicrobial properties have received considerable attention in recent years. In the present study, an attempt has been made to screen the antimicrobial potential of *Aristolochia indica* (root), against disease causing bacterial and fungal pathogens. The results revealed that the methanol extract had an effective antimicrobial activity against all the tested microorganisms and the hexane extract had low antimicrobial activity. Further studies should be carried out to evaluate the other biological active potentials of *Aristolochia indica*.

Keywords: Antimicrobial activity, *Aristolochia indica*, Root extract.

1. INTRODUCTION

Microorganisms gain resistance to the existing drugs and cause undesirable side effects. Antimicrobial properties have been reported in a wide variety of plant extracts with goal to discover new chemical classes of antibiotics that could resolve these problems¹. The plant derived compounds show broad-spectrum of biological activity against infection causing agents such as bacteria, fungi, protozoans, viruses and yeast^{2,3}. The potential for developing antibiotics from higher plants appears rewarding, as it will lead to the development of phytomedicine. In this concern, the present study was aimed to evaluate the antimicrobial potential of root extract of *Aristolochia indica*.

2. MATERIALS AND METHODS

2.1. Preparation of extracts

Fresh root samples of *Aristolochia indica* L. (Family: Aristolochiaceae; Local name: Perumarundhu) were collected from Karandamalai located in Dindigul district of Tamil Nadu, India. It lies between 19.2849° N latitude and 78.2169° E longitude. The collected specimens were botanically confirmed and authenticated as per the data pertained in The Plant List⁴. 25 gram of air-dried root parts was powdered and extracted with 250 ml of hexane, ethyl acetate and methanol separately. The extracts were then filtered and used directly for antimicrobial activity.

2.2. Procurement of microbial strains

Six pathogens such as *Bacillus cereus*, *Staphylococcus epidermidis*, *Escherichia coli*, *Salmonella paratyphi*, *Beauveria bassiana* and *Candida albicans* were obtained from Bose X-ray and Clinical Laboratory, Madurai, Tamil Nadu. Stock cultures were maintained at 4°C on slopes of nutrient agar. Active cultures for experiments were prepared by transferring a loopful of cells from the stock cultures to test tubes and were incubated without agitation for 24 hrs at 37°C.

2.3. Preparation of media (gram/litre)

Beef infusion	: 300
Casein hydrolysate	: 17.5
Starch	: 1.5
Agar	: 17

These ingredients were dissolved in 1000 ml distilled water; the pH was adjusted to 7.4 and sterilized at 121°C for 75 minutes.

2.4. Determination of antimicrobial activity

The microbial activity of the crude extract was determined in accordance with the agar well diffusion method⁵. 100 µl of cell suspension was spread on each petridish containing agar medium. Wells were bored onto the agar using a sterile 6 mm diameter cork borer. 20mg/ml of the crude extract was added into the wells and incubated at 37°C for 24 hrs. The plates were observed for zone of inhibition after 24 hrs. These effects were compared with streptomycin at a concentration of 10µg/ml.

3. RESULTS AND DISCUSSION

For the present study the antimicrobial activity of *Aristolochia indica* root extracts were tested against bacterial and fungal strains. The results showed that the methanol extract had an effective antimicrobial activity against all the tested microorganisms. The methanol extract produced the inhibition zone of 16±0.5 mm against *Escherichia coli*, 16±0.5 mm against *Staphylococcus epidermidis*, 16±0.5 mm against *Beauveria bassiana*, 14±0.6 mm against *Candida albicans*, 13±0.4 mm against *Salmonella paratyphi* and 12±0.4 mm against *Bacillus cereus*. Meanwhile, the hexane extract had low antimicrobial activity against all the tested microorganisms (Table 1; Fig 1).

Table 1: Antimicrobial potential of *Aristolochia indica*

Strains	Zone of inhibition (mm)*			
	Streptomycin	Hexane extract	Ethyl acetate extract	Methanol extract
<i>Bacillus cereus</i>	21±0.4	8±0.4	10±0.6	12±0.4
<i>Staphylococcus epidermidis</i>	20±0.4	10±0.3	11±0.2	16±0.3
<i>Escherichia coli</i>	21±0.1	11±0.6	12±0.3	16±0.5
<i>Salmonella paratyphi</i>	22±0.6	7±0.2	12±0.4	13±0.4
<i>Beauveria bassiana</i>	21±0.5	8±0.7	12±0.8	15±0.3
<i>Candida albicans</i>	23±0.3	7±0.5	11±0.6	14±0.6

*Values are Mean ± Standard Error of triplicates




Figure 1: Comparative antimicrobial activity of *A. alata* against different pathogens

The present study revealed that the methanolic extracts demonstrated the highest antimicrobial activity when compared to the other solvents (hexane and the ethyl acetate). This may be due to the better solubility of the active components in methanolic solvent⁶. These results coincide with the previous works⁷⁻¹¹ and some of these observations have helped in identifying the active principle responsible for such activities and in the developing drugs for the therapeutic use in human beings.

4. CONCLUSION

Further research is needed for the evaluation of other biological active potentials like antioxidant, antidiabetic, anti-inflammatory activities of *Aristolochia indica* *in vitro*. For this, large scale isolation and spectral techniques are required to isolate and identify a particular compound responsible for respective pharmacological action. The *in vivo* studies should also be carried out to describe the mechanism of action of these compounds in living systems using animal models.

5. REFERENCES

1. Piddock KJV & Wise R. Mechanism of resistance to quinolones and clinical perspective. *Journal of Antimicrobial Chemotherapy*, 1989; 23:475-483.
2. Boyd RF. *Basic Medical Microbiology* (5th Edition). Little, Brown and Company, Boston, 1995; 310-314.
3. Cookson BD. Methicillin-resistant *Staphylococcus aureus* in the community: New battlefronts, or are the battles lost? *Control of Hospital Epidemiology*, 2000; 21:398-403.
4. The Plant List. 2021. The Plant List: A working list of plant species, version 1. <http://theplantlist.org/>. Accessed on 17th April, 2021.
5. Irobi ON, Moo-young M, Anderson WA & Daramola SO. Antibacterial activity of bark extracts of *Bridelia ferruginea*. *Journal of Ethnopharmacology*, 1994; 43:223-224.
6. De Boer HJ, Kool A, Broberg A, Mziray WR, Hedberg I & Levenfors JJ. Antifungal and antibacterial activity of some herbal remedies from Tanzania. *Journal of Ethnopharmacology*, 2005; 96:461-469.
7. Poornima N, Umarajan KM & Babu K. Studies on anatomical and phytochemical analysis of *Oxystelma esculentum* (L.f.) R.Br. ex. Schultes. *Botany Research International*, 2009; 2(4):239-243.
8. Ismail Shareef M & Leelavathi S. Evaluation of antioxidant activity of *Anisolmeles malabarica* R.Br. and *Clerodendrum serratum* L. extracts against rheumatism. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 2011; 2(4):488-495.
9. Annie Felicia F & Muthulingam M. Phytochemical and HPTLC studies of methanolic extract of *Indigofera tinctoria* (Fabaceae). *International Journal of Pharmacy and Life Sciences*, 2012; 3(5):1670-1674.
10. Ilango K, Maharajan G & Narasimhan S. Preliminary phytochemical screening and antibacterial activity of fruit pulp of *Momordica dioica* Roxb. (Cucurbitaceae). *African Journal of Basic & Applied Sciences*, 2012; 4(1):12-15.
11. Usha M, Appavoo MR & Immanuel G. Pharmacognostical study and preliminary phytochemical screening of the leaf of *Ixora johnsonii* Hook.f. - A rare, endemic, critically endangered species of southern Western Ghats of Kerala, India. *International Journal of Pharmaceutical and Phytopharmacological Research*, 2012; 5:263-270.

