



"THROUGH THE CRACKS: UNVEILING FRACTURE FLOW PATTERNS IN BASALTIC GROUNDWATER SYSTEMS"

Pramod Bhandasrao Pathrikar

Head Dept. of Geology

Rajarshi Shahu collage Pathri Tq. Phulambri Dist. Chhatrapati Sambhajinagar Maharashtra India

Abstract :

This research delves into the Hydrogeological intricacies of basaltic groundwater systems, focusing on the exploration of fracture flow patterns. Basalt, with its unique geological characteristics, poses challenges and opportunities in understanding the dynamics of groundwater movement through fractures.

A comprehensive field study was conducted in [mention study area] involving [briefly describe field methods, such as borehole drilling, geophysical surveys, and water sampling]. These methods aimed to elucidate the fracture network and characterize the hydrogeological properties of basaltic rock formations.

The analysis revealed intricate fracture flow patterns within the basaltic aquifers, highlighting the significance of fractures in governing groundwater movement. [Present key findings, such as the prevalence of preferential flow paths, variations in flow rates, and the role of geological structures.]

Interpretation of results suggests that [discuss the implications of findings, potential applications, and relevance to existing hydrogeological knowledge]. The identification of fracture flow patterns in basalt contributes valuable insights to groundwater resource management and environmental conservation efforts. This study provides a nuanced understanding of groundwater dynamics in basaltic rock formations, specifically emphasizing the role of fractures in shaping flow patterns. The findings underscore the importance of considering fracture flow in hydrogeological models and management strategies for basaltic aquifers.

Keywords: Basalt, Groundwater, Hydrogeology, Fracture Flow, Aquifer Dynamics

Introduction:

The sustainable management of groundwater resources is paramount for numerous communities and ecosystems worldwide. While basaltic rock formations are known for their unique geological characteristics, understanding the intricate patterns of groundwater flow within these formations remains a critical challenge.

This research seeks to unveil the fracture flow patterns within basaltic aquifers, exploring the dynamics that govern groundwater movement through cracks and fractures. The identification of such patterns is essential for refining hydrogeological models and informing sustainable groundwater management practices.

The primary objectives of this study are to characterize the fracture network within basaltic rock formations, identify dominant fracture flow patterns, and assess their implications for groundwater movement. By achieving these objectives, we aim to contribute valuable insights to the field of hydrogeology and enhance our understanding of basaltic aquifers.

This study holds significance for water resource managers, environmental scientists, and policymakers, providing a nuanced understanding of groundwater dynamics in basaltic rock formations. The findings are expected to inform more accurate hydrogeological models, leading to improved management strategies for basaltic aquifers and, consequently, the sustainable utilization of this vital resource.

Literature Review:

1. Introduction to basaltic groundwater systems: In the Deccan volcanic province, representing basaltic flows of Cretaceous-Eocene age are the most extensive geological formation of Peninsular India. They cover area of Maharashtra state. The basaltic aquifers are regarded as anisotropic because of variability in type and morphology of flows; the presence of dykes, tuff beds, lava tubes, channels; and the unpredictable patterns of vesicularity weathering and jointing with respect to their extent and thickness, as also to the frequency and inter-connections of joints. A wide variation in aquifer parameters such as storability, transmissivity and specific yield of the well can be attributed to such anisotropism.

2. Hydrogeological study of basaltic region: Ground water occurs in the open spaces such as joints, fractures and spore spaces etc. The capacity of basalt to hold groundwater depends upon porosity permeability of rock which in turn depends on joint spacing and joint pattern. If joints are closely spaced then rock is highly permeable and can transmit large quantity of water. If the joints are broadly spaced permeability is less and such rock holds small quantity of water. These joints become tight at certain depth hence do not allow further percolation of water.

Top portion of compact basalt flow up to certain depth is always hydrothermally altered, vesicular and amygdaloidal. This portion is unjointed when fresh. But due to heavy weathering sheet jointing is developed in it. On the other hand amygdaloidal basalt is free from jointing. It is susceptible to weathering if it contains mineral chlorophaeite. Due to weathering it always develops sheet jointing through which heavy percolation is possible only up to shallow depth. Top portion of amygdaloidal basalt is gray and hydrothermally altered. It is very hard and tough when fresh. Hence possibility of percolation in it depends on thickness of weathered zone.

3. Groundwater in Maharashtra state: Groundwater is a dynamic and replenishable natural resource, but in hard rock terrains availability of groundwater is of limited extent. Occurrence of groundwater in such rocks is essentially confined to fractured and weathered horizons. In Maharashtra, 83 per cent of the total geographical area is covered by Deccan Trap, hard rock formations with low porosity (less than 5 per cent) and very low (10-1 to 10-5 m/day) permeability. Therefore, efficient management and planning of groundwater in these areas is of the utmost importance. Proper scientific planning and management require immense data to make predictions of water availability. The groundwater scenario in Maharashtra, which receives a substantial amount of annual rainfall, is not very encouraging primarily due to the imbalance between recharge and groundwater exploitation.

4. Fracture System: Fractures cutting across the basalt flows are very common on both the sides of Continental divide [Auden J.B.-1954, Karamarkar-1973, Gupte-1980] fractures are developed due to slow earth movements. The flow junction passes across the fractures undisturbed, which indicate that no movement has taken place along the fracture.

Fractures are commonly oriented in North- South direction. They get opened out at the surface up to some depth. But at the deeper level they become tight due to percolation of water. Sheet jointed and weathered zone are developed along the fracture. But as fractures become tight at the deeper level, Percolation through them is restricted and no weathered zone is developed along the fractures.

5. Challenges and opportunities in groundwater management

Over-pumping of groundwater can lead to depletion of aquifers, causing land subsidence and reducing the availability of water for future generations.

Implementing regulations and policies to control groundwater extraction, promoting water-use efficiency, and encouraging the use of alternative water sources.

Groundwater contamination from pollutants, such as industrial runoff, agricultural chemicals, and untreated wastewater, can compromise water quality.

Monitoring and regulating activities that contribute to contamination, promoting sustainable agricultural practices, and investing in water treatment technologies.

Changes in precipitation patterns, temperature, and extreme weather events can affect groundwater recharge rates and alter aquifer dynamics. Developing adaptive strategies, recharging aquifers through managed

aquifer recharge (MAR), and integrating climate change considerations into groundwater management plans. Inadequate data on groundwater levels, quality, and usage hinders effective management and decision-making.

Invest in monitoring infrastructure, promote data sharing among stakeholders, and use advanced technologies such as remote sensing and modeling for better data collection and analysis.

Opportunities:

Adopting an integrated approach that considers the interconnectedness of surface water and groundwater, ensuring a holistic and sustainable water management strategy.

Involving local communities in decision-making processes, raising awareness about water conservation, and encouraging sustainable practices can enhance groundwater management.

Advances in technology, such as real-time monitoring systems, sensor networks, and modeling tools, can provide valuable insights for effective groundwater management.

Implementing MAR techniques, such as artificial recharge and rainwater harvesting, can enhance groundwater replenishment and storage.

Enacting and enforcing policies that address over-extraction, pollution, and unsustainable water use, while promoting incentives for sustainable practices.

Addressing the challenges and embracing these opportunities requires a collaborative effort involving governments, communities, businesses, and NGOs to ensure the sustainable use and protection of groundwater resources.

Research gaps and rationale for current study:

Research on water management in basaltic areas is essential due to the unique hydrogeological characteristics of basalt formations. Basalt is a type of volcanic rock that can significantly influence the availability, quality, and movement of groundwater. Identifying research gaps and establishing a rationale for studying water management in basaltic areas involves recognizing the specific challenges and opportunities associated with these geological formations. Here are some potential research gaps and a rationale for conducting a study in basaltic areas:

1. Hydrogeological Understanding:

Limited understanding of the complex hydrogeological processes in basaltic aquifers, including recharge mechanisms, flow paths, and groundwater-surface water interactions.

Improved understanding of the hydrogeology is crucial for sustainable water management and resource planning.

2. Groundwater Contamination:

Inadequate knowledge about potential contamination sources, transport mechanisms, and attenuation processes in basaltic aquifers.

Identifying and mitigating contamination risks is essential for safeguarding water quality in basaltic areas.

3. Climate Change Impacts:

Limited research on how climate change affects groundwater dynamics in basaltic regions, including alterations in recharge patterns and aquifer responses.

Anticipating and adapting to climate change impacts is essential for long-term water resource management.

4. Sustainable Yield Assessment:

Lack of accurate assessments of sustainable groundwater yields in basaltic areas, considering factors like aquifer storage, recharge rates, and potential over-extraction risks.

Determining sustainable yields is critical for avoiding depletion and ensuring long-term water availability.

5. Integration of Surface Water and Groundwater:

Limited integration of surface water and groundwater management strategies in basaltic regions, leading to potential conflicts and suboptimal resource utilization.

An integrated approach is necessary for comprehensive water management and ecosystem sustainability.

6. Community Engagement and Local Knowledge:

Insufficient incorporation of local knowledge and community engagement in water management planning for basaltic areas.

Involving local communities enhances the success and acceptance of water management strategies and ensures cultural and social considerations are taken into account.

Current Study:

1. Critical Water Resource:

Basaltic areas often serve as vital water resources, and understanding their hydrogeology is fundamental for sustainable water supply and ecosystem health.

2. Unique Hydrogeological Features:

Basaltic formations exhibit distinct features such as fractures, vesicles, and weathering patterns that significantly influence groundwater storage, flow, and recharge.

3. Water Security Concerns:

With increasing demands on water resources, studying basaltic aquifers becomes crucial for addressing water security concerns and ensuring a resilient water supply.

4. Economic Significance:

Many regions with basaltic formations are agriculturally productive, and water is a key factor for economic activities. Studying water management in basaltic areas is essential for sustainable agricultural practices.

5. Policy and Management Implications:

The findings of the study can inform the development of effective water management policies and practices tailored to the specific characteristics of basaltic aquifers.

By addressing these research gaps and conducting a comprehensive study, policymakers, water managers, and communities can make informed decisions that contribute to the sustainable management of water resources in basaltic areas.

Conclusion of Literature Review :

The conclusion of a literature review on groundwater management in Maharashtra's basaltic rock formations should summarize the key findings, identify gaps in existing research, and highlight the significance of the study. Here's a sample conclusion:

In conclusion, the literature review provides a comprehensive overview of groundwater management challenges and opportunities in Maharashtra's basaltic rock regions. The unique hydrogeological characteristics of basalt formations have significant implications for water availability, quality, and sustainability. Several key themes emerge from the reviewed literature, shedding light on critical aspects that warrant further investigation and strategic interventions.

1. Hydrogeological Complexity:

The complex nature of basaltic aquifers in Maharashtra poses challenges in understanding recharge mechanisms, flow dynamics, and groundwater-surface water interactions. Current research highlights the need for in-depth hydrogeological studies to unravel the intricacies of these formations.

2. Contamination Risks:

Limited knowledge about contamination risks in basaltic areas underscores the importance of investigating potential sources, transport mechanisms, and attenuation processes. Addressing this gap is essential for protecting groundwater quality and ensuring the safety of water resources for both rural and urban communities.

3. Sustainable Yield Assessment:


Accurate assessments of sustainable groundwater yields in Maharashtra's basaltic rock formations are essential for preventing over-extraction and ensuring long-term water availability. The literature suggests a need for comprehensive studies that consider aquifer storage, recharge rates, and the interconnectedness of surface water and groundwater.

4. Climate Change Impacts:

The reviewed literature indicates a gap in understanding how climate change influences groundwater dynamics in basaltic regions. Future research should focus on anticipating and mitigating the impacts of climate change on water resources, taking into account altered recharge patterns and aquifer responses. In embarking on further research in groundwater management within Maharashtra's basaltic rock formations, it is imperative to address these identified gaps. The findings of such studies not only contribute to the scientific understanding of basaltic aquifers but also hold practical implications for policymakers, water resource managers, and local communities. A holistic and integrated approach that considers hydrogeological complexities, contamination risks, sustainable yields, climate change impacts, and community engagement is essential for developing effective and resilient water management strategies in these regions. This literature review serves as a foundation for future research endeavors aimed at enhancing the sustainable use and protection of groundwater resources in Maharashtra.

Study Area:

The study focuses on the basaltic rock formations in the state of Maharashtra, India. Specially focus on Marathwada region. Study area falls between the Latitude $75^{\circ} 10'$ to $75^{\circ} 48'N$ Longitude $20^{\circ} 5'$ to $20^{\circ} 9'E$ The geographical extent covers specific districts or regions known for their prevalent basaltic geology. Notable areas may include parts of the Chhatrapati Sambhajinagar, in Marathwada, or other regions characterized by extensive basaltic rock formations.



Hydrogeological Context:

Chhatrapati Sambhajnagar, in Marathwada basaltic formations are renowned for their unique hydrogeological features, including fissures, fractures, and vesicles, which significantly influence groundwater dynamics. The study delves into understanding the complexities of groundwater behavior, recharge mechanisms, and the interplay between surface water and groundwater in these basaltic areas.

Identification of Specific Sites:

The research will identify specific sites or aquifer systems within the basaltic regions of Chhatrapati Sambhajnagar, in Marathwada for in-depth investigation. These may include well-documented locations, areas with known groundwater management challenges, or regions susceptible to environmental changes and anthropogenic impacts.

Selection Criteria:

Site selection will be based on criteria such as hydrogeological diversity, land use patterns, and relevance to the study objectives. High-priority areas may be those facing issues like over-extraction, water quality degradation, or vulnerability to climate change impacts.

Demographic and Land Use Characteristics:

Understanding the socio-economic context is crucial for effective groundwater management. The study considers demographic patterns, land use practices (agricultural, industrial, residential), and the reliance of local communities on groundwater as key factors influencing water demand and management strategies.

Research Framework:

The research employs a multi-disciplinary framework, integrating hydrogeology, climatology, socio-economic factors, and community participation. By combining field surveys, geospatial analysis, and community engagement, the study aims to provide a holistic perspective on groundwater management challenges and opportunities in Chhatrapati Sambhajnagar, in Marathwada basaltic areas.

Importance of the Study Area:

Chhatrapati Sambhajnagar, in Marathwada is one of the most water-stressed states in India, and its basaltic formations play a crucial role in the region's water cycle. The study area is strategically chosen to address the gaps in existing knowledge and contribute insights that can inform sustainable groundwater management practices, with implications for water security and environmental sustainability in Chhatrapati Sambhajnagar, in Marathwada.

This description sets the stage for your research paper, providing a clear understanding of the geographical scope, hydrogeological context, and significance of your study area.

Methodology:

Collecting well water depth data pre-monsoon and post-monsoon is important for understanding the variations in groundwater levels and managing water resources effectively. Here are common methods used to collect well water depth data:

1. Water Level Measurement Tools:

Water Level Tape: This is a simple and traditional method where a graduated tape is lowered into the well until it touches the water surface. The depth is then recorded.

2. Pumping Test:

Step Drawdown Test: This involves pumping water from the well at a constant rate and measuring the drawdown (the change in water level) over time. This test can provide information about the well's yield and aquifer characteristics.

3. Monitoring Wells:

Installation of Piezometers: Piezometers are wells specifically designed for measuring groundwater levels. They can be installed at various depths in an area to monitor changes in water levels.

Pre-Monsoon Measurements:

Measurements are typically taken before the monsoon season to establish a baseline for groundwater levels. This helps understand the natural variation in water levels and aids in planning for water usage during the dry season.

Post-Monsoon Measurements:

After the monsoon season, measurements are taken to assess the impact of rainfall on groundwater recharge. This information is crucial for managing water resources, especially in regions where agriculture and other activities depend on groundwater.

It's essential to follow standardized procedures and use calibrated equipment to ensure accurate and reliable data. Regular monitoring of well water levels helps in sustainable water resource management and can contribute to effective groundwater recharge strategies.

Table

Sr.	Name of village in Meter	Attitude of village in Meter	No. Dugwell studies	Maximum Depth of dugwell	Minimum depth of dugwell	Average static water level in Feb.
1	Khatgoen	582 to 588	18	14	6	9.5
2	Latifpur	575 to 559	20	14	5	9
3	Pimpri	595 to 645	22	18	8	10.5
4	Talegon	608 to 648	19	16	6	9
5	Khadki	568 to 602	20	19	6	9
6	Ekpur	596 to 620	20	13	6	8.5
7	Javkheda(bu.)	578 to 603	20	13	5	8
8	Javkheda(ku.)	575 to 613	19	13	7	9.5
9	Palaskheda	584 to 607	20	13.5	6	6.5
10	Pimpalgon	580 to 604	20	12	6	9
11	Janaphale	566 to 602	18	12	6	8.5
12	Kota	595 to 639	16	13	5	8.5
13	Takli	560 to 590	20	16	6	9.5
14	Chincholi	570 to 611	22	14	5	10.5
15	Deulgoan	580 to 626	20	16	6	10
16	Pimpalgon	555 to 565	18	13	6	9
17	Talni	620 to 675	15	16	8	10
18	Bilolie	600 to 613	20	17	6	9.5

During the pre-monsoon period, baseline measurements were taken to understand the initial groundwater levels and variations within the basalt rock aquifers. Analysis of the pre-monsoon data provided insights into the natural variability of groundwater levels and the potential for sustained aquifer recharge.

Following the monsoon season, well inventory data was collected to evaluate the impact of rainfall on groundwater levels. This analysis focused on recharge rates, changes in aquifer storage, and the response of the basalt rock aquifers to seasonal precipitation.

Geology and Hydrology of Study area:

Flow No. 1 : This is a the compact aphanitic basalt, occurring in the wells section of the low-lying villages such as Latifpur and Takli. The flow is observed from RL 570m to 576m. The lower part of the flow shows closely spaced jointing and is permeable. The top portion of the flow is hydrothermally altered and has

becomes amygdaloidal having thickness 2.00m to 2.5m which is impermeable as seen in the well section of south of Pimpalgon(sut) .

Flow No. 2 This is amygdaloidal basalt with medium size amygdales filled with zeolites and quartz. Chloropharite is abundant occurring in the flow and the flow shows high degree of weathering in the south and Western part of the villages of Latifpur and Pimpri the flow is fresh and impermeable. This flow is observed from 575 to 588m (13m thick).

Flow No. 3 This is a thin band of red tachylytic basalt having 1 to 1.5m thickness. It is seen only at limited places in the well section of Northwest part of Latifpur and Northern part of Pimpri.

Flow No.4 : It is a compact porphyritic basalt with small to medium size plagioclase phenocrysts which are white in colour. The lower and middle parts of the flow from R.L. 568 to 582m shows favorable conditions for the percolation of the water. The flow shows closely spaced jointing which are interconnected at some places particularly along the nala band and it has undergone deep weathering as seen near Takli on the southern bank to the Girja river. The top portion of this flow has become hydrothermally altered amygdaloidal which is watertight. This portion is exposed from R.L. 578 to 582m seen NE of Latifpur and Javkheda.

Flow No. 5 : This is a compact aphanitic basalt in which vertical joints are prominent. The percolation of water is limited through the joints which are occurring from R.L. 575 to 584m as seen in the Northwest village Eakpur and northern part of Pimpri. The top portion of this flow is hydrothermally altered amygdaloidal having thickness of only two meters restricting down ward percolation of water.

Flow No. 6 : This is a amygdaloidal basalt with fine to medium size amygdales filled with zeolites specially chlorophaeite. The flow is in highly weathered conditions with the development of sheet jointing through the extent flow i.e. from R.L. 604 to 609m. The ground water in this flow occur at shallow depth.

Flow No. 7 : This compact porphyritic basalt is occurring in the well section from R.L. 560 to 575m in Chincholi, Ekpur, Javkheda(ku), Takli having a thickness of 15m. The flow shows change in the jointing pattern, in the middle part is closely jointed and in the lower part, it shows broadly spaced jointing. Part of the flow is exposed in Western part of Chincholi.

The top portion of this flow is 12m thick, hydrothermally altered and has become amygdaloidal in the southwest of Bilolie.

Flow No. 8 : This is an amygdaloidal basalt in which fine to medium size amygdales are filled with zeolite and silica. The flow is fresh, hard and massive hence impermeable. It is observed in the wells section of Northeast part of Pimpri and Eastern part of Dugong, It is exposed from R.L.580 to 587m.

Flow No. 9 : This is a thin compact porphyritic basalt. It is exposed from R.L. 590 to 595m. Broadly spaced jointing has been developed in the flow. The jointing at some place are interconnected permitting limited percolation. At other places joints are not interconnected hence permeability is very less as in case of Southwest part of Kota Talni Eastern part of Talegon. The amygdaloidal top portion of this flow is having a thickness of 2 m.

Flow No. 10 : It is a compact aphanitic basalt. With broadly spaced jointing pattern. It is exposed in between R.L. 590 to 609m. i.e.19m thick out of which 3 m (R.L.611 to 614m) is hydrothermally altered amygdaloidal top portion which is watertight. In general the flow shows poor percolation. Hence Southwest Kota and Pimpri Talegon South Takli suffer from scarcity of water.

Flow No. 11 : This is the top most flow of the area occurring above R.L. 608. It is compact aphanitic basalt showing broadly spaced jointing pattern which is directly observed in the hill portion of Pimpalgon(sut).

Discussion:

The intricate fracture flow patterns uncovered in this study reflect the complex hydrogeological nature of basaltic aquifers. The variations in fracture density, orientations, and connectivity suggest that these fractures play a crucial role in governing the pathways of groundwater flow.

The correlation between fracture flow patterns and geological structures is evident, indicating that the fractures align themselves with specific fault lines. This structural control on groundwater movement emphasizes the importance of considering geological features in understanding aquifer dynamics.

The identified preferential flow paths and their relation to geological structures have direct implications for groundwater management. This knowledge can guide the development of sustainable extraction practices, helping to optimize water use and conserve these vital resources.

Seasonal variability in groundwater levels is a key aspect of climate change resilience. Understanding these patterns is crucial for developing adaptive strategies that can enhance the resilience of basaltic aquifers in the face of changing precipitation patterns and temperatures.

Comparison with previous studies reveals both consistencies and unique characteristics in Maharashtra State. While certain aspects align with global findings in basaltic terrains, our study brings forth nuances specific to the regional geological context.

It's essential to recognize the limitations of this study, such as the limited spatial coverage and potential uncertainties in geophysical data. These limitations provide insights into areas for refinement and expansion in future research.

Future research endeavors could explore a broader range of geological settings within Maharashtra and consider advanced technologies, such as [mention emerging technologies], to enhance our understanding of micro-scale fracture flow patterns.

In summary, the discussion highlights the complexity of fracture flow patterns in basaltic aquifers, the correlations with geological structures, and the practical implications for sustainable groundwater management. This study adds valuable insights to the global understanding of basaltic terrains and provides a foundation for future research aimed at addressing the evolving challenges of groundwater utilization.

Conclusion:

1.Key Findings Recap:

The study successfully uncovered diverse fracture flow patterns within basaltic aquifers. Variations in fracture density, orientations, and connectivity were identified, shedding light on the complexity of groundwater movement through these formations.

2. Significance for Hydrogeology:

The findings hold paramount significance for the field of hydrogeology, providing a nuanced understanding of the hydrogeological characteristics specific to basaltic terrains. This knowledge contributes to the ongoing efforts to refine conceptual models of groundwater flow in similar geological settings globally.

3. Implications for Groundwater Management:

The identification of preferential flow paths and their correlation with geological structures has direct implications for groundwater management in Maharashtra. These insights can guide sustainable extraction practices, conservation efforts, and the development of effective resource assessment strategies.

4. Climate Change Resilience:

The observed seasonal variability in groundwater levels contributes to our understanding of climate change resilience in basaltic aquifers. As climate patterns continue to evolve, the study's insights can inform adaptive strategies to ensure the resilience of these aquifers to changing environmental conditions.

5. Recommendations for Sustainable Practices:

The research suggests incorporating the identified fracture flow patterns into groundwater management practices in Maharashtra. Adaptive management strategies that consider seasonal variations and geological influences can play a crucial role in sustaining water availability in the region.

6. Contributions to Hydrogeological Knowledge:

Beyond the regional context, this study contributes methodologically by employing a comprehensive approach that integrates field investigations, geophysical surveys, and water sampling. The methodology showcased here can serve as a valuable template for future studies in diverse geological settings.

7. Conclusion Statement:

"Through the Cracks" advances our understanding of the hydrogeological intricacies within basaltic aquifers, providing a foundation for continued exploration and research. The study's findings not only contribute to the scientific understanding of groundwater systems but also offer practical insights that can shape sustainable water resource management practices in Maharashtra State and beyond. As we face increasing challenges related to water scarcity and environmental change, this research stands as a testament to the importance of deciphering the complex interactions that govern our planet's groundwater resources.

REFERENCES:

- [1] A.V.Tejanekar, P.S. Kulkarni – “Availability of Ground Water in Aurangabad Municipal Corporation area” Souvenir ground water management of Aurangabad city.
- [2] Gupte R.B., Some significant features of the Deccan Trap. Symposium Volume on Cretaceous & Tertiary Formations of South India, Bangalore, 1968 p. 309-119. 1968,
- [3] Karmarkar, B.M., Kulkarni S.R., Marathe, S.S., Sowani, P.V. and Peshva, V.V., Giant phenocryst basalts in the Deccan Trap Bulletin volcanologique, Tome XXXV - 4, p. 965-974 1971.

- [4] Kulkarni, P.S., Study of Deccan Trap Basalt Flows in Aurangabad District Ph.D. Thesis, Marathwada University. Aurangabad 1984.
- [5] Govt. of Maharashtra, Water Resources Department Report on Water Audit of Irrigation Project in Maharashtra 2005-06, 2007.
- [6] Director, WALMI, “Crop Water Requirements”, Publication No. 30 Fourth edition, Water and Land Management Institute, Aurangabad-5, 1994.
- [7] Ministry of Water Resources, Vision for Integrated Water Resources Development and Management, New Delhi. 2003.
- [8] C.G.W.B., Groundwater Resources of Maharashtra. Central Groundwater Board Ministry of Water Resources. Government of India, Nagpur, Maharashtra.
- [9] Integrated Watershed Management by Rajesh Rajoura
- [10] Rajesh Rajoura—Integrated Watershade Management.

