



Review on Assistive Technology and Mobility Impairment

Priya M. Raut¹, Mohan S. Raut²

- Assistant Professor, Dept. Of Agadtantra Evum Vidhi Vaidyak, DMAMCHRC, Nagpur
- Associate Professor, Dept. Of Kaumarbhritya, DMAMCHRC, Nagpur

Corresponding Author-

1) Dr. Mohan S. Raut

Associate Professor, Dept. Of Kaumarbhritya, DMAMCHRC, Nagpur

Abstract-

For older persons or people with impairments, assistive technology (AT) may also refer to assistive, adaptive, and rehabilitative technologies. Activities of daily living (ADLs) can be challenging for people with disabilities to complete without help or even with assistance. The term "ADLs" refers to activities of daily living, which also include grooming, eating, clothing, bathing, and personal device maintenance. When a disability prevents someone from performing ADLs, assistive technology can help. Through improvements to or new ways of interacting with the technology required to complete certain tasks, assistive technology encourages increased independence by allowing users to carry out tasks that they were previously unable to complete or had a great deal of trouble completing¹. One essential exercise might be moving around the home and the neighbourhood, practitioners in rehabilitation should pay close attention since it is essential to widespread human participation. The work of rehabilitation practitioners can be informed by differentiating the restrictions and supports that clients see as having an impact on their present and desired life outcomes, as well as any impediments that are normally out of the rehabilitation observer's field of view¹. The ability to move around, as defined by the Oxford Wordbook of English², may be a fundamental aspect of human capability. The rehabilitation literature views freedom of movement as a crucial component of living, aboard communication, and self-care, ideally without the need for assistive technology (AT)³. Health-related quality-of-life evaluations also view the ability to move around independently as an important sign of life satisfaction.^{4,5}

Keywords – accessibility, assistive technology devices, mobility impairment, rehabilitation

Introduction –

With medical advances and growing understanding of the value of good health, diet, and hygiene, life expectancy is continuing to increase. There will be more elderly people in society as a result of longer anticipation, which will result in an increase in the number of people with age-related health disorders who may require assistance to manage their independence and daily activities. A further constraint is that care must be provided with limited financial resources because to the declining caregiver-to-care-recipient ratio. There is also a strong desire for older people to age in place and for this to be encouraged in order to preserve their independence for as long as feasible⁶.

An growth in anticipation has been accompanied by a continuous advancement in data and communication technology, which has resulted in an increase in efficiency and a decrease in cost. In instances when resources are limited and new healthcare requirements create a challenge that seems amenable to a technical solution, at least in part, technology is frequently used to produce solutions. There is an opportunity to use and shape the development of technology to deliver scalable and inexpensive healthcare⁷. Healthcare systems are frequently given through the use of appropriate technologies, and they may offer high-quality care within new resource restrictions. Activity monitoring is one of a variety of new technologies with applications. More frequently, especially for providing future care or management, the solutions that arise contain some technical component. Activity monitoring is the practise of keeping an eye on a person's behaviour and activity patterns over a period of time and using the data gathered to help with treatment. A constant assessment of activity levels is crucial to detect changes in health status because an individual's activity is linked to their health⁸. Additionally, a quantitative or direct measure of activity level is preferred over self-reported measures in order to meet the guidelines for older individuals set forth by the World Health Organization (WHO) for the amount of moderate and vigorous physical activity.

A type of sensor, loosely categorised as ambient or wearable, is used to keep track of a person's activity. Environmental sensors provide information about a user's activities at home or at work. Examples of environmental sensors include door/window switches, motion detectors (also referred to as space occupancy sensors) such as passive infrared (PIR) sensors, motion capture systems (video/camera systems, Kinect), power consumption monitors, as well as sensors that measure the environment's physical properties such as light, temperature, sound, and pressure. Information about the user's activity in their environment is gathered by using a mix of such sensors⁹. Body-worn sensors or wearable sensors provide information immediately about the user's activity. The wearable sensors include accelerometers, temperature, pressure, light, and camera sensors, as well as GPS (location) sensors. Accelerometers are widely used to monitor activity, and they work by placing a triaxial measuring instrument on the chest, waist, or several other sites.¹⁰ suggests that the approach to live PA, step counts, and energy estimation be included when choosing a measuring device to be employed in investigations with older persons. A simple approach of wearing sensors attached to the wrist is typically the most popular, especially for older people. Data from several wearable sensors will provide an excellent picture of the person's activity.¹¹ Activity monitoring has many different applications, ranging from general health to rehabilitation. The general well-being of any person, active living, sports, exercise, and watching daily activities all have their uses. It is helpful to monitor activity when treating

polygenic disorder, obesity, stroke rehabilitation, and other semi-permanent chronic illnesses including chronic obstructive pulmonary disease (COPD). Learning disorders like chronic fatigue syndrome and fibromyalgia benefit from activity observation. It is helpful to observe someone's capacity to execute Activities of Daily Living (ADL) in old age¹², which is useful in observing independent living in older people.

Role of Adaptive Technology -

Technology that is adaptive and assistive are two separate things. While adaptive technology refers to products that are especially made for impaired people and are rarely used by non-disabled people, assistive technology refers to objects that are known to help disabled individuals. In other words, assistive technology is anything that aids individuals with disabilities, whereas adaptive technology is made expressly for persons with disabilities. Thus, adaptive technology might be a division of assistive technology. Electronic and data technology access is one area where adaptive technology is frequently used.¹³ Therapeutic Occupational A medical specialty called occupational therapy (OT) aims to preserve or raise the quality of life for those who have difficulty carrying out their daily tasks on their own. Occupations include activities related to relaxation and sleep, education, work, recreation, leisure, and social interaction, in accordance with the physiotherapy Practice Framework: Domain and Process¹⁴ (3rd ed.; AOTA, 2014). For the enhancement and maintenance of optimal, functional participation in jobs, occupational therapists have the particular competence of using assistive technology (AT). A person can modify problematic features of the environment for the user using an AT appliance, maximising their functional participation in such occupations. Thus, occupational therapists may instruct advocate the use of ATs to raise the quality of life for their clientele.

Physical Mobility Impairments-

The category of incapacity known as mobility impairment includes people with a variety of physical limitations. This type of incapacity comprises loss or incapacity of the upper or lower limbs, facility, and incapacity in coordination with completely separate body organs. With age-related limitations, incapacity in quality might either be innate or not transmissible. This problem could possibly be a side effect of disease. Additionally, those among us who have structural flaws constitute this group of incapacity. Crutches, canes, wheelchairs, and artificial limbs are common quality aids used by people with physical disabilities to promote quality. The person's physical disability could even be inherited, caused by an injury, a neurological ailment, an amputation, MS, or pneumonic sickness, cardiovascular disease, or other factors may be to blame. Non-visible disabilities, such as diseases of the metabolism or epilepsy, may be experienced by some people¹⁵. Volunteers from the People with Disabilities Program wheel a chair around the US Air Force National Depot lobby. Each volunteer chooses a disability from the Associate in Nursing envelope after making the deposit that corresponds to the disability they anticipate. This icon depicts a disability that has an impact on the participant's legs. (Photo by the U.S. Air Force/Kimberly Gaither). Physical and qualitative impairment is defined by NANDA as a restriction on the free, intentional physical mobility of the body or of one or more extremities. According to them, the change in the individual's quality could also be either transitory

or more permanent. A degree of immobility is involved in the majority of illnesses and rehabilitative conditions that cause physical and quality deficits.

These measurements are typically linked to conditions like MS, limb fractures, strokes, morbid avoirdupois, trauma, and so on. According to NANDA, if Americans live longer, there will be more illnesses and incapacities in the country. Shorter hospital stays are leading to more patients being discharged to their homes for medical treatment or being moved to rehabilitation centres. It is said that when people age, their bodies lose some of their mobility. Loss of muscle mass and strength, stiffer and less movable joints, as well as changes in gait, all affect a person's balance and may greatly degrade it. For elders to continue working as freelancers, quality is essential. When a person's quality is constrained, it advances to have some measure of control their routine everyday tasks¹⁶.

Students with mobility issues may have some effects at this time. Some students might require more time to access buildings, move about in tight spaces, or transition from one class to another. Physical obstacles may occasionally make it difficult to enter a structure or space. Additionally, accessible transportation may be needed to get school students to construction sites. A quality impairment may be permanent or transient. A student's ability to manoeuvre carefully and move between nearby buildings in an exceptionally very timely manner can be swiftly affected by a broken bone or surgery. For short distances within a space, some kids may even be mobile with a walker, but for longer trips, they may have a chair or scooter¹⁷.

Conditions for Mobility Impairment-

Cerebral palsy-

Several chronic diseases that impact people with the disorder's ability to move freely and coordinate their muscles are collectively referred to as "Cerebral Palsy." Spastic paralysis harms at least one or more specific parts of the person's brain. It often occurs during vertebrate development or just before, during, or shortly after delivery, though the impairment could also occur during infancy.

Spastic paralysis disorders don't appear to be brought on by problems with the person's nerves or muscles. The ability of a person's brain to control posture and movement is disrupted by developmental flaws or damage to their motor areas.

Spastic paralysis is not a progressive disorder, but it may coexist with other conditions like muscle jerkiness that may become worse or get better with time might not change Although there is no known cure for spastic paralysis, training and medical attention may help in improving performance.¹⁸

Spina Bifida-

Possibly a mesoderm deficiency, Spina Bifida. Mesoderm abnormalities, which result from the fetus's spine failing to close properly during the first month of physiological state, involve the inadequate development of the brain, funiculus, and/or their protective coverings. Children who are born with a congenital condition or an associated congenital defect may have an open lesion on their spine where significant damage to their nerves and funiculus

occurred. Although the breach in the spine can be surgically closed, the damage to the nerves is permanent. The child's lower limbs could become dysfunctional to varying degrees as a result of the nerve damage. There is always the possibility of misshaped or missing vertebrae, as well as nerve damage, in cases where there is no lesion gift. People with birth defects frequently have a form of impaired learning disorder in addition to physical and mental impairments. Spina bifida has no known cure as the animal tissue cannot be replaced or regenerated. A congenital defect may be treated with surgery, treatment, and medication. Many people who have birth defects utilise aids in addition to braces, crutches, or wheelchairs.¹⁹

Muscular Dystrophy –

The term "Muscular Dystrophy" refers to a group of hereditary illnesses that cause the skeletal or voluntary muscles that control movement to gradually weaken and degenerate. Similar to other involuntary muscles, the heart muscles have some type of hereditary disease. Some types of hereditary disorders have a comparable impact on a person's organs. The most common genetic disease affecting children is called duchene, while the most prevalent disease affecting adult populations is tonicity inherited condition. Some types of hereditary illnesses manifest in early childhood or adolescence, whilst other types may not manifest until a person is an adult or older. People of all ages may be affected by the genetic illness. No particular treatment exists for hereditary disorders of any kind. A physiatrist's services and corrective orthopaedic surgery both have the potential to enhance a person's quality of life.²⁰

Heart

Defects:

A cardiac defect may also form in the early physiological state when the heart is still developing. The functionality and mechanics of a person's heart could be impacted by these flaws. Many children with heart abnormalities are diagnosed by the World Health Organization and exhibit symptoms as soon as they are born. Others don't exhibit signs until early childhood. Some cardiac conditions prevent a person's heart from pumping enough blood to their lungs or other parts of their body, which most likely leads to cardiopathy. Alternative cardiac abnormalities may result in the person's skin appearing pale grey or blue in colour at birth or during infancy; this ailment is known as "Cyanosis."

It's unclear why some people have intrinsic cardiac abnormalities.

Some environmental factors, such as microbial infections that the mother narrows during the earliest phases of the physiological state, are well-known to contribute to the development of innate heart abnormalities. Some drugs may potentially result in inherent heart abnormalities in children when taken by the mother throughout her physiological condition. A number of drugs, including Accutane, bind anti-seizure drugs, Lithium, Trimethoprim, and folacin, have been linked to inherent cardiac problems in children.²¹

Physical and quality impairments have many different aspects.

- People with physical and mental disabilities may have problems with their fine motor skills, movement, and non-locomotor functioning.

Similar to impairments in language, vision, hearing, or other alternative sensory domains, there is the possibility of psychological characteristics, social, and accommodating behaviour deficiencies.

- People with physical and mental disabilities may experience stiffness, jerkiness, and/or loss of muscle strength. They like making learning and daily activities easier.
- People who are physically or mentally impaired may need assistance with quality, transfers, and walking.
- They may experience a sensory activity or psychological characteristic impairment, have a limited range of motion, and be unwilling to undertake movement.
- People with these impairments may experience pain, discomfort, depression, or worry; they may also need to spend a lot of time in bed and may be subject to certain medical limitations.
- They could have comparable contractor or fasciculus deficits.²²

Mobility Impairments Devices-

Wheelchair:

Wheelchairs are devices that can be driven manually or electrically, incorporate a seating system, and are intended to be a replacement for the typical mobility that most people have. Wheelchairs are propelled by an associated hand cycle. People who use wheelchairs and other mobility aids can undertake activities of daily life relating to movement, such as eating, using the restroom, dressing, grooming, and taking a shower. The devices come in a number of forms where the occupant employs electrical controls to regulate the motors and seating control actuators through a joystick, sip-and-puff control, head switches, or other input devices. The gadgets can be driven either by hand or by motors. There are frequently handles or input devices behind the seat for anyone trying to push. People who find it difficult or impossible to walk due to disease, injury, or handicap use wheelchairs. People who have trouble standing up and walking frequently need to use a wheelchair or walker. Using Segway technology or other add-ons like hand bikes or power assistance, wheelchairs can now climb stairs, travel off-road, or move forward thanks to more recent improvements in wheelchair design.

a wheelchair with an attached power accessory²³

Transfer

Patient

Devices-

lift:

Patient transfer equipment often enable caretakers to transfer patients with limited mobility between beds, wheelchairs, toilets, chairs, stretchers, shower benches, cars, swimming pools, and other patient support systems (i.e., radiology, surgical, or examining tables). The most frequently used equipment includes patient lifts (for vertical transfers), transfer benches, stretchers, or convertible chairs (for lateral, supine transfers), sit-to-stand lifts (for transferring patients from one seated position to another, such as from wheelchairs to commodes), air bearing inflatable mattresses (for supine transfers, such as transfers from a gurney to an operating room table), and sliding boards (usually used for transfer from a bed to a wheelchair). OSHA and the American Nursing Association still regard the patient lift, also known as a floor or ceiling-suspended sling lift, which was developed in 1955 and has been widely used since the 1960s, as the most advanced transfer device for highly dependent patients who can't help their carer move them.²⁴

Walkers:

For people with disabilities who require extra assistance to maintain balance or stability while walking, a walker, walking frame, or Rollator may be the right aid. It comprises of a frame that is somewhat wider than the user, about waist high, and about twelve inches deep. There are several sizes of walkers available, such as those for children or large individuals. Height-adjustable walkers are available today. Depending on the user's strength and competence, the front two legs of the walker may or may not have wheels connected. A walker with wheels on the front frequently has glides or caster wheels on the back legs as well.²⁵

Prosthesis:

A device that fills in for a missing part may be referred to as a prosthesis, prosthetic, or prosthetic limb. It belongs to the field of biomechatronics, which studies the use of mechanical devices with human muscle, bone, and neurological systems to supplement or restore control that has been lost due to injury, illness, or defect. Prostheses are frequently used to augment damaged body parts, replace missing or absent from birth (congenital) body parts, and replace lost or injured body parts. Implant valves are frequently used inside the body, but artificial hearts and lungs are less frequently used but are still actively being developed. Additionally, hearing aids, artificial eyes, palatal obturators, gastric bands, and dentures will all be regarded as prostheses. Prostheses are not orthoses in the strictest sense, yet under certain conditions they might end up serving some or all of the same functional purposes as one. In a sense, prostheses are the full finished product. For instance, a C-Leg knee is merely a prosthetic component and not a complete prosthesis. The entire prosthesis would include the connection system to the residual limb, which is typically a "socket," as well as every piece of attachment hardware up to and including the terminal device. As terminology is frequently used interchangeably, keep this in mind.

Prosthetic and orthotic are adjectives commonly used to describe artificial joints, such as a prosthetic knee. Both allied health professions are commonly referred to as "prosthetics" and "orthotics." The duties of an occupational therapist in the field of prosthetics include rehabilitation, instruction, and assessments. Orientation to prosthetic

components and terminology, donning and doffing, wearing schedule, and how to care for the residual limb and the prosthesis are all included in prosthetic training.²⁶

Exoskeletons

A wearable mobile device known as a powered exoskeleton could be powered by a system of electrical motors, pneumatics, levers, hydraulics, or a combination of technologies that give limb movement greater strength and endurance. Its construction seeks to provide back support, take into account the user's motion, and communicate with the motors controlling the gears. The exoskeleton helps with movement for lifting and gripping big objects while reducing back strain. It also supports the shoulder, waist, and thigh. Adaptive Positioning and Seating Specialized equipment is frequently required for people with balance and motor function issues to sit down or stand up securely. This equipment is frequently tailored for use in particular environments, like a classroom or institution. In sitting arrangements, positioning is typically crucial to ensure that users' body pressure is distributed equally without impairing movement in an undesirable way.

Positioning aids are made to make it safer for people to stand square and support their weight on their legs. These standers are typically divided into two groups based on how they supported the occupant's stance. Prone standers typically have a tray in front of them and spread their weight forward. They are therefore advantageous for users who are actively attempting to complete a task. In situations where the user has less mobility or is recovering from an injury, supine standers are useful because they distribute the user's weight to the back.²⁷

Discussion –

The International Organization for Standardization (ISO) defines assistive products more broadly as any product that is specifically produced or widely accessible and that is used by or for people with disabilities in order to participate, to protect, aid, train, level up, or replace frame functions/systems and activities, or to avoid impairments, activity boundaries, or participation restrictions. In particular, children and persons through process therapy who anticipate future growth may outgrow their assistive devices may find the expenditures of purchase and maintenance to be significant. There may be a few assistive devices that can be paid for with the help of your government's healthcare or with the help of private insurers, depending on your country's healthcare system. The Standard Rules on the Equalization of Opportunities for Persons with Disabilities also call on States to support the creation, manufacturing, distribution, and upkeep of assistive devices and equipment as well as the spread of knowledge about them.

Conclusion -

Improvements made in the assistive era led to the refinement of a present assistive era in clever ways so that its capabilities integrated better with the person's skills. These improvements are being made on a utility-through-means-of basis by developing advanced technology mechanics (such as knee-ankle-foot orthoses, kinetic

manipulation of prosthetic limbs), advanced user interfaces (such as tongue or whole-frame controllers, electrodes implanted in valuable and peripheral nervous systems), and automating goal-control functions in a way that blends the machine's assistance with the natural capabilities. Better integrated control structures reduce user strain, enabling more nuanced manipulation of particularly cutting-edge prosthetics or enabling those with the most severe physical limitations. independent mobility powered wheelchairs.



References-

1. American Speech-Language-Hearing Association. (2005). "Roles and Responsibilities of Speech-Language Pathologists With Respect to Augmentative and Alternative Communication: Position Statement". Archived from the original on February 13, 2009. Retrieved January 23, 2009.
2. DeCoste, Denise C. (1997). "Chapter 10: Introduction to Augmentative and Alternative Communication Systems". In Glennen, Sharon; DeCoste, Denise C. (eds.). Handbook Of Augmentative And Alternative Communication. San Diego, CA: Singular Publishing Group. ISBN 978-1-56593-684-3.
3. Schlosser, R. W.; Wendt, O. (2008). "Effects of augmentative and alternative communication intervention on speech production in children with autism: a systematic review". American Journal of Speech-Language Pathology. 17 (3): 212–230. doi:10.1044/1058-0360(2008/021). PMID 18663107.
4. Beukelman, David R.; Mirenda, Pat (2005). Augmentative & alternative communication: supporting children & adults with complex communication needs (3rd ed.). Paul H. Brookes Publishing Company. ISBN 978-1-55766-684-0.

5. Galvão Filho, T. (2009). "Tecnologia Assistiva para uma Escola Inclusiva: apropriação, demandas e perspectivas. Tese (Doutorado em Educação) – Faculdade de Educação, Universidade Federal da Bahia, Salvador, Brazil".
6. K. L. Rust and R. O. Smith, "Assistive technology in the measurement of rehabilitation and health outcomes: a review and analysis of instruments," *American Journal of Physical Medicine and Rehabilitation*, vol. 84, no. 10, pp. 780–793, 2005.
7. N. B. Oldridge, "Outcomes measurement: health-related quality of life," *Assistive Technology*, vol. 8, no. 2, pp. 82–93, 1996.
8. G. Hawthorne, J. Richardson, and R. Osborne, "The Assessment of Quality of Life (AQoL) instrument: a psychometric measure of health-related quality of life," *Quality of Life Research*, vol. 8, no. 3, pp. 209–224, 1999.
9. T. Shakespeare, *Disability Rights and Wrongs*, Routledge, New York, NY, USA, 2006.
10. A. Cook and S. Hussey, Eds., *Assistive Technologies: Principles and Practice*, vol. 3, Mosby Elsevier, St. Louis, Mo, USA, 2008.
11. N. Layton and E. Wilson, "Re-conceptualizing disability and assistive technology: Australian consumers driving policy change," *Technology and Disability*, vol. 21, no. 4, pp. 135–141, 2010.
12. F. Heywood and L. Turner, *Better Outcomes, Lower Costs: Implications for Health and Social Care Budgets of Investment in Housing Adaptations, Improvements and Equipment: a Review of the Evidence*, University of Bristol Office for Disability Issues, Bristol, UK, 2007.
- [24] N. Layton, E. Wilson, S. Colgan, M. Moodie, and R. Carter, *The Equipping Inclusion Studies: Assistive Technology Use and Outcomes in Victoria*, Deakin University, Melbourne, Australia, 2010.
13. "Assistive Technology Act of 1998 | Section508.gov". Section508.gov. Archived from the original on March 29, 2016. Retrieved April 4, 2016.
- 14.[^] <https://www.aota.org/Publications-News/AmericanJournalOfOccupationalTherapy/access-framework-domain-otpf-download.aspx>
15. Journal: Disabled World. Language: English (U.S.). Author: Disabled World. Revised Publication Date: 2019-01-17. Title: Physical and Mobility Impairment Information, Source: <https://www.disabled-world.com/disability/types/mobility/> Mobility Impairments. Retrieved 2021-06-02, from <https://www.disabled-world.com/disability/types/mobility/> - Reference: DW#264-17.172.98-6c.

16. M. P. J. M. Dijkers, "Correlates of life satisfaction among persons with spinal cord injury," *Arch. Phys. Med. Rehabil.*, vol. 80, no. 8, pp. 867–876, 1999.
17. N. Sezer, "Chronic complications of spinal cord injury," *World J. Orthop.*, vol. 6, no. 1, p. 24, 2015.
18. World Health Organization, "Towards a Common Language for Functioning, Disability and Health ICF," *Int. Classif.*, vol. 1149, pp. 1–22, 2002.
19. J. Wee and R. Lysaght, "Factors affecting measures of activities and participation in persons with mobility impairment," *Disabil. Rehabil.*, vol. 31, no. 20, pp. 1633–1642, 2009.
20. J. A. Lenker, F. Harris, M. Taugher, and R. O. Smith, "Consumer perspectives on assistive technology outcomes," *Disabil. Rehabil. Assist. Technol.*, vol. 8, no. 5, pp. 373–380, 2013.
21. de G. I. Cardol M, de Haan RJ, van den Bos GA, de Jong BA, "Impact on Participation and Autonomy-E (IPA)," *Clin. Rehabil.* vol. 13, pp. 411–419, 1999.
22. D. B. Gray, H. H. Hollingsworth, S. L. Stark, and K. A. Morgan, "Participation Survey / Mobility," pp. 1–13, 2006.
23. J. W. Jutai and H. Day, "Psychosocial Impact of Assistive Devices Scale (PIADS)," *Technol. Disabil.*, no. JANUARY 2002, pp. 107–111, 2002.
24. A. Martins, J. Pinheiro, B. Farias, and J. Jutai, "Psychosocial Impact of Assistive Technologies for Mobility and Their Implications for Active Ageing," *Technologies*, vol. 4, no. 3, p. 28, 2016.
25. R. Fomiatti, J. Richmond, L. Moir, and J. Millstead, "A Systematic Review of the Impact of Powered Mobility Devices on Older Adults' Activity Engagement," *Phys. Occup. Ther. Geriatr.*, vol. 31, no. 4, pp. 297–309, 2013.
26. M. J. Scherer and R. Glueckauf, "Assessing the Benefits of Assistive Technologies for Activities and Participation," *Rehabil. Psychol.*, vol. 50, no. 2, pp. 132–141, 2005.
27. Hermansson, Liselotte N.; Turner, Kristi (October 2017). "Occupational Therapy for Prosthetic Rehabilitation in Adults with Acquired Upper-Limb Loss: Body-Powered and Myoelectric Control Systems". *JPO Journal of Prosthetics and Orthotics*. 29 (4S): P45–P50. doi:10.1097/JPO.000000000000154. ISSN 1040-8800. S2CID 79986334.