



# EFFECT OF NaCl SALINITY ON INORGANIC CONSTITUENTS OF MULBERRY. VARIETY M5

LATADEVI S KARIKAL

SHARNBASVESHVAR COLLEGE OF SCIENCE, KALABURAGI. KARNATAKA.

Email: [latadevikarikal@gmail.com](mailto:latadevikarikal@gmail.com)

## Abstract

Mineral elements are important constituents of protoplasm and cell wall. Mineral salts dissolved in the cell sap partially influence the osmotic pressure of the cell. Imbalance of mineral ions under saline condition generally results into poor crop growth and lesser productivity. The metabolic reactions also get affected due to disturbed ionic concentration. The minerals sodium and chloride are listed among essential trace elements, sodium regulates the transport of amino acids to the nucleus, whereas chloride plays an important role in the transfer of electrons from water to chlorophyll. Even though the higher levels of these elements alter the plant metabolism and reduce the yield. The other macronutrients such as Potassium, Calcium, Magnesium, Iron and micronutrients are essential for various metabolic activities. If all the necessary nutrients available in required amount, then the growth and yield of the plant will be optimum, if any disturbance in the nutrition disturbs the entire metabolic activity of the plant. The present investigation is to analyse the levels of various mineral nutrients in the different parts of NaCl treated Mulberry plants because salinity is known to influence the ion uptake through change in the osmotic potential and ion interaction at root zone.

Key words: NaCl salinity, Mineral ions, Inorganic components, Mulberry.

## Introduction:

Soil salinity is one of the major abiotic stresses throughout the world. According to present statistics 1125 million hectares of land is affected by salinity in the world, of which 76.6 million hectares are by anthropogenic activities (Oldmen.,1991; Mashali,1995; Shahid et al.,2018). Every year a large area is getting salinized and probably by 2050 around 50% of arable land would be salt affected. In India saline soil occupy 44% area covering 12 states and one union territory. According to Singh (1980) the introduction of irrigation projects has been the main cause of soil salinity in many countries. At present Agriculture sector has two threats that is population explosion and unavailability of cultivable land (Shahbaz and Ashraf 2013). In this context ICAR, Central soil salinity Research Institute and many States Agricultural Universities are engaged in the studies related to salt affected land and developing several innovative technologies in the country. Among them are screening and cultivation of salt tolerant crop varieties. Because Soil salinity reduce net cultivable area and also

affects productivity and quality of the crop (Munns 2005, Jamil et al 2011). An estimate suggested the global economic losses are about 11.4 billion USD in an irrigated land and 1.2 billion USD in non-irrigated land per year (Ghassemi et al 1995) due to soil salinity. Salinity affects almost all parts of the plant and also growth and reproduction. They face osmotic and nutrient stress. The agriculture crops exhibit many complex responses and interaction of physiological, morphological and biochemical processes which ultimately leads to low quality and production ((Ghassemi et al 1995)). In addition to this soil salinization also affects physicochemical properties of the affected area which leads to ecological imbalances (Liu et al 2019). Salinity stress leads to accumulation of various types of metabolites, these metabolites helps to overcome the stress in plants by developing tolerance (Gomez et al 2019; Pyne et. al 2019). The present study aimed to understand the influence of soil salinity on various organic metabolites under increasing concentration of salinity in two varieties of Mulberry.

Mulberry (*Morus alba. L*) is a deciduous woody tree, which has great economic importance and cultivated as commercial crop, the leaves are used to feed silkworms *Bombyx mori. L* to get superior quality of silk. The tender parts of the plant are used as fodder and also the ripe fruits are nutritious and edible. Mulberry is cultivated as both rainfed and irrigated crop. Mulberry cultivation is associated with rearing of silkworms known as sericulture. Sericulture is practised by small and marginal land holders; it gives economy and employment for rural women. The present study aimed to investigate effects of salinity on the inorganic constituents such as some macro and micro elements which may help to know the effect of increased sodium chloride on mineral ion concentration in different parts of the plants. Which may help to screen new salinity resistant varieties, this may help to utilize salt affected lands.

### **Materials and Methods**

During present investigation mulberry cultivar M5 was selected. Mulberry cuttings were obtained from Regional Sericulture Research Station, Kadaganchi. Kalaburagi. Healthy cuttings with 15-18 cm length with 3-4 axillary buds were selected. The selected cuttings were immersed in 10ppm solution of NAA for 20-30 min to initiate rooting. The cuttings were planted in earthen pots (22 x 20 CMS) filled with equal quantities of (4 kg/pot) garden soil and sand. The saplings were established. The well-established three months old healthy and uniform saplings were transplanted to experimental pots, each pot with 4 saplings, after recovery of saplings due to transplantation shocks the pots were irrigated with 1 lit of tap water to control, 1lit of 0.25 M, 0.5 M, and 1.0 M NaCl solution to other experimental pots respectively. Three replications for each treatment were maintained. The salt treatment was given by the method of Black (1956) The plants were harvested and collected after 45 days for analysis of organic metabolites. EC and pH of soil extract was measured up to 45 days. After 45 days the plants were harvested, the different plant parts like Roots, Stems and leaves were collected and used for estimation of mineral ions. The acid digestion was carried as per the method of AOAC hand book (Horwitz 1975)

Sodium and Potassium elements were determined by using Flame photometer (Jackson 1956) Calcium was estimated by titrimetric EDTA method (Richards 1954). The minerals like Fe, Mn, Zn, Cu and Mg were analysed by Atomic spectrophotometer, at the regional sophisticated instrumentation center, IIT Bombay. Chloride was estimated by the method of Volhard (1956).

**Results:**

**Table – 1**  
**Effect of NaCl on Mineral Components of Leaf**

NaCl(M)	Na	K	Ca	Mg	Fe	Mn	Zn	Cu	Cl	Na/K	Na/Ca	K/Ca
Control	28.0 ±0.06	14.1 ±0.04	78.1 ±0.13	8.7 ±0.03	1.19 ±0.09	0.11 ±0.008	0.012 ±0.003	0.02 ±0.002	7.2 ±0.03	0.019	0.35	18.21
0.25	40.0 ±0.05	13.4 ±0.06	106.1 ±0.14	11.3 ±0.03	1.91 ±0.03	0.15 ±0.002	----	0.02 ±0.002	11.6 ±0.05	0.029	0.037	12.71
0.5	56.0 ±0.06	18.0 ±0.05	128.2 ±0.25	18.3 ±0.04	2.42 ±0.03	0.14 ±0.005	----	0.02 ±0.003	29.9 ±0.18	0.031	0.43	14.06
1.0	150.0 ±0.02	24.0 ±0.08	108.1 ±0.12	16.8 ±0.05	0.07 ±0.002	0.20 ±0.01	0.015 ±0.003	0.03 ±0.003	72.3 ±0.03	0.061	1.38	22.66

Values are expressed as mg/g of dry weight

Values are significant at 1% level

In table 1 shows various inorganic contents of leaf. The sodium and chloride as usual increased with increase in salinity level. The potassium accumulation was stimulated at 0.25M level but gradually decreased with the increase in salinity level. Calcium accumulation increased upto 0.5 m level but lowered at 1m level, Magnesium accumulation was lower at 0.25 M level but increased at higher NaCl level iron content remained same in control and 0.25 level whereas at higher concentration iron level increased and more or less remained same. Manganese was low at lower concentration but high at higher concentration. Salinity did not affect uptake of Zinc and copper to a large extent.

Na/K and Na/Ca ratio showed increasing tendency but K/Ca ratio increased at 0.25M and 1M but declined at 0.5 m NaCl level.

**Table – 2**  
**Effect of NaCl on Mineral Components of Stem**

NaCl (M)	Na	K	Ca	Mg	Fe	Mn	Zn	Cu	Cl	Na/K	Na/Ca	K/Ca
Control	47.6 ±0.03	14.4 ±0.05	66.1 ±0.14	10.9 ±0.08	0.87 ±0.05	0.07 ±0.008	--	0.02 ±0.001	1.2 ±0.003	0.033	0.72	21.85
0.25	70.8 ±0.06	12.5 ±0.08	60.0 ±0.07	12.2 ±0.02	0.92 ±0.03	0.08 ±0.009	---	0.01 ±0.001	2.0 ±0.10	0.057	1.19	20.70
0.5	114.2 ±0.06	14.1 ±0.05	27.2 ±0.12	12.6 ±0.12	1.94 ±0.06	0.09 ±0.005	---	0.02 ±0.003	16.3 ±0.06	0.080	4.17	51.92
1.0	156.1 ±0.10	88.4 ±0.01	38.1 ±0.06	81.9 ±0.06	2.47 ±0.11	0.08 ±0.007	----	0.02 ±0.005	18.8 ±0.04	0.017	4.12	32.10

Values are expressed as mg/g of dry weight

Values are significant at 1% level

-- Values are negligible

The table -2 shows the results of effects of NaCl salinity on mineral components. The sodium and chloride accumulated in the stem tissues, accumulation increased with increase in salinity level. potassium accumulation decreased slightly in 0.25 M level ,whereas remained same as control but at 1.0 M rate of accumulation increased suddenly. Calcium has shown decreasing tendency and highly affected by salinity. Magnesium increased upto second level but at 1M eight times more than control. Iron deposition increased with increase in salt concentration. Whereas Mn and copper did not affect but Zn is found in traces.




**Table – 3**  
**Effect of NaCl on Mineral Components of Root**

NaCl( M)	Na	K	Ca	Mg	Fe	Mn	Zn	Cu	Cl	Na/K	Na/Ca	K/Ca
Control	35.2 ±0.03	100.1 ±0.11	38.0 ±0.05	6.9 ±0.05	31.3 ±0.10	0.16 ±0.03	0.06 ±0.004	0.07 ±0.003	5.1 ±0.07	0.35	0.92	2.63
0.25	30.4 ±0.02	103.1 ±0.03	32.0 ±0.07	5.3 ±0.04	26.7 ±0.02	0.14 ±0.01	0.04 ±0.006	0.06 ±0.005	7.2 ±0.03	0.29	0.95	3.22
0.5	28.1 ±0.02	57.8 ±0.05	36.0 ±0.11	7.6 ±0.03	43.5 ±0.03	0.16 ±0.02	0.03 ±0.003	0.06 ±0.01	17.1 ±0.05	0.48	0.78	1.60
1.0	27.1 ±0.04	92.64 ±0.05	66.0 ±0.08	7.5 ±0.02	27.3 ±0.04	0.12 ±0.01	0.02 ±0.002	0.04 ±0.003	5.1 0.05	0.29	0.41	1.40

Values are expressed as mg/g of dry weight

Values are significant at 1% level

The table -3 shows the results of mineral ion deposition in root under various levels of NaCl concentration. Na uptake decreased with increased salt concentration. Whereas Cl increased up to 0.5 M but declined suddenly at 1M concentration. The potassium accumulation decreased at 0.5M and 1M concentration. Ca accumulation was decreased upto0.5 NaCl but enhanced at 1Msalt concentration.



**Discussion:**

Salinity can damage the plant through osmotic effects of ions and by disturbing uptake of essential nutrients. According to Greenway and Munns (1980) salt sensitivity in non-halophytes may results from, injury caused by ions which are absorbed by the cell and are not compartmentalized. Disturbed inorganic nutrition is one of the factors responsible for the suppression of plant growth in saline or alkaline soil (Singh and Singh,1985).

According to Pitman (1975) ion distribution within plant is dependent on two main factors namely uptake and its radical and its radical and long distance transport NaCl stress has been related to increase contents of Na, Cl or both ions in the cytosol (Lauchil and Wiencke,1979) and to a decreased internal availability of nutrients arising from a lower uptake rate and the competition between nutrient ions and Na or Cl (Black,1956;Greenway,1962).The ionic imbalances affects plant metabolism through repression or induction (Hason-Porath and Poljakoff Mayber,1969;Boucaud and Billard, 1979)

The maintenance of internal positive turgor potential of plants exposed to saline conditions is an important factor for maintaining growth. This accomplished by uptake of ions, chiefly K, Na and Cl as well as by synthesizing organic metabolites (Yeo,1983). Potassium is the most abundant cat ion in the cytoplasm and in glycophytes plays an important role in osmotic adjustment (Marschner,1986).and maintenance of photosynthetic activity ( Bal et al.,1978; Behboudian and Anderson,1990).Mg is an integral constituent of chlorophyll pigments and Ca is required essentially for maintenance of physicochemical nature, permeability and structural configuration of membranes(Greenway and Munns,1980) The role of Ca as second messenger in plants or of calcium containing proteins has been well documented in literatures.

Large differences in ion concentrations occurs between species. The increase in Na in salt tolerant species grown at 100mM is generally associated with a decrease in K. In contrast salt sensitive species which includes Na from their shoots usually increase in K by 20 to 30% Greenway et al.,1981)There are also differences in ion concentration between different parts of the same plant.

Generally the different plant parts like leaf, stem and root showed a differential rate of uptake of both macro and micro nutrients and also the rate of accumulation or exclusion of minerals show varietal differences. Salinity generally decreased mineral transport especially Fe, Mn ,Mg and Cu from seed to seedlings, except for potassium. Transport Fe, Mg and Ca to the aerial part was also markedly reduced in sunflower (Sanchez Raya and Delgado,1996) The response of moderately tolerant genotypes had higher concentration of Na and Cl in the shoot tissue and also showed lower K/Na ratio under stress as compared to tolerant genotypes. The ratio of K/Na was decreased progressively on salinization.

Kulkarni and Karadage (1991) have reported that Ca suppressed Na uptake and probably contributed to the ionic balance under stress conditions. Uptake and distribution of Mg was favourable, Fe and Mn were accumulated to the toxic level at 100mM NaCl in moth bean, Mn uptake tended to be higher. Ruiz et al., (1997) have found that a significant effect on the leaf concentration of Cl, Na,K, Ca ,Mg ,P ,Fe ,Mn and Zn in citrus after 60 days of salinization. Xiaohang Weng et.al (2024) have reported salt stress shifted the calcium absorption focus from leaves to trunks. Exogenous calcium mitigation on mulberry stoichiometry under salt stress shows initial decrease then increase, the N: P ratio showed that salt treatment could change the P stress to N stress in mulberry seedlings. Meanwhile, there was a significant positive correlation between the biomass of each organ with C: P and N: P ( $p < 0.01$ ,  $0.001$ ), but a extremely significant negative correlation between the biomass with others stoichiometric ratios ( $p < 0.001$ )

In contrast to this there are several reports showing decreased ratio of Na/K and Na/Ca. Datta et al.,(1991) have reported decreased K/Na ratio in three tropical foliage crops under salinity in Maiza , Sudan grass ,Teosinte). Lahiri et al (1996) has showed the moderately tolerant genotypes of cluster bean had lower K:Na ratio under stress. The ratio of K/Na was decreased progressively on salinization in sunflower plants by Gadallah (1996). The soil sodicity decreased K/Na and Ca/Na ratio in plants in tissues considerably (Porcelli et al .,1995). El-Agrodi (1991) has found the K/Na in plant decreased with increasing soil salinity in sorghum.The decreased K/Na in seven flax cultivars under saline treatment was reported by Beke and Graham (1995). Ritwik Acharya et.al (2024) reporte exposure to NaCl increased the Na concentration in the plant tissues. Na concentrations were affected in the shoots after only 3 days under the 0.1% NaCl treatment. Similar trends were observed for changes in the Na concentrations in taproots and lateral roots. In leaves, Na concentrations began to increase significantly under the 0.2% NaCl treatment. After 21 days, the Na content increased only in plants subjected to the 0.3% NaCl treatment. K concentrations in leaves were unaffected by the NaCl treatments, but increased in other organs after 3 days of NaCl treatment and decreased after 21 days of treatment. Mg concentrations increased in both lateral roots and taproots after 21 days of NaCl treatment, but decreased in leaves and shoots. Ca concentrations increased in response to NaCl treatment in shoots, taproots, and lateral roots after 3 days, but remained unchanged in leaves by NaCl concentrations less than 0.5%. After 21 days, the Ca concentrations in

taproots and lateral roots increased in response to the high salinity (0.5% NaCl) treatment. The K:Na, Mg:Na, and Ca:Na ratios decreased with increasing salinity levels.

In the present investigation accumulation of mineral constitution under salt stress in mulberry, variety M5 showed that the Na and Cl accumulation was more in stem and leaves than in roots. It seems that root has excluded the accumulation of toxic ions. It has effectively translocated the toxic ions to stem and leaves. More amount of Na and Cl was found in stem than the leaf. This shows that stem has accumulated more Na and Cl. At the same time the Ca accumulation in stem and leaf increased and decreased in root. Whereas roots have accumulated more K and the K content was less in stem and leaf. This variable distribution of mineral ions namely Na, K, Ca, and Cl seems to have role in the salt tolerance metabolism of mulberry. Roots have developed higher tolerance than aerial parts.

**Conclusion:** The present investigation may help to know the uptake and accumulation of various minerals in different parts of the saline treated plants. Which may help to screen salt tolerant varieties which in turn helps to undertake Mulberry cultivation in the salt affected lands.

### References:

- 1] Oldeman LR, Hakkeling RTA, Sombroek W G (1991). World map of the status of human-induced soil degradation. An explanatory note. Second revised edition. International Soil Reference and Information Centre (ISRIC), Wageningen, 35 pp
- 2] Mashali AM (1995) Integrated soil management for sustainable use of salt-affected soils and network activities. Proceedings of the international workshop on integrated soil management for sustainable use of salt-affected soils. 6–10 November 1995. Bureau of Soils and Water Management, Manila, Philippines, pp 55–75
- 3] Shahid, S.A., Zaman, M. and Heng, L. (2018) Soil Salinity: Historical Perspectives and a World Overview of the Problem. Eds., Guideline for Salinity Assessment, Mitigation and Adaptation Using Nuclear and Related Techniques. Springer International Publishing, Cham.
- 4] Singh (1980). Growth of Guava Plants (*Psidium gajuava* L.) as affected by salinity Indian Journal of Horticulture - indianjournals.com.
- 5] Shahbaz and Ashraf (2013). Improving salinity tolerance in cereals, Crit. Rev. Plant Sci.32.pp237-249
- 6] Munns (2005). Genes and salt tolerance: bringing them together. New Phytol.167, pp. 645-663.
- 7] A.Jamil,S.Riaz,M.Ashraf and M.R.Foolad (2011). Pages 435-458 | Published online <https://doi.org/10.1080/07352689.2011.605739>
- 8] Ghassemi F, Jakeman AJ, Nix HA (1995) Salinisation of land and water resources: human causes, extent, management and case studies. CABI Publishing, Wallingford 526 pp.
- 9] Black R F (1956). Effect of NaCl in water culture on the ion uptake and growth of *Atriplex hastata* L. Aust. J. Biol. Sci. 9:67-80
- 10] Horwitz Willam (1975). (Ed) Official methods of analysis of the Association of official analytical chemistry, Ph. Washington. D.C. pp.34.
- 11] Greenway, H and Munns, R (1980). Mechanism of salt tolerance in nonhalophytes. Ann. Rev. Plant. Physiol, 31:149-190

- 12] Singh and Singh (1995). Sodicty inducedchangesin saccharides,free proline and Protein Content of different Pea genotypes.Ind.J.Plant Physiol,38(2):196-200
- 13] Pitman, M. G (1975).In:Ion Transport in plant cell and tissue.Eds.D.A.Barder and J.L.Hall,North Holland
- 14] Lauchil,A and Wiencke,J (1979) Studies on growth and distribution of Na,K and Cl in Soyabean Varieties Differing in Salt tolerance.Z.Phanzenernaehr Bodenk.142:3-13
- 15] Black,R.F(1956) Effect of NaCl in water culture on the ion uptake and growth of *Atriplex hastata*. *L.Aust.J.Biol.Sci.*9:67-80
- 16] Greenway,H (1962) Plant response to saline substrate. Growth and ion uptake of several varieties of *Hordeum vulgare* during and after sodium chloride treatment.*Aust.J.Biol.Sci.*15:16-38
- 17] Hason-Porath,E and Poljakoff Mayber, A(1969). The effect of salinity on malic dehydrogenase of Pea roots.*Plant. Physiol.*,44: 1031-1034
- 18] Yeo,A.R (1983). Salinity resistance; physiologies and prices.*Physiol Plant.*58; 214- 222
- 19] Marschner,H (1986). Mineral nutrition of higher plants. Academic press, London, Orlando, Sandigo, New York, Austin, Boston, Sydney, Tokya, Toronto.
- 20] Bal et al (1978). Grain yield analysis and N.R activity in rice. *Ann.Rep. Centr.Soil salinity. Inst.karnal, India.* 69-71.
- 21] Behboudian M.H and Anderson, D. R (1990). Effect of potassium deficiency on water relations and photosynthesis of tomato plant. *Plant and soil.* 127; 137-139.
- 22] Greenway, H and Munns, R (1980). Mechanism of salt tolerance in non-halophytes. *Ann.Rev.Plant. Physiol.* 31; 141-190
- 23] Greenway,H Munns,R and Kirst,G.O (1981). Halophytes, higher plants and algae. *Encyclopedia of plant physiology; physiological plant ecology*, ed.O.L.Lange, C.B.Osmond. P.S.Nobel, H.Zigler. in preparation.
- 24] Sanchez-Raya, A.J and Delgado,I.C (1996). Mineral nutrient transport by sunflower seedlings grown under saline conditions (Nacl). *J.Plant. Nutr.*19 (10-11); 1463-1475.
- 25] Kulkarni,H and Karadage,B.A (1991). Growth and mineral nutrition of moth bean(*Phaseolus aconotifolius*.Jacq) under saline conditions.*Ind.J.Plant Physiol.*34(1); 14-24.
- 26] Ruiz.D., Martinez and Cerda A (1997). Citrus response to salinity; growth and nutrient uptake. *Tree Physiol.* 17(3); 141-150.
- 27] Xiaohang Weng, Hui Li, Yongbin Zhou, Chengshuai Ren, Yan Huo, Songzhu Zhang & Liying Liu (2024).Exogenous Calcium can Promote Nutrient Uptake and Stoichiometric Balance of Mulberry Seedlings Under Salt Stress. *Journal of Soil Science and Plant Nutrition*
- 28] Datta,.K.S., Asok kumar., Verma ,S.K and Angrish,R (1996). Delimitation of salt tolerance in barlie on the basis of water relations and ion distrinution under chloride and sulphate salinity. *Ind.J.Plant Physiol.* 33(3); 253-256.

- 29] Gadallah,M.A.A. (1996). Abscissic acid temperature and salinity interactions on growth and some mineral elements in Carthamus plants. *Plant growth regulation*. 20(3); 225-236.
- 30] Porcelli,C.A.,Gutierrez Boem,F.H and levado,R.S (1995) The K/Na and Ca/Na. ratios and rapeseed yield under soil salinity and sodicity. *Planta and Soil*. 175(2); 251-255.
- 31] El-Agrodi (1991)-M.W., EL- Sirafy, Z.M., Taha, A.A and EL- Hadidi, E.M (1991). Sorghum growth and nutrients content in relation to soil salinity and nitrogen fertilisation. *Egyptian. J.Soil. Sci.* 31(2); 181-194.
- 32] Beke. G.J and Graham,D.P (1995). Growth and chemical composition of flax cultivors on artificially salinized soil. *Canadian.J.P.SCI.* 75(1); 159-162.
- 33] Ritwik Acharyaa , Debnirmalya Gangopadhyaya deb, Ganguly,Phalguni Bhattacharyya B, Amitava Ghosh C. (2024). Evaluation of salt-tolerant germplasm of mulberry (*Morus L.*) through in vitro and field experiments under different salinity stresses. *Volume 10, Issue 16 e35868 August 30, 2024 Open access*

