



Wetland Birds as Bioindicators of Ecological Health: Conservation Significance of Ujani Reservoir, Maharashtra, India

Ms. P.D Rahate¹ and Dr. P. V. Patil²

¹Assistant Teacher, Shankarrao Mohite Mahavidyalya, Akluj

²Associate Professor, K.S.K College, Beed

Abstract

Wetlands are sensitive ecosystems and changes in water availability, habitat condition, pollution and human activities can affect their biological communities. Birds are closely associated with wetland habitats and can provide useful information about changes in ecological conditions. Ujani Reservoir in Solapur District, Maharashtra, supports a variety of resident and migratory wetland birds and contains different habitats such as open water, shallow water, mudflats, marshes and surrounding agricultural areas. The present paper discusses the use of wetland birds as indicators of ecological condition and the conservation importance of Ujani Reservoir based on published information. Earlier studies have reported considerable aquatic-bird diversity from wetlands associated with the reservoir, including resident, migratory and resident-migratory species. Differences in bird distribution and habitat use have been related to water availability, food resources, habitat conditions and human interference. Sand mining, fishing, livestock grazing, agricultural runoff, tourism, vehicular movement, habitat encroachment, water extraction and solid-waste disposal are among the pressures reported around the reservoir. These activities may affect habitats used by wetland birds. Regular monitoring of bird communities along with habitat conditions and human activities can help in assessing changes in the wetland. Protection of important bird habitats and reduction of unnecessary disturbance are therefore important for maintaining the ecological value of Ujani Reservoir.

Keywords: Wetland Birds, Bioindicators, Biomonitoring, Ujani Reservoir, Conservation, Habitat Disturbance, Migratory Birds

Introduction

Wetlands provide important habitats for many plants and animals. For birds, they provide feeding areas, breeding grounds, roosting sites and places for seasonal movements. Changes in water level, vegetation, food availability and habitat condition can influence the birds that use these areas.

Birds are useful in ecological studies because different species depend on different habitats and food resources. Changes in species richness, abundance, community composition and seasonal occurrence can therefore provide information about changes in habitat condition. Wetland birds are consequently useful for monitoring wetland ecosystems.

Ujani Reservoir is situated on the Bhima River in Solapur District, Maharashtra. The reservoir and surrounding areas contain open water, shallow-water areas, mudflats, marshes, agricultural fields, grasslands and islands. The variety of habitats allows birds with different feeding and habitat requirements to occur in the same wetland system.

Published studies have reported 81 aquatic bird species from 15 wetland sites and 38 wader species from five sites associated with Ujani Reservoir. The bird community includes resident, migratory and resident-migratory species.

The reservoir is also exposed to several human activities, including sand mining, fishing, livestock grazing, agricultural runoff, tourism, vehicular movement, habitat encroachment, water extraction and waste disposal. These pressures make regular monitoring important.

The aim of this paper is to discuss the importance of wetland birds as bioindicators and the conservation significance of Ujani Reservoir using available published information.

Materials and Methods

i.Study area

Ujani Reservoir is located on the Bhima River in Solapur District, Maharashtra. The reservoir and its surrounding wetlands contain different aquatic and terrestrial habitats used by wetland birds. The locations included in the available synopsis are Palasdeo, Bhigwan, Diksal, Kondhar Chincholi, Katraj, Takali and Old Railway Bridge.

ii.Sources of information

Information was obtained from published studies on wetland birds of Ujani Reservoir and from the available research synopsis. Bird diversity, migratory status, habitat use and reported human disturbances were considered.

The available methodology identifies species richness, abundance, community composition, seasonal variation, habitat utilization, migratory assemblages and response to anthropogenic disturbance as useful parameters for biomonitoring.

Since the original observation-level data are not available, no new diversity index, abundance estimate or statistical relationship has been calculated.

iii.Wetland birds as bioindicators

Wetland birds depend on several components of their habitat. A change in water level, food availability, vegetation or habitat structure can affect the species using a particular wetland.

Species richness and abundance are simple measures that can be followed over time. Changes in community composition and seasonal occurrence can also help identify changes in habitat conditions. The presence of migratory birds can provide additional information because many migrants depend on suitable feeding and resting habitats during particular seasons.

The available synopsis suggests species richness, abundance, community composition, seasonal variation, habitat use, migratory assemblages and response to human disturbance as useful components of wetland-bird biomonitoring.

However, birds can be used effectively as indicators only when monitoring is carried out regularly and with similar methods. A single species list cannot show whether the condition of a wetland is improving or deteriorating. Repeated observations over different seasons and years are more useful for this purpose.

iv. Ecological significance of Ujani Reservoir

The published records show that Ujani Reservoir supports a diverse bird community. One study recorded 81 aquatic bird species belonging to 59 genera, 11 orders and 23 families from 15 wetland sites. Another recorded 38 wader species belonging to 14 families from five sites.

The aquatic-bird study recorded 26 resident, 19 migratory and 36 resident-migratory species. This mixture of birds indicates that the reservoir provides suitable habitat during different periods of the year.

Birds also use different parts of the wetland. The published study recorded divers, swimmers, large waders, small waders and aerial foragers. The availability of different habitats therefore appears to be important for maintaining the bird community.

v. Human pressures and their possible effects

The ecological value of Ujani Reservoir is closely connected with the condition of its wetland habitats. Several activities reported around the reservoir can affect these habitats.

Human activity	Possible effect
Sand mining	Alteration of wetland margins
Fishing	Disturbance near feeding and resting sites
Livestock grazing	Habitat disturbance
Agricultural runoff	Possible deterioration of habitat and water quality
Tourism	Disturbance to birds
Vehicular movement	Noise and physical disturbance
Habitat encroachment	Loss of wetland habitat
Water extraction	Change in water availability
Solid-waste disposal	Habitat degradation

The synopsis reports that these disturbances were intended to be observed during field visits and related to bird occurrence where possible.

Because the original field measurements are not available, the present paper does not claim that any one of these activities caused a measured decline in bird abundance. They are discussed as reported pressures that may affect wetland habitats.

vi. Conservation significance

The diversity of birds recorded from Ujani Reservoir shows the importance of maintaining suitable wetland habitats. Conservation should include not only the open-water area but also shallow-water zones, mudflats, marshes, islands and other areas used by birds.

Migratory birds need suitable feeding and resting areas during their seasonal movements. Disturbance at important sites may reduce their suitability. Identification of important bird areas and control of unnecessary disturbance should therefore be part of wetland management.

Regular bird surveys should be carried out along with observations of water level, aquatic vegetation, habitat condition and human activities. This would provide a more complete picture of changes in the wetland than bird observations alone.

The available synopsis also recommends long-term bird monitoring, protection of sensitive areas, control of sand mining and encroachment, water-quality and vegetation assessment, awareness among local communities and sustainable tourism.

Conclusion

Ujani Reservoir supports a diverse community of resident and migratory wetland birds. Published records of 81 aquatic bird species and 38 wader species show the importance of the reservoir and its associated wetlands.

Wetland birds can be useful indicators of habitat condition when they are monitored regularly. Changes in species composition, abundance, seasonal occurrence and habitat use can provide useful information about changes in the wetland.

The human pressures reported around Ujani Reservoir highlight the need for continued monitoring and conservation. Protecting important bird habitats, reducing disturbance and involving local communities can help maintain the ecological value of the reservoir.

References

- Bibby, C. J., Burgess, N. D., Hill, D. A., & Mustoe, S. H. (2000). *Bird census techniques* (2nd ed.). Academic Press.
- Bharucha, E., & Gogte, P. (1990). Avian profile of a man-modified aquatic ecosystem in the backwaters of the Ujjani Dam. *Journal of the Bombay Natural History Society*, 87(1), 73–90.
- BirdLife International. (2018). *State of the world's birds: Taking the pulse of the planet*. BirdLife International.
- Chavan, S., & Kumbhar, A. (2020). Breeding ecology of River Tern (*Sterna aurantia*) at Ujjani-Bhigwan Reservoir Bird Sanctuary near Pune-Solapur Highway, India. *Asian Journal of Conservation Biology*, 9(2), 352–358.
- Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z.-I., Knowler, D. J., Lévêque, C., Naiman, R. J., Prieur-Richard, A.-H., Soto, D., Stiassny, M. L. J., & Sullivan, C. A. (2006). Freshwater biodiversity: Importance, threats, status and conservation challenges. *Biological Reviews*, 81(2), 163–182.
- Gregory, R. D., van Strien, A., Vorisek, P., Gmelig Meyling, A. W., Noble, D. G., Foppen, R. P. B., & Gibbons, D. W. (2005). Developing indicators for European birds. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1454), 269–288.
- Gregory, R. D., & van Strien, A. (2010). Wild bird indicators: Using composite population trends of birds as measures of environmental health. *Ornithological Science*, 9, 3–22.
- Islam, M. Z., & Rahmani, A. R. (2004). *Important bird areas in India: Priority sites for conservation*. Indian Bird Conservation Network, Bombay Natural History Society, & BirdLife International.
- Islam, M. Z., & Rahmani, A. R. (2012). *Threatened birds of India: Their conservation status*. Indian Bird Conservation Network & Bombay Natural History Society.
- Karikar, S. P., Mali, S. V., Kulkarni, P., & Aphale, P. (2019). An assessment of bird communities across Ujjani and its five satellite wetlands in Solapur District of Maharashtra, India. *Journal of Threatened Taxa*, 11(15), 14989–14997. <https://doi.org/10.11609/jott.4741.11.15.14989-14997>
- Kumar, P., & Gupta, S. K. (2009). Diversity and abundance of wetland birds around Kurukshetra, India. *Our Nature*, 7, 212–217.

- Ma, Z., Cai, Y., Li, B., & Chen, J. (2010). Managing wetland habitats for waterbirds: An international perspective. *Wetlands*, 30, 15–27.
- Markert, B. A., Breure, A. M., & Zechmeister, H. G. (Eds.). (2003). *Bioindicators and biomonitoring: Principles, concepts and applications*. Elsevier.
- Mitsch, W. J., & Gosselink, J. G. (2009). *Wetlands* (4th ed.). John Wiley & Sons.
- Newton, I. (2008). *The migration ecology of birds*. Academic Press.
- Noss, R. F. (1990). Indicators for monitoring biodiversity: A hierarchical approach. *Conservation Biology*, 4(4), 355–364.
- Ramsar Convention Secretariat. (2021). *Global wetland outlook: Special edition 2021*. Ramsar Convention Secretariat.
- Sekercioglu, C. H. (2006). Increasing awareness of avian ecological function. *Trends in Ecology & Evolution*, 21(8), 464–471.
- Weller, M. W. (1999). *Wetland birds: Habitat resources and conservation implications*. Cambridge University Press.

