

OPERATION APPROACHES ON g_s - OPEN SETS IN TOPOLOGICAL SPACES

Jayashree R¹, Dr. Sivakamasundari K²

¹Department of Mathematics, Avinashilingam Institute for Home Science and Higher Education for Women Coimbatore, Tamilnadu, India.

²Professor, Department of Mathematics, Avinashilingam Institute for Home Science and Higher Education for Women Coimbatore, Tamilnadu, India.

Abstract: The concept of an operation κ on the family of g_s -open sets $GSO(X, \tau)$, in a topological space (X, τ) is introduced. The concepts of κ -open sets, κ -regular and κ -closed are introduced using the operation κ and their related topological properties are studied.

Key words: κ -interior, κ -open set, κ -regular set, κ -closed set, $g_s\kappa$ closure, g_s -closure $_{\kappa}$, $regular_{\kappa}$ operation and $open_{\kappa}$ operation.

1. Introduction

In 1963, Levine [4] introduced the concept of semi-open set. Following this the notion of generalized semi closed sets was introduced by Arya and Nour [1] in 1990.

Kasahara [3] defined the concept of an operation on a topological space and introduced the concept of a α -closed graphs of functions in 1979. Following his work, Jankovic [2], developed the concept of α -closed sets and investigated functions with α -closed graphs in 1983. Ogata [5] defined and investigated the concept of operation-open sets.

In this paper, we shall introduce operation κ on generalized semi open sets. Section 3 of this paper deals with the definition and properties of κ -interior, κ -open set, κ -regular set, κ -closed set, $g_s\kappa$ closure, g_s -closure $_{\kappa}$, $regular_{\kappa}$ operation and $open_{\kappa}$ operation.

2. Preliminary

Definition 2.1

Let (X, τ) be a topological space. A subset A of a space (X, τ) is called **generalized semi closed (g_s -closed)** set if $scl(A) \subseteq U$ whenever $A \subseteq U$ and U is semi-open in (X, τ) .

Definition 2.2

Let (X, τ) be a topological space. A subset A of a space (X, τ) is called **generalized semi open (g_s -open)** set if $X \setminus A$ is g_s -closed. The collection of all g_s -open sets is denoted by $GSO(X, \tau)$. Clearly $\tau \subseteq GSO(X, \tau)$.

Remark 2.3

Every closed set is g_s -closed but the converse not true.

Definition 2.4 [2]

Let (X, τ) be a topological space. An **operation** $\gamma: \tau \rightarrow P(X)$ is a mapping from τ into the power set of X such that $V \subseteq V^\gamma$ for each $V \in \tau$, where V^γ denotes the value of γ at V .

Definition 2.5 [5]

A subset A of a space (X, τ) will be called a **γ -open set** of (X, τ) if for each $x \in A$, there exists an open set U such that $x \in U$ and $U^\gamma \subseteq A$. τ_γ will denote the set of all γ -open sets. Clearly we have $\tau \supseteq \tau_\gamma$.

Definition 2.6 [5]

A subset B of (X, τ) is said to be **γ -closed** in (X, τ) if $X \setminus B$ is γ -open in (X, τ) .

Definition 2.7 [5]

A point $x \in X$ is in the **γ -closure** of a set $A \subseteq X$ if $U^\gamma \cap A \neq \emptyset$ for each open set U of x . The γ closure of a set A is denoted by $Cl_\gamma(A)$.

Definition 2.8[5]

An operation $\gamma: \tau \rightarrow P(X)$ is a mapping from τ into the power set $P(X)$.

$$\tau_\gamma\text{-}Cl(A) = \cap \{F: A \subseteq F, X \setminus F \in \tau_\gamma\}.$$

Where τ_γ denotes the set of all γ -open sets in (X, τ) .

3. κ - Open sets.

Definition 3.1

Let (X, τ) be a topological space. A mapping $\kappa: GSO(X, \tau) \rightarrow P(X)$ the family of generalized semi open sets $GSO(X, \tau)$ to the power set of X such that $V \subseteq V^\kappa$ for every $V \in GSO(X, \tau)$ where V^κ denotes the value of V under the operation κ .

Example 3.2

Let $X = \{a, b, c\}$ and $\tau = \{X, \phi, \{a\}, \{b\}, \{a, b\}, \{a, c\}\}$. Then $\kappa: GSO(X, \tau) \rightarrow P(X)$ defined by $A^\kappa = \begin{cases} A & \text{if } b \in A \\ cl(A) & \text{if } b \notin A \end{cases}$ is a κ -operation on (X, τ) as $A \subseteq A^\kappa$, for every $A \in GSO(X, \tau) = \{X, \phi, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}\}$.

Definition 3.3

A subset A of a space (X, τ) will be called a κ -open set of (X, τ) if for each $x \in A$, there exists a gs -open neighbourhood U of x and $U^\kappa \subseteq A$.

$\kappa O(X, \tau)$ will denote the set of all κ -open sets.

Example 3.4

Let (X, τ) and κ be defined as in Example(3.2). Then the κ -open sets are $\{X, \phi, \{b\}, \{a, b\}, \{b, c\}\}$.

Theorem 3.5

If κ is an operation on $GSO(X, \tau)$, then the following results are true.

- (a) Every κ -open set of (X, τ) is gs -open in (X, τ) .
- (i.e) $\kappa O(X, \tau) \subseteq GSO(X, \tau)$.
- (b) Every γ -open set of (X, τ) is κ -open.
- (c) Arbitrary union of κ -open sets in (X, τ) is also κ -open set.

Proof

(a) Consider a κ -open set A in (X, τ) and a point $x \in A$. By Definition(3.3), there exists a gs -open neighbourhood B of x such that $B^\kappa \subseteq A$. By Definition(3.1), $B \subseteq B^\kappa$ and hence $x \in B \subseteq B^\kappa \subseteq A$. (i.e.) $x \in B \subseteq A$ implying that A is a gs -open set. Thus $\kappa O(X, \tau) \subseteq GSO(X, \tau)$.

(b) Consider a γ -open set C of (X, τ) with $x \in C$. By Definition(2.7), there exists an open set U such that $x \in U \subseteq U^\kappa \subseteq C$. Since every open set is gs -open, C is κ -open.

(c) Consider $\{B_\alpha: \alpha \in J\}$, a collection of κ -open sets in (X, τ) . Let $x \in B = \cup B_\alpha$. Hence $x \in B_\alpha$ for some α and since B_α is κ -open, there exists a gs -open neighborhood U of x such that $U^\kappa \subseteq B_\alpha \subseteq B$. Therefore B is κ -open.

Example 3.6

This example shows that a gs -open set need not be κ -open.

Let $X = \{a, b, c\}$, $\tau = \{x, \phi, \{a\}, \{a, b\}\}$ and $gs\text{-open} = \{x, \phi, \{a\}, \{b\}, \{a, b\}, \{a, c\}\}$. Let κ be an operation on $GSO(X, \tau)$ such that $\kappa(A) = \begin{cases} A & , \text{if } b \in A \\ \{a, c\} & , \text{if } b \notin A \end{cases}$.

Then the κ -open sets are $\{X, \phi, \{b\}, \{a, b\}, \{a, c\}\}$. Here $\{a\}$ is gs -open but not κ -open.

Example 3.7

This example shows that a κ -open set need not be γ -open.

Let $X = \{a, b, c\}$, $\tau = \{x, \phi, \{a\}, \{b\}, \{a, b\}\}$ and $gs\text{-open} = \{x, \phi, \{a\}, \{b\}, \{a, b\}, \{a, c\}\}$. Let κ be an operation on $GSO(X, \tau)$ such that $\kappa(A) = gcl(A)$. Then the κ -open sets are $\{X, \phi, \{b\}, \{a, c\}\}$. And γ be an operation on τ . The γ -open sets are $\{X, \phi, \{b\}\}$. Here $\{a, c\}$ is κ -open but not γ -open.

Remark 3.8

Intersection of any two κ -open sets need not be κ -open.

Counter Example 3.9

The following example shows that intersection of κ -open sets need not be κ -open.

Let $X = \{a, b, c\}$, $\tau = \{x, \phi, \{a\}, \{a, b\}\}$ and $GSO(X, \tau) = \{x, \phi, \{a\}, \{b\}, \{a, b\}, \{a, c\}\}$. Let κ be an operation on $GSO(X, \tau)$ such that $\kappa(A) = \begin{cases} A & , \text{if } b \in A \\ \{a, c\} & , \text{if } b \notin A \end{cases}$.

Then the κ -open sets are $\{X, \phi, \{b\}, \{a, b\}, \{a, c\}\}$.

Intersection of the κ -open sets $\{a, b\}$ and $\{a, c\}$ is not a κ -open.

Definition 3.10

A κ -operation $\kappa: GSO(X, \tau) \rightarrow P(X)$ is called **regular κ -operation** given $x \in X$ and for each pair of gs -open neighbourhoods A and B of x , there exists a gs -open neighbourhood C of x such that $A^\kappa \cap B^\kappa \supseteq C^\kappa$.

Definition 3.11

A topological space (X, τ) is called **κ -regular** if for given $x \in X$ and each gs -open neighbourhood U of x , there exists a gs -open neighbourhood V of x such that $V^\kappa \subseteq U$.

Theorem 3.12

Let (X, τ) be a topological space and κ an operation on $GSO(X, \tau)$. Then the following results are equivalent.

- (a) $GSO(X, \tau) = \kappa O(X, \tau)$.
- (b) (X, τ) is a κ -regular space.
- (c) Given $x \in X$ and every gs -open set B of (X, τ) containing x there exists a κ -open set W of (X, τ) such that $x \in W$ and $W \subseteq B$.

Proof

(a) ⇒ (b)

Let x in X and V , a gs -open neighbourhood of x . By (a), V is κ -open in (X, τ) . By Definition(3.3), there exists a gs -open neighbourhood U of x such that $U^\kappa \subseteq V$. Hence by Definition(3.11), (X, τ) is κ -regular.

(b) ⇒ (c)

Consider $x \in X$ and a gs -open neighbourhood B of x . By(b), (X, τ) is a κ -regular space. Hence by Definition (3.11), there exists a gs -open neighbourhood W of x such that $W^\kappa \subseteq B$. By Definition(3.1), $W \subseteq W^\kappa$. Hence $x \in W \subseteq W^\kappa \subseteq B$.

Claim: W is κ -open.

Let $y \in W$. Implies

$y \in X$ and W , be the gs -open neighbourhood of y . Then By (b), there exists a gs -open neighbourhood U of x such that $U^\kappa \subseteq W$. By Definition(3.3), W is κ -open.

Hence, there

exists a κ -open set W such that $x \in W \subseteq B$, proving (c).

(c) ⇒ (a)

In Theorem(3.5)(i), it is proved that $\kappa O(X, \tau) \subseteq GO(X, \tau)$. It is left to prove $GSO(X, \tau) \subseteq \kappa O(X, \tau)$.

Let A be a gs -open set in (X, τ) and $x \in A$. Then $x \in X$ and By (c), there exists a

κ -open set W of (X, τ) such that $x \in W \subseteq A$.

(1) Since W is a κ -open set there exists

a gs -open set V such that $x \in V^\kappa \subseteq W$. (2)

(1) and (2) implies $x \in V^\kappa \subseteq A$. Implies A is κ -open.

Therefore, $GSO(X, \tau) \subseteq \kappa O(X, \tau)$.

Hence, $GSO(X, \tau) = \kappa O(X, \tau)$.

In general the intersection of two κ -open sets is not κ -open. The following theorem proves that if κ is a regular κ -operation then the intersection of κ -open sets is κ -open.

Theorem 3.13

Let κ be a regular κ -operation on $GSO(X, \tau)$. If A and B are κ -open sets in (X, τ) then $A \cap B$ is κ -open.

Proof

Let A and B be κ -open sets in (X, τ) . Consider $C = A \cap B$. Let $x \in C$ implies $x \in A$ and $x \in B$. Since A and B are κ -open sets, there exists a gs -open neighbourhoods U and V of x such that $U^\kappa \subseteq A$ and $V^\kappa \subseteq B$. Since the operation κ is regular κ , by Definition (3.10), there exists a gs -open neighbourhood C of x such that $C^\kappa \subseteq U^\kappa \cap V^\kappa \subseteq A \cap B$. Therefore $A \cap B$ is κ -open.

Remark 3.14

$\kappa O(X, \tau)$ forms a topology whenever κ is regular κ .

Definition 3.15

A subset A of a topological space (X, τ) is called κ -closed whenever $X - A$ is κ -open.

Example 3.16

From the Example(3.2) and Example(3.4), the κ -closed sets are $\{X, \phi, \{a\}, \{c\}, \{a, c\}\}$.

Definition 3.17

Let κ be an operation on $GSO(X, \tau)$. A point $x \in X$ is said to be a

κ -closure point of the set A if $U^\kappa \cap A \neq \phi$ for each gs -open neighbourhood U of x .

$$gs Cl_\kappa(A) = \{x \in X / U^\kappa \cap A \neq \phi, \forall U, gs\text{-open neighborhood of } x\}$$

open neighborhood of x

Example 3.18

Let X, τ and κ be defined as in Example(3.2). Let $A = \{b, c\}$ then $Cl_\kappa(A) = \{a, b, c\} = X$.

Remark 3.19

Let κ be an operation on $GSO(X, \tau)$. Then $gs Cl(A) \subseteq gs Cl_\kappa(A)$.

Proof

Let $x \in gs cl(A)$. Implies $A \cap V \neq \phi$, for every gs -open neighbourhood V of x . Now, $V \subseteq V^\kappa$ implies $A \cap V^\kappa \neq \phi$. By Definition (3.17), $x \in Cl_\kappa(A)$. Hence, $gs Cl(A) \subseteq gs Cl_\kappa(A)$.

Definition 3.20

Let κ be an operation on $GSO(X, \tau)$. Then $gs_\kappa Cl(A)$ is defined as the intersection of all κ -closed sets containing A .

$$gs_\kappa Cl(A) = \cap \{F \subseteq X / A \subseteq F \text{ and } X \setminus F \in \kappa O(X, \tau)\}$$

Theorem 3.21

Let (X, τ) be a topological space and A a subset of X and κ be an operation on $GSO(X, \tau)$. Then for a given $y \in X$, $y \in gs_\kappa Cl(A)$ if and only if $V \cap A \neq \phi$ for every $V \in \kappa O(X, \tau)$ such that $y \in V$.

Proof

Define: $F = \{y \in X / V \cap A \neq \phi \text{ for every } V \in \kappa O(X, \tau) \text{ and } y \in V\}$. It is to be proved that $gs_\kappa Cl(A) = F$.

Take $x \notin F$. By the construction of F , there exists a κ -open set V containing x such that $V \cap A = \phi$. Then $X \setminus V$ is κ -closed and $A \subseteq X \setminus V$. Taking $gs_\kappa Cl(A)$ on both sides, $gs_\kappa Cl(A) \subseteq gs_\kappa Cl(X \setminus V) = X \setminus V$. Since $x \in V$, $x \notin X \setminus V$ implies $x \notin gs_\kappa Cl(A)$. Hence $gs_\kappa Cl(A) \subseteq F$.

Take $x \notin gs_\kappa Cl(A) = \cap \{E / A \subseteq E \text{ and } X \setminus E \in \kappa O(X, \tau)\}$. Then there exists κ -closed set E such that $A \subseteq E$, but $x \notin E$ implies $x \in X \setminus E \in \kappa O(X, \tau)$ and $(X \setminus E) \cap A = \phi$ implies $x \notin E$. Therefore $E \subseteq gs_\kappa Cl(A)$. Hence $gs_\kappa Cl(A) = E$.

Theorem 3.22

Let (X, τ) be a topological space. Let A and B be subsets of X and κ be an operation on $GSO(X, \tau)$. The statements below are true.

- (a) The set $gs_{\kappa}Cl(A)$ is κ -closed and $A \subseteq gs_{\kappa}Cl(A)$.
- (b) A is κ -closed if and only if $A = gs_{\kappa}Cl(A)$.
- (c) If $A \subseteq B$ then $gs_{\kappa}Cl(A) \subseteq gs_{\kappa}Cl(B)$.
- (d) $gs_{\kappa}Cl(A) \cup gs_{\kappa}Cl(B) \subseteq gs_{\kappa}Cl(A \cup B)$.
- (e) If κ is *regular* $_{\kappa}$, then $gs_{\kappa}Cl(A) \cup gs_{\kappa}Cl(B) = gs_{\kappa}Cl(A \cup B)$.
- (f) $gs_{\kappa}Cl(A \cap B) \subseteq gs_{\kappa}Cl(A) \cap gs_{\kappa}Cl(B)$.
- (g) $gs_{\kappa}Cl(gs_{\kappa}Cl(A)) = gs_{\kappa}Cl(A)$.

Proof

(a) Let $A \subseteq X$. Consider $gs_{\kappa}Cl(A) = B$, say. Claim: B is κ -closed.
 prove: $X - B$ is κ -open. Now $B = gs_{\kappa}Cl(A) = \cap \{ \text{all } \kappa\text{-closed sets containing } A \}$.

$X \setminus B = B^c = (gs_{\kappa}Cl(A))^c = \cup \{ \text{all } \kappa\text{-open sets contained in } A \} = \text{a } \kappa\text{-open set (by Theorem (3.5) (c)).}$

$A \subseteq gs_{\kappa}Cl(A)$ follows directly from the Definition(3.20).

(b) A is κ -closed if and only if $A = gs_{\kappa}Cl(A)$

Necessity: Since A is κ -closed, $gs_{\kappa}Cl(A) = \cap \{ \text{all } \kappa\text{-closed sets containing } A \} = A$

Sufficiency: Since $A = gs_{\kappa}Cl(A)$ and from (a) $gs_{\kappa}Cl(A)$ is κ -closed. We get A is κ -closed.

(c) Let $A \subseteq B$.

$gs_{\kappa}Cl(A) = \cap \{ \text{all } \kappa\text{-closed sets containing } A \} = \cap \mathcal{A}$ where \mathcal{A} is the collection of all κ -closed sets containing A

$gs_{\kappa}Cl(B) = \cap \{ \text{all } \kappa\text{-closed sets containing } B \} = \cap \mathcal{B}$ where \mathcal{B} is the collection of all κ -closed sets containing B

Since $\mathcal{A} \subseteq \mathcal{B}$, $(\cap \mathcal{A}) \subseteq (\cap \mathcal{B})$. Therefore $gs_{\kappa}Cl(A) \subseteq gs_{\kappa}Cl(B)$.

(d) Let $A \subseteq A \cup B$ and $B \subseteq A \cup B$. Therefore By (c) $gs_{\kappa}Cl(A) \subseteq gs_{\kappa}Cl(A \cup B)$ and $gs_{\kappa}Cl(B) \subseteq gs_{\kappa}Cl(A \cup B)$. Hence $gs_{\kappa}Cl(A) \cup gs_{\kappa}Cl(B) \subseteq gs_{\kappa}Cl(A \cup B)$.

(e) Suppose κ is *regular* $_{\kappa}$. Let $y \notin gs_{\kappa}Cl(A) \cup gs_{\kappa}Cl(B)$. Then $y \notin gs_{\kappa}Cl(A)$ and $y \notin gs_{\kappa}Cl(B)$. Then there exist two κ -open sets U and V such that $U \cap A = \phi$ and $V \cap B = \phi$. Since κ is *aregular* $_{\kappa}$ operation, by Theorem(3.11), $A \cap B$ is κ -open in (X, τ) . Therefore $(U \cap A) \cap (V \cap B) = (U \cap V) \cap (A \cap B) = \phi$. Implies $x \notin gs_{\kappa}Cl(A \cup B)$.

Hence, $gs_{\kappa}Cl(A \cup B) \subseteq gs_{\kappa}Cl(A) \cup gs_{\kappa}Cl(B)$.

(f) We know $A \cap B \subseteq A$ and $A \cap B \subseteq B$. Hence by (c), $gs_{\kappa}Cl(A \cap B) \subseteq gs_{\kappa}Cl(A)$
 and $gs_{\kappa}Cl(A \cap B) \subseteq gs_{\kappa}Cl(B)$. Therefore $gs_{\kappa}Cl(A \cap B) \subseteq gs_{\kappa}Cl(A) \cap gs_{\kappa}Cl(B)$.

(g) From (a), $gs_{\kappa}Cl(A) \subseteq gs_{\kappa}Cl(gs_{\kappa}Cl(A))$. Now by Theorem(3.19) for every κ -open set containing z in (X, τ) , $V \cap gs_{\kappa}Cl(A) \neq \phi$. Therefore there exist a point $y \in V$ and $y \in gs_{\kappa}Cl(A)$. Again by Theorem (3.19), $V \cap A \neq \phi$. Which implies $z \in gs_{\kappa}Cl(A)$. Hence $gs_{\kappa}Cl(gs_{\kappa}Cl(A)) = gs_{\kappa}Cl(A)$.

Definition 3.23

An operation κ on $GSO(X, \tau)$ is said to be **open** $_{\kappa}$ operation if for every gs -open neighbourhood U of $x \in X$, there exists a κ -open set V such that $x \in V$ and $V \subseteq U^{\kappa}$

Theorem 3.24

Let $\kappa: GSO(X, \tau) \rightarrow P(X)$ be an operation on $GSO(X, \tau)$ and P and Q are subsets of X . Then the results below are true.

- (a) $gsCl_{\kappa}(A)$ is gs -closed and $A \subseteq gsCl_{\kappa}(A)$.
- (b) A is κ -closed if and only if $A = gsCl_{\kappa}(A)$.
- (c) If (X, τ) is a κ -regular space then $gsCl_{\kappa}(A) = gscl(A)$.
- (d) If $A \subseteq B$ then $gsCl_{\kappa}(A) \subseteq gsCl_{\kappa}(B)$.
- (e) $gsCl_{\kappa}(A) \cup gsCl_{\kappa}(B) \subseteq gsCl_{\kappa}(A \cup B)$.
- (f) If κ is *regular* $_{\kappa}$, then $gsCl_{\kappa}(A) \cup gsCl_{\kappa}(B) = gsCl_{\kappa}(A \cup B)$.
- (g) $gsCl_{\kappa}(A \cap B) \subseteq gsCl_{\kappa}(A) \cap gsCl_{\kappa}(B)$.
- (h) If κ is *open* $_{\kappa}$ operation then $gsCl_{\kappa}(A) = gs_{\kappa}Cl(A)$ and $gsCl_{\kappa}(gsCl_{\kappa}(A)) = gsCl_{\kappa}(A)$.

Proof

(a) Let $A \subseteq X$. Consider $x \in gscl(Cl_{\kappa}(A))$. Then for every gs -open neighbourhood V of x , $V \cap gsCl_{\kappa}(A) \neq \phi$. Let $y \in V \cap gsCl_{\kappa}(A)$. Since $V \subseteq V^{\kappa}$, $V^{\kappa} \cap gsCl_{\kappa}(A) \neq \phi$. Implies $x \in gsCl_{\kappa}(A)$. Therefore,
 $gscl(gsCl_{\kappa}(A)) \subseteq gsCl_{\kappa}(A)$. Hence $gsCl_{\kappa}(A)$ is gs -closed.

Let $x \in A$ and U be any gs -open neighbourhood of x , then $x \in U \cap A$. Now $U \subseteq U^{\kappa}$ implies $x \in U^{\kappa} \cap A$ which $U^{\kappa} \cap A = \phi$. Therefore, $x \in gsCl_{\kappa}(A)$. Hence, $A \subseteq gsCl_{\kappa}(A)$.

(b) **Necessity:** Let A be κ -closed. Then $X \setminus A$ is κ -open. Claim: $gsCl_{\kappa}(A) \subseteq A$. Let $x \notin A$. Then $x \in X \setminus A$. Since $X \setminus A$ is a κ -open set containing x , by Definitin(3.3), there exists a gs -open set containing x such that $U^{\kappa} \subseteq X \setminus A$ which implies $U^{\kappa} \cap A = \phi$. Therefore $x \notin gsCl_{\kappa}(A)$. Hence $gsCl_{\kappa}(A) \subseteq A$.

Sufficiency: Let $A = gsCl_{\kappa}(A)$. Let $x \in X \setminus A$. Then $x \notin A = gsCl_{\kappa}(A)$, there exists a gs -open neighbourhood W of x such that $W^{\kappa} \cap A = \phi$ which implies $W^{\kappa} \subseteq X \setminus A$. By Definition(3.3), $X \setminus A$ is κ -open (i.e.) A is κ -closed.

(c) $gscl(A) \subseteq gsCl_{\kappa}(A)$ is proved in Remark(3.17). Let $x \notin gscl(A)$. Then there exists a gs -open neighbourhood U of x such that $U \cap A = \phi$. Since (X, τ) is a κ -regular space by Definition(3.11), for every $x \in X$, there exists a gs -neighbourhood V of x such that $V^{\kappa} \subseteq U$ and so $V^{\kappa} \cap A = \phi$ which implies $x \notin gsCl_{\kappa}(A)$. Therefore, $gsCl_{\kappa}(A) \subseteq gscl(A)$.

(d) Given $A \subseteq B$. Let $x \in gsCl_{\kappa}(A)$. By Definition(3.17), there exists a gs -open neighbourhood U of x such that $U^{\kappa} \cap A \neq \phi$ which implies $V^{\kappa} \cap B \neq \phi$. Therefore $x \in gsCl_{\kappa}(B)$. Hence $gsCl_{\kappa}(A) \subseteq gsCl_{\kappa}(B)$.

(e) $A \subseteq (A \cup B)$ implies $gsCl_{\kappa}(A) \subseteq gsCl_{\kappa}(A \cup B)$ From(d). Similarly $B \subseteq (A \cup B)$ implies $gsCl_{\kappa}(B) \subseteq gsCl_{\kappa}(A \cup B)$.

Hence, $gsCl_{\kappa}(A) \cup gsCl_{\kappa}(B) \subseteq gsCl_{\kappa}(A \cup B)$.

(f) Let κ is *aregular* $_{\kappa}$ operation. Let $x \notin gsCl_{\kappa}(A) \cup gsCl_{\kappa}(B)$. $x \notin gsCl_{\kappa}(A)$ and $x \notin gsCl_{\kappa}(B)$. There exists *gs*-open neighbourhoods U and V of x such that $U^{\kappa} \cap A = \phi$ and $V^{\kappa} \cap B = \phi$. Since κ is *aregular* $_{\kappa}$ by Definition (3.10), there exists a *gs*-open neighbourhood C of x such that $C^{\kappa} \subseteq U^{\kappa} \cap V^{\kappa}$. Which implies $C^{\kappa} \cap (A \cup B) \subseteq (U^{\kappa} \cap V^{\kappa}) \cap (A \cup B) = \phi$ and $x \notin gsCl_{\kappa}(A \cup B)$. Therefore $gsCl_{\kappa}(A \cup B) \subseteq gsCl_{\kappa}(A) \cup gsCl_{\kappa}(B)$.

Hence $gsCl_{\kappa}(A) \cup gsCl_{\kappa}(B) = gsCl_{\kappa}(A \cup B)$.

(g) $A \cap B \subseteq A$ implies $gsCl_{\kappa}(A \cap B) \subseteq gsCl_{\kappa}(A)$ and $A \cap B \subseteq B$ implies $gsCl_{\kappa}(A \cap B) \subseteq gsCl_{\kappa}(B)$ Therefore, $gsCl_{\kappa}(A \cap B) \subseteq gsCl_{\kappa}(A) \cap gsCl_{\kappa}(B)$.

(h) Let $y \notin gsCl_{\kappa}(A)$, there exists a *gs*-open neighbourhood U of y such that $(U^{\kappa} \cap A) = \phi$. Since the operation κ is *open* $_{\kappa}$, there exists a κ -open set V containing y such that $V \subseteq U^{\kappa}$ implies $V \cap A = \phi$ and $x \notin gs_{\kappa}Cl(A)$ by Theorem(3.19). Hence $gs_{\kappa}Cl(A) \subseteq gsCl_{\kappa}(A)$.

Let $x \in gsCl_{\kappa}(A)$. Suppose $x \notin F$ where F is κ -closed and $A \subseteq F$. Then $x \in X \setminus F$, and $A \cap (X \setminus F) = \phi$. Since $X \setminus F$ is κ -open and $x \in X \setminus F$, there exists *ags*-open set W such that $x \in W$ and $W^{\kappa} \subseteq X \setminus F$. Which implies $A \cap W^{\kappa} = \phi$. Thus $x \notin gsCl_{\kappa}(A)$.

Hence, if the operation κ is *open* $_{\kappa}$ operation then $gsCl_{\kappa}(A) = gs_{\kappa}Cl(A)$.

References

- [1] Arya, S.P. and Nour, T.M., Characterizations of S-normal spaces, Indian J. Pure. Appl. Math., (1990), 21,717-719.
- [2] Jankovic, D.S., On functions with α -closed graphs, Glasnic Mat., (1983), 18, 161-169.
- [3] Kasahara, S., Operation-compact spaces, Math. Japon., (24) 1979, 97-105.
- [4] Levine, N., Semi-open sets and semi-continuity in topological spaces, Amer. Math. Monthly., (1960), 67, 269-275.
- [5] Ogata, H., Operations on topological spaces and associated topology, Math. Japon., (1991), 36, 175-184.
- [6] Sanjay Tahiliani., Operation approach to β -open sets and applications, Math. Commun., (2011), 16, 577-591

