

Evaluation of mungbean (*Vigna radiata* L.) variety performance by growing in different nutrients and sowing method

Jaskirat Singh, A.H. Madakemohekar* and Ranjit Singh

School of Agriculture, Lovely Professional University, Phagwara-144 411 (Punjab)

Abstract

A field experiment was conducted at Agricultural Research Farm, School of Agriculture, Lovely Professional University, Phagwara, Punjab, during Rabi 2017-2018. The soil texture was sandy loam, with pH 7.83 to 7.98 and 0.4dSm^{-1} . The study was conducted to evaluate the performance of mung bean, by growing in different nutrients and sowing methods. In this experiment cultivar used was SML 668. The experiment was conducted using Two Factorial Random Block Design with three sowing patterns (flat sowing, raised bed sowing, ridge sowing) and 6 treatments T₀ (control), T₁ (Compost), T₂ (NPK), T₃ (Micronutrients), T₄ (NPK + Micronutrients), T₅ (NPK + Micronutrients + Compost) and were replicated three times. It was found that Ridge sowing shows. T₅ showed better results in growth and yield parameters (NPK + Micronutrients + Compos). However ridge sowing and T₅ interaction performed better results in higher plant height, number of leaves, number of branches and grain yield whereas interaction between flat sowing and T₅ sowing shows better results in pod length, number of grains per pod, pods per plant and 1000 grains weight.

Key words: NPK, micronutrients, compost, grain weight, yield, fertilizer

Introduction

Mung bean (*Vigna radiate* L.) is one of important kharif as well as summer, short duration crop, grown mainly for pulse. It fixes 200 to 300 kg/ha/year, nitrogen through symbiotic relation (Geletu and Mekonnen, 2018) and fulfill plant requirement for available nitrogen (Mahmood and Athar, 2008; Mandal *et al.*, 2009). Because of good palatability, easy digestibility, short term of crop and high cost of market, mung bean is a good choice for agriculturists (Choudhary, 2010). Many factors are responsible for the low yield of mung bean at farmer's field which include factors such as unawareness of farmers about optimum date of sowing, improper proper planting patterns, insufficient weed control, insect and pest management practices and imbalanced use of fertilizers. Among these factors, proper sowing time

and planting pattern are of great importance. Planting pattern influences radiation interception and utilization of moisture from soil (Malik *et al.*, 2003). Application of N, P, K, S and Mg, etc. and inorganic fertilizer in the form of compost in balanced form is essential for major processes like plant development and yield. Among other agronomic practices, proper method of planting may considerably increase the production of green gram. Ideal planting geometry is precious and important for better and efficient utilization of available plant growth resources in order to get maximum productivity (Sanjay *et al.*, 2014). Foliar use of supplements alongside soil application has great benefit for providing nutritional requirements of crops such as rapid and productive response by the plants, less product needed and independence of soil conditions. Foliar nutrition designed to mitigate the problems like fixation and immobilization of nutrients. This method provides utilization of nutrients more effectively and for overcoming deficiencies rapidly especially for short duration crops (Shruthi, 2013). The foliar application is one of the best methods because plants can fully utilize fertilizers, and these cause less environmental pollution (Das, 1999; Barik *et al.*, 1994). Vermicompost is a sustainable bio-fertilizer made from the organic wastes with the help of earthworms. Vermicompost is a rich source of N, P, K and micronutrients. It is also rich in growth hormones, vitamins and acts as powerful biocide against diseases and nematodes besides improving physical condition of soil. The application of vermicompost at increasing rates significantly increased nutrient content and uptake of N, P and K in grain and straw of green gram crop (Anita, 2017).

Materials and Methods

Experiment location

The research experiment was conducted at Agricultural Research Farm, School of Agriculture, Lovely Professional University, Phagwara, Punjab during Rabi season of the year 2017-2018, August to December. Geographically, farm is situated 31.25°N latitude and 75°E longitude with an altitude of 232 meter above the mean sea level. The soil of experimental field was found to be sandy clay loam and pH of the soil varied from 7.83 to 7.98. , EC (0.4dSm⁻¹), available phosphorus (248 kg/ ha), available nitrogen (235 Kg /ha). Geographically, farm is situated 31.25°N latitude and 75°E longitude with an altitude of 232 meter above the mean sea level. It has annual rainfall of 1919.5mm, subtropical with

cool winters, hot summers and a distinct rainy season. Supplemental irrigation to the field provided through canal water as per crop water demand.

Experimental design and treatments

The experimental design was laid out on two factorial Randomized Block Design with 3 sowing methods viz., S₁ (flat sowing), S₂ (raised bed sowing), S₃ (ridge sowing) and 6 treatments viz., T₀ (control), T₁ (NPK), T₂ (compost), T₃ (micronutrients), T₄ (NPK + micronutrients), T₅ (NPK + micronutrients + compost) with three replications. The Blocks were divided into 3 blocks to represent 3 replications. Irrigation and all other agronomic practices were carried out uniformly throughout the growing season. Application of NPK and micronutrients was done by foliar spraying.

Statistical analysis

The data was recorded for various growth, yield and quality attributes and then analyzed variance as described by Cochran and Cox (1987) for two factorial Randomized Block Design (RBD).

Results and discussion:

Plant height: It was observed that at harvest maximum plant height was recorded for treatment T₅ (NPK + micronutrients + compost) i.e. and S₃ (ridge method) i.e. 79.20 cm (Table 2a), because of proper aeration and availability of nutrient with organic matter. **Wasma et al., (2016)** reported that application of inorganic fertilizer and compost has beneficial effect on the growth of mung bean. These findings are in agreement with this study that on application of NPK+ micronutrients + compost shows a positive impact on the height of mung bean as well as obtaining good grain yield after combined application of fertilizers and micronutrients when grown on ridges. It was observed that plant height was found to be maximum in S₃ ridges sowing i.e. 74.64. (Table 1)

Number of leaves per plant: It was observed that number of leaves was found to be maximum in S₃ (ridge sowing) i.e. 54.68 (Table 1). Among interaction the maximum number of leaves were found maximum for T₅ (NPK + micronutrients + compost) and S₃ (ridge method) i.e. 59.20 (Table 2a). Same result was observed for number of leaves per plant and number of branches and these findings are in agreement with **Romel et al., (2014)**.

Number of branches per plant: It was observed that number of branches was found to be maximum in ridge sowing i.e. 26.40 T₅ (NPK + micronutrients + compost) has a maximum number of branches record

of 27.54 (Table 1). Among interaction of branches were found maximum for T₅ (NPK + micronutrients + compost) and S₃ (ridge method) i.e. 79.20 (Table 2a). **Delbar et al., (2014)** concluded that foliar application of micronutrients results in increase in plant height, number of leaves. It has been observed that application of T₅ (NPK + Micronutrients + Compost) has attained maximum number of leaves.

Number of grains per pod: The number of grains per pod is very important character which affect on the yield per plant. More number of grains per pod was found maximum in S₁ (flat sowing) because germination was less and plant showed more number of branches and healthiness. For fertilizer treatment T₅ (NPK + micronutrients + compost) i.e. 12.16 (Table 1), it was observed effective response on mung bean. Whereas, interaction between sowing method and different treatment combinations were found to be maximally higher among the combination T₅ (NPK + micronutrients + compost) and S₁ (flat sowing). These present findings can be further supported by findings of **Hammad et al., (2014)**.

Pod length: In the study conducted by **Mondal et al., (2010)** found that foliar application on micronutrients at reproductive stage appears effective in improving pod length of mung bean which is comparable to this current study in which pod length was found to be maximum in application of T₅ (NPK + Micronutrients + Compost) i.e. 10.70 cm. and in sowing method it was found maximum for flat sowing (Table 1). For interaction pod length was maximum in combination T₅ (NPK + micronutrients + compost) and S₁ (flat sowing) 12.00 (Table 2a). **Dilip et al. (2016)** observed significant increase in pod length after application of micronutrients.

Number of pods per plant: The combined application of T₅ (NPK + micronutrients + compost) has resulted in highest number of pods per plant i.e. 27.55 (Table 2b). This finding is substantiated by the study done by **Das et al., (2015)** which showed that combined application of N:P:K (19:19:19) fertilizer spray increases the yield components of mung bean.

1000 grain weight: It was found that maximum grain weight was found in S₁ (flat sowing) i.e. 53.81 g (Table 1). Combination of T₅ (NPK + micronutrients + compost) and S₁ (flat sowing method) resulted in highest results i.e. 55.36 g (Table 2b).

Table 1 shows the effect of different treatment combinations and sowing methods on yield parameters. The different sowing methods along with different treatments showed significant influence on grain yield as well as straw yield. Higher yield per plot was observed in ridge sowing. Among the treatments T₅ (NPK + micronutrients + compost) showed better results as compared to other treatments. Among

interaction T₅ (NPK + micronutrients + compost) and S₃ (ridge method) resulted in better yield i.e. 330 g (Table 2b). Straw yield was observed maximum in sowing method for S₂ (raised bed) and in fertilizer treatment for T₅ (NPK + micronutrients + compost), whereas, interaction T₅ (NPK + micronutrients + compost) and S₃ (ridge sowing) found maximum for straw yield (Table 2b). **Netwal (2003)** reported the application of vermicompost and compost had significant increasing effect of straw yield.

Table 1. Influence of different sowing method and treatments

Traits Treatments	Plant height ⁻¹	Number of leaves ⁻¹	Number of branches ⁻¹	Number of grains per pod	Pod length (cm)	Number of Pods ⁻¹	1000 grains Weight (g)	Grain yield per plot (g)	Straw yield per plot	Biological yield per plot
T ₀	63.34	45.63	23.82	9.54	7.50	20.15	50.55	166.00	393.33	559.33
T ₁	69.01	49.53	24.16	9.77	8.61	22.03	52.14	220.55	553.33	773.88
T ₂	68.42	48.47	24.70	10.58	9.33	22.82	54.08	238.33	576.66	814.99
T ₃	70.25	51.87	25.17	11.10	9.54	22.90	54.00	283.33	610.33	893.66
T ₄	71.18	54.67	24.97	11.51	10.11	26.60	53.98	293.88	625.66	919.54
T ₅	71.57	56.87	27.54	12.16	10.70	27.55	53.90	299.77	643.33	943.1
Sowing Methods										
S ₁	63.77	47.87	24.30	11.34	10.67	25.78	53.81	219.33	455.55	674.88
S ₂	68.49	50.97	24.50	10.15	8.61	21.19	52.00	241.33	627.94	869.27
S ₃	74.64	54.69	26.40	10.83	8.61	24.06	53.72	290.27	617.83	908.1

S₁ : Flat sowing

S₂ : Raised bed sowing

S₃ : Ridge sowing

Compost

T₀ : Control

T₁ : Compost

T₂ : NPK

T₃ : Micronutrients

T₄ : NPK + Micronutrients

T₅ : NPK + Micronutrients +

Table 2a. Interaction of different sowing method and treatments

Traits Treat ment	Plant height ⁻¹ (cm)			Number of Leaves ⁻¹			Number of branches ⁻¹			Number of grains per pod			Pod length (cm)		
	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃
T ₀	59.53	63.77	66.73	36.90	50.00	50.00	23.17	23.80	24.50	9.66	9.00	9.96	8.30	7.00	7.20
T ₁	64.70	69.60	72.73	48.50	47.00	53.10	24.13	23.77	24.60	10.06	9.06	10.20	9.63	8.10	8.10
T ₂	61.67	68.40	75.20	46.00	46.40	53.00	22.60	24.20	27.30	11.20	10.23	10.30	11.20	8.30	8.50
T ₃	63.70	69.47	77.60	50.20	51.20	54.20	25.70	24.20	25.60	12.00	10.10	11.20	11.33	8.50	8.80
T ₄	65.60	68.77	79.20	50.20	55.20	58.60	24.37	24.33	26.20	12.00	11.16	11.36	11.53	9.70	9.10
T ₅	67.40	70.90	76.40	55.40	56.00	59.20	25.80	26.67	30.17	13.13	11.36	12.00	12.00	10.10	10.00

Table 2b. Interaction of different sowing method and treatments

Traits Treat ment	Number of Pods ⁻¹			1000 grains Weight (g)			Grain yield per plot (g)			Straw yield per plot (g)			Biological yield per plot (g)		
	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃
T ₀	22.16	18.29	20.00	51.06	49.26	51.33	151.66	163.00	183.33	325.00	435.00	420.00	476.66	598.00	603.33
T ₁	23.33	20.25	22.52	52.33	51.40	52.70	185.00	191.66	285.00	405.00	636.66	618.33	590.00	828.32	903.33
T ₂	24.43	21.34	22.69	54.46	53.33	54.46	200.00	216.66	298.33	455.00	645.00	630.00	655.00	861.66	928.33
T ₃	26.20	16.96	25.55	54.46	54.26	54.46	250.00	283.33	316.66	486.66	678.33	666.00	736.66	961.66	982.66
T ₄	28.36	25.13	26.32	55.20	52.20	54.56	260.00	293.33	328.33	520.00	684.33	672.66	780.00	977.66	1000.9
T ₅	30.23	25.16	27.27	55.36	51.53	54.80	269.33	300.00	330.00	541.66	688.33	700.00	810.99	988.33	1030.0

S₁ : Flat sowing
S₂ : Raised bed sowing
S₃ : Ridge sowing

T₀ : Control
T₁ : Compost
T₂ : NPK

T₃ : Micronutrients
T₄ : NPK + Micronutrients
T₅ : NPK + Micronutrients + Compost

Conclusion:

The study has evaluated that ridge sowing method showed higher growth parameters as well as yield and yield attributes in comparison to other sowing methods. The study has evaluated that foliar spraying of micronutrients and NPK and application of compost has shown a synergistic effect on the growth and yield of mung bean as compared to other treatments. The combination of compost with foliar application of micronutrients and NPK showed increase in plant height, number of leaves, branches per plant, pod length, pods per plant, grains per pods well as grain yield and straw yield. Therefore, it can be concluded that the combination of compost with application of micronutrients and NPK should be applied by the use of ridge sowing method to get good yield.

References:

Anita Todawat (2017). Effect of vermicompost and zinc on yield and quality of green gram (*Vigna radiata* L.) grown under loamy sand soil. *Annals of Agricultural Research*, 24(1): 203-204.

- Barik, A., Sunda, P. and Mukherjee, A. (1994). Influence of nitrogen, phosphorus and potash fertilization on growth, yield and oil of kharif groundnut. *Indian J. Agric. Sci.* 38: 105-111.
- Chaudhary, H.R. (2010). Integrated Nutrient Management in Mungbean (*Vigna radiata*). Swami Keshwanand Rajasthan Agricultural University, Bikaner.
- Cochran, W.G. and Cox, G.M. 1987. Experimental Designs. 2nd ed. Bombay, Asia, 610 p.
- Das, P. (1999). Plants Nutrients. In: Manures and Fertilizers. 2nd Edition ed. Kalyani Publishers, New Delhi, India.
- Delbar Chehel Amirani, Pourang Kasraei (2015). The effect of foliar application of microelements on phenological and physiological characteristics of Mungbean under drought stress. *International Journal of Agronomy and Agricultural Research*, 7: 01-08.
- Geletu, T. and Mekonnen, F. (2018). The effect of bio and inorganic fertilizer on yield, nutrient uptake and economics of mungbean (*Vigna radiata* L. Wilczek) Varieties in Ethiopia. *J. Agri. Res.*, 3(11): 1-11.
- Hammad Hafiz Mohkum, Saeed Ahmad Ashfaq, Farhad Wajid and Nasim Wajid (2014). Sources of nutrients influence mungbean crop under that environmental college of agriculture, Layyah sub campus Bahauddin Zakariya University, Multan, Pakistan. *Applied sciences and business economics*, 4: 4-48.
- Mahmood, A. and Athar, M. (2008). Cross inoculation studies: Response of *Vigna mungo* to inoculation with rhizobia from tree legumes growing under arid Environment. *Int. J. Environ. Sci. Technol.*, 5: 135-139.
- Malik, M.A., Saleem, M.F., Asghar, A. and Ijaz M. (2003). Effect of nitrogen and phosphorus application on growth yield and quality of mungbean (*Vigna radiata* L.). *Pak. J. Agri. Sci.*, 40:3-4.
- Mandal, S., Mandal, M. and Das, A. (2009). Stimulation of indoleacetic acid production in a Rhizobium isolate of *Vigna mungo* by root nodule phenolic acids. *Arch. Microbiol.*, 191: 389-393.
- Mondal, M.M.A., Rahman, M.A., Akter, M.B. and Fakir, M.S.A. (2010). Effect of foliar application of nitrogen and micronutrients on growth and yield in mungbean. Bangladesh Institute of Nuclear Agriculture (BINA), *Bangladesh. Legume Res.*, 34 (3): 166-171.
- Netwal, L.C. (2003). Effect of FYM and vermicompost on nutrient uptake and phosphorus application on growth, yield and quality of mungbean (*Vigna radiata* L.). M.Sc. (Ag.), Thesis, Rajasthan Agricultural University, Bikaner.
- Romel Biswash, Md., Wasal Rahman, Md., Maksudul Haque, Md., Mashuka Sharmin and Rajesh Barua (2014). Effect of Potassium and vermicompost on the growth, yield and nutrient contents of mungbean (Bari Mung 5). *Open Science Journal of Bioscience and Bioengineering*, 1: 33-39.
- Sanjay Yadav and Singh, B.N. (2014). Effect of irrigation schedules and planting methods on yield, attributes and economics of green gram (*Phaseolus radiata* L.) under rice – wheat – green gram cropping system. *Plant Archives*, 14(1): 521-523

- Sharma, S., Upadhyay, R.S. and Sharma, C.R. (2003). Response of various levels of nitrogen and phosphorus application on growth, physiological parameters and yield of (*Vigna radiata* L.) Wilczek under rainfed and mid hill conditions of Himachal Pradesh. *Indian Journal of Agricultural Research*, 37(1): 52-55.
- Shruthi M.K. (2013). Effect of foliar application of water soluble NPK fertilizer on growth, yield and quality of lima bean. Master of Science (Agriculture) in Agronomy Bengaluru.
- Wasma Armin, Kh. Ashraf-Uz-Zaman, Sheikh Shawkat Zamil , Mominul Haque Rabin, Asim Kumar Bhadra, Fahima Khatun. (2016). Combined effect of organic and inorganic fertilizers on the growth and yield of mungbean (Bari Mung 6). *International Journal of Scientific and Research Publications*, 6(7): 557-561.

